

FUNCTIONAL CALCULUS AND SPECTRAL ASYMPTOTICS FOR HYPOELLIPTIC OPERATORS ON HEISENBERG MANIFOLDS. I.

RAPHAËL PONGE

ABSTRACT. This paper is part of a series papers devoted to geometric and spectral theoretic applications of the hypoelliptic calculus on Heisenberg manifolds. More specifically, in this paper we make use of the Heisenberg calculus of Beals-Greiner and Taylor to analyze the spectral theory of hypoelliptic operators on Heisenberg manifolds. The main results of this paper include: (i) Obtaining complex powers of hypoelliptic operators as holomorphic families of Ψ_H DO's, which can be used to define a scale of weighted Sobolev spaces interpolating the weighted Sobolev spaces of Folland-Stein and providing us with sharp regularity estimates for hypoelliptic operators on Heisenberg manifolds; (ii) Criteria on the principal symbol of P to invert the heat operator $P + \partial_t$ and to derive the small time heat kernel asymptotics for P ; (iii) Weyl asymptotics for hypoelliptic operators which can be reformulated geometrically for the main geometric operators on CR and contact manifolds, that is, the Kohn Laplacian, the horizontal sublaplacian and its conformal powers, as well as the contact Laplacian. For dealing we cannot make use of the standard approach of Seeley, so we rely on a new approach based on the pseudodifferential approach representation of the heat kernel. This is especially suitable for dealing with positive hypoelliptic operators. We will deal with more general operator in a forthcoming paper using another new approach. The results of this paper will be used in another forthcoming paper dealing with an analogue for the Heisenberg calculus of the noncommutative geometry which, in particular, will allow us to make use in the Heisenberg setting of Connes' noncommutative geometry, including the operator theoretic framework for the local index formula of Connes-Moscovici.

1. INTRODUCTION

This paper is part of a series papers devoted to geometric and spectral theoretic applications of the hypoelliptic calculus on Heisenberg manifolds. Recall that a Heisenberg manifold (M, H) consists of a manifold M together with a distinguished hyperplane bundle $H \subset TM$. This definition covers many examples: Heisenberg group and its quotients by cocompact lattices, (codimension 1) foliations, CR and contact manifolds and the confoliations of [ET].

The name Heisenberg manifold comes from the fact that the relevant tangent structure for a Heisenberg manifold (M, H) is rather that of a bundle GM of two-step nilpotent Lie groups, whose Lie group structure is encoded by an intrinsic Levi form $\mathcal{L} : H \times H \rightarrow TM/H$ (see [BG], [Be], [EMM], [FS1], [Gro], [Po6], [Ro2]).

In this context the most natural operators include Hörmander's sum of squares, the Kohn Laplacian, the horizontal sublaplacian and its conformal powers, as well as the contact Laplacian (see Section 3 and the references therein for an overview of these operators). Although these operators may be hypoelliptic, they are definitely not elliptic. Therefore, the classical pseudodifferential calculus cannot be used efficiently to study these operators.

The relevant substitute to the standard pseudodifferential calculus is provided by the Heisenberg calculus of Beals-Greiner [BG] and Taylor [Tay] (see also [Bo], [CGGP], [Dy1], [Dy2], [EMM], [FS1], [RSt]). The idea in the Heisenberg calculus, which goes back to Elias Stein, is the following. Since the relevant notion of tangent structure for a Heisenberg manifold (M, H) is that of a bundle

2000 *Mathematics Subject Classification.* Primary 58J40, 58J50; Secondary 58J35, 32V10, 35H10, 53D10.

Key words and phrases. Heisenberg calculus, complex powers, heat equation, spectral asymptotics, analysis on CR and contact manifolds, hypoelliptic operators.

GM of 2-step nilpotent graded Lie groups, it stands for reason to construct a pseudodifferential calculus which at every point $x \in M$ is well modelled by the calculus of convolution operators on the nilpotent tangent group $G_x M$.

The result is a class of pseudodifferential operators, the $\Psi_H\text{DO}$'s, which are locally ΨDO 's of type $(\frac{1}{2}, \frac{1}{2})$, but unlike the latter possess a full symbolic calculus and makes sense on a general Heisenberg manifold. In particular, a $\Psi_H\text{DO}$ admits a parametrix in the Heisenberg calculus if, and only if, its principal symbol is invertible, and then the $\Psi_H\text{DO}$ is hypoelliptic with a loss of derivatives controlled by its order (see Section 4 for a more detailed overview of the Heisenberg calculus).

In [Po6] a tangent groupoid has been associated to any Heisenberg manifold (M, H) as the differentiable groupoid encoding a smooth deformation of $M \times M$ (see also [Va]). This is the analogue for Heisenberg manifolds of Connes' tangent groupoid of a manifold which plays a pivotal role in his proof in [Co] of the index theorem of Atiyah-Singer [AS]. The results of [Po6] have been used subsequently in [Po8] to give an intrinsic definition of the principal symbol for the Heisenberg calculus, which was not done in [BG] or in [Tay].

In this paper, and its sequel [Po10], we make use of the Heisenberg calculus to analyze the spectral theory of hypoelliptic operators on Heisenberg manifolds. The main results of this paper include:

- Obtaining complex powers of hypoelliptic operators as holomorphic families of $\Psi_H\text{DO}$'s, which can be used to define a scale of W_H^s , $s \in \mathbb{R}$, of weighted Sobolev spaces interpolating the weighted Sobolev spaces of Folland-Stein and providing us with sharp estimates for $\Psi_H\text{DO}$'s.
- Criteria on the principal symbol of P to invert the heat operator $P + \partial_t$ and to derive the small time heat kernel asymptotics for P .
- Weyl asymptotics for hypoelliptic operators which can be reformulated geometrically for the main geometric operators on CR and contact manifolds, that is, the Kohn Laplacian, the horizontal sublaplacian and its conformal powers, as well as the contact Laplacian.

These results will be important ingredients in [Po11] to construct an analogue for the Heisenberg calculus of the noncommutative residue trace of Wodzicki ([Wo1], [Wo2]) and Guillemin [Gu1] and to study the zeta and eta functions of hypoelliptic operators. In turn this has several geometric consequences. In particular, this allows us to make use of the framework of Connes' noncommutative geometry, including the operator theoretic framework for the local index formula of Connes-Moscovici [CM].

Let us also mention that the lack of microlocality of the Heisenberg calculus does not allow us to carry out in the Heisenberg setting the standard approach of Seeley [Se] (see also [Sh], [Gru]) to complex powers of elliptic operators. Instead, we rely on a new approach, based on the pseudodifferential representation of the heat in [BGS], which is quite suitable for dealing with positive differential operators. A similar approach has also been used by Mathai-Melrose-Singer [MMS] and Melrose [Me3] in the context of projective pseudodifferential operators on Azamaya bundles. In [Po10] we will use another approach to deal with $\Psi_H\text{DO}$'s which are not positive differential operators.

1.1. Holomorphic families of $\Psi_H\text{DO}$'s. Prior to dealing with complex powers of hypoelliptic operators we define holomorphic families of $\Psi_H\text{DO}$'s and check their main properties.

In a local Heisenberg chart $U \subset \mathbb{R}^{d+1}$ the definition of a holomorphic family of $\Psi_H\text{DO}$'s parametrized by an open $\Omega \subset \mathbb{C}$ is similar to that of the standard definition of a holomorphic family of ΨDO 's in [Wo1, 7.14] and [Gu2, p. 189] (see also [KV]). In particular, we allow the order of the family of ΨDO 's to vary analytically. Several properties of $\Psi_H\text{DO}$'s on U extend *mutatis standis* to the setting of holomorphic families of $\Psi_H\text{DO}$'s. In particular, concerning the product of $\Psi_H\text{DO}$'s we have:

Proposition 1.1. *For $j = 1, 2$ let $(P_{j,z})_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E})$ be a holomorphic family of Ψ_H DO's and supposed that $(P_{1,z})_{z \in \Omega}$ or $(P_{2,z})_{z \in \Omega}$ is uniformly properly supported with respect to z . Then the family of products $(P_{1,z}P_{2,z})_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's.*

There is, however, a difficulty when trying to extend the definition to arbitrary Heisenberg manifolds. More precisely, the proof of the invariance of the Heisenberg calculus by Heisenberg diffeomorphisms relies on a characterization of the distribution kernels of Ψ_H DO's by means of a suitable class of kernels $\mathcal{K}^*(U \times \mathbb{R}^{d+1}) = \sqcup_{m \in \mathbb{C}} \mathcal{K}^m(U \times \mathbb{R}^{d+1}) \subset \mathcal{D}'(U \times \mathbb{R}^{d+1})$. Each distribution $K \in \mathcal{K}^m(U \times \mathbb{R}^{d+1})$ admits an asymptotic expansion in the sense of distributions,

$$(1.1) \quad K \sim \sum_{j \geq 0} K_{m+j}, \quad K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1}),$$

where $\mathcal{K}_l(U \times \mathbb{R}^{d+1})$ consists of kernels which are smooth for $y \neq 0$ and homogeneous of degree l if $l \notin \mathbb{Z}$ and homogeneous of degree l up to logarithmic terms otherwise. In particular, the definition of $\mathcal{K}_l(U \times \mathbb{R}^{d+1})$ depends upon whether l is an integer or in not, which causes trouble for defining holomorphic families of kernels in $\mathcal{K}^*(U \times \mathbb{R}^{d+1})$ whose order may take integer values.

We resolve this issue by giving an alternative description of the class $\mathcal{K}^*(U \times \mathbb{R}^{d+1})$ in terms of what we call *almost homogeneous* kernels, as they are homogenous modulo smooth terms and, under the Fourier transform, they correspond to the almost homogeneous symbols considered in [BG].

Since the definition of an almost homogeneous symbol of degree l does not depend on whether l is an integer or not, there is no trouble anymore to define holomorphic families of almost homogeneous kernels. Therefore, we can make use of the characterization of $\mathcal{K}^*(U \times \mathbb{R}^{d+1})$ in terms of almost homogenous kernels to define holomorphic families with values in $\mathcal{K}^*(U \times \mathbb{R}^{d+1})$ (see Definition 5.24 for the precise definition).

We show that the distribution of kernel holomorphic families of Ψ_H DO's can be characterized in terms of holomorphic families with values in $\mathcal{K}^*(U \times \mathbb{R}^{d+1})$. As a consequence we can extend to the setting of holomorphic families of Ψ_H DO's the arguments in the proof in [BG] and [Po8] of the invariance by Heisenberg diffeomorphisms of the Heisenberg calculus, so that we get:

Proposition 1.2. *Let $\phi : U \rightarrow \tilde{U}$ be a change of Heisenberg chart and let $(\tilde{P}_{z \in \Omega})_{z \in \Omega}$ be a holomorphic family of Ψ_H DO's on $\tilde{U}$. Then the family $(P_z)_{z \in \Omega} := (\phi^* \tilde{P}_z)_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's on U .*

This allows us to define holomorphic families of Ψ_H DO's on a general Heisenberg manifold (M, H) and acting on the sections of a vector bundle on $\mathcal{E}$. In this setting the main properties of holomorphic families of Ψ_H DO's are summarized below.

Proposition 1.3. *Let $(P_z)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E})$ be a holomorphic family of Ψ_H DO's. Then:*

- 1) *The family of principal symbols $(\sigma_*(P_z))_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(\mathfrak{g}^*M \setminus 0, \text{End } \mathcal{E}))$.*
- 2) *The family of transpose operators $(P_z^t)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E}^*)$ is a holomorphic family of Ψ_H DO's.*
- 3) *The family of adjoints $(P_z^*)_{z \in \Omega}$ is an anti-holomorphic family of Ψ_H DO's, that is, $(P_{\bar{z}}^*)_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's.*

Finally, let us mention that the almost homogeneous approach to the Heisenberg calculus will have further applications in [Po10] for constructing a class of Ψ_H DO's with parameter containing the resolvents of hypoelliptic Ψ_H DO's (see also [Po12]).

1.2. Complex powers of hypoelliptic operators. One the main aims of this paper is to realize complex powers of hypoelliptic operators as holomorphic families of Ψ_H DO's. As alluded to above we cannot carry out the standard approach of Seeley [Se] in the Heisenberg setting. Therefore, we have to rely on another approach based on the pseudodifferential representation of the heat kernel of [BGS]. This is quite suitable for dealing differential operators, which is enough to cover may

examples. The general case will be dealt with in [Po10] using a different approach using a new kind of pseudodifferential representation of the resolvent.

Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a hypoelliptic selfadjoint differential operator and assume that P is bounded from below. It is well known the heat semigroup e^{-tP} , $t \geq 0$, allows us to invert the heat operator $P + \partial_t$. Conversely, constructing a suitable pseudodifferential calculus nesting parametrices for $P + \partial_t$ allows us to give a pseudodifferential representation for the heat kernel of P .

In the elliptic setting the relevant pseudodifferential calculus is the Volterra calculus of Greiner [Gre] and Piriou [Pi]. The latter is only a simple modification of the classical pseudodifferential calculus in order to take into account the parabolicity and the Volterra property with respect to the time variable of the heat equation. Moreover, this approach holds in fairly greater generality and has many applications ([BGS], [BS1], [BS2], [Gre], [Kr1], [Kr2], [KS], [Me2], [Pi], [Po4], [Po7]).

The Greiner's approach has been extended to the Heisenberg calculus in [BGS], mostly with the purpose of deriving the small time heat kernel asymptotics for the Kohn Laplacian on CR manifolds. In particular, a class of Volterra $\Psi_H DO$'s is obtained that contains parametrices for the heat operator $P + \partial_t$. As a consequence, once the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol, the inverse of $P + \partial$ is a Volterra- $\Psi_H DO$ which, in turn, yields a pseudodifferential representation of the heat kernel of P .

Assume now that P is positive. Thanks to the spectral theorem we can define the complex power of P^s , $s \in \mathbb{C}$, as an unbounded operator on $L^2(M, \mathcal{E})$ which is bounded for $\Re s \leq 0$. Moreover, for $\Re s < 0$ the Mellin formula holds,

$$(1.2) \quad P^s = \Gamma(s)^{-1} \int_0^\infty t^s (1 - \Pi_0(P)) e^{-tP} \frac{dt}{t},$$

where $\Pi_0(P)$ denotes the orthogonal projection onto the kernel of P . Combining this formula with the pseudodifferential representation of the heat kernel of P allows us to prove:

Theorem 1.4. *Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a positive differential operator of Heisenberg order m such that the principal symbol of the heat operator $P + \partial_t$ is an invertible Volterra-Heisenberg symbol. Then the family $(P^s)_{s \in \mathbb{C}}$ of the complex powers of P , defined by functional calculus on $L^2(M, \mathcal{E})$, is a holomorphic 1-parameter group of $\Psi_H DO$'s such that $\text{ord} P^s = ms$ for any $s \in \mathbb{C}$.*

In [BGS] it was shown that for a selfadjoint sublaplacian $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ the principal symbol of $\Delta + \partial_t$ is invertible under a condition closely related to the Rockland condition and the invertibility of the principal symbol of Δ (see [BGS, 5.23] and Section 6). Therefore, in the case of a sublaplacian we get:

Theorem 1.5. *Let $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a positive sublaplacian satisfying the condition of [BGS, 5.23]. Then the family $(\Delta^s)_{s \in \mathbb{C}}$ of the complex powers of Δ is a holomorphic 1-parameter group of $\Psi_H DO$'s such that $\text{ord} \Delta^s = 2s$ for any $s \in \mathbb{C}$.*

This theorem holds for the following kinds of sublaplacians:

- (a) A selfadjoint sum of squares $\nabla_{X_1}^* \nabla_{X_1} + \dots + \nabla_{X_m}^* \nabla_{X_m}$, where the vector fields $X_1, \dots, X_m$ linearly span H and ∇ is a connection on $\mathcal{E}$, provided that the Levi form of (M, H) is nonvanishing;
- (b) The Kohn Laplacian on a CR manifold acting on (p, q) -forms under condition $Y(q)$.
- (c) The horizontal Laplacian on a Heisenberg manifold acting on horizontal forms of degree k under condition $X(k)$.

For more general operators we show that, provided that the Levi form of (M, H) has constant rank, the Rockland condition for P is enough to deal with invertibility of the principal symbol of $P + \partial_t$ (see Theorem 1.11 below). This allows us to get:

Theorem 1.6. *Suppose that the Levi form of (M, H) has constant rank and let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a positive differential operator of Heisenberg order m such that P satisfies the Rockland condition at every point. Then the family $(P^s)_{s \in \mathbb{C}}$ of the complex powers of P is a holomorphic 1-parameter group of $\Psi_H DO$'s such that $\text{ord} P^s = ms$ for any $s \in \mathbb{C}$.*

In particular, Theorem 1.6 is valid for the contact Laplacian on a contact manifold. In this context this allows us to fill a technical gap in the proof of by Julg-Kasparov [JK] of the Baum-Connes conjecture for $SU(n, 1)$ (see [Po9]).

Finally, let us mention that a similar approaches to complex powers of (hypo)elliptic operators have been used by Mathai-Melrose-Singer [MMS] and Melrose [Me3] in the context of projective pseudodifferential operators on Azamaya bundles.

1.3. Weighted Sobolev spaces. Assuming that the Levi form of (M, H) is nowhere vanishing, we can construct a scale $W_H^s(M)$, $s \in \mathbb{R}$, of weighted Sobolev spaces obtained as follows.

Let $X_1, \dots, X_m$ be vectors fields spanning H and let $\Delta_X = X_1^* X_1 + \dots + X_m^* X_m$. Then by Theorem 1.5 the complex powers $(1 + \Delta_X)^s$ gives rise to a 1-parameter group of $\Psi_H DO$'s. We then define $W_H^s(M)$ as the Hilbert space consisting of distributions $u \in \mathcal{D}'(M)$ such that $(1 + \Delta_X)^{\frac{s}{2}} u \in L^2(M)$ together with the Hilbertian norm,

$$(1.3) \quad \|u\|_{W_H^s} = \|(1 + \Delta_X)^{\frac{s}{2}} u\|_{L^2}, \quad u \in W_H^s(M).$$

Neither the underlying space of $W_H^s(M)$, $s \in \mathbb{R}$, nor its topology depend on the choice of the vector fields $X_1, \dots, X_m$ (Proposition 7.2). Moreover, these weighted Sobolev spaces are nicely related to the standard Sobolev spaces $L_s^2(M)$ and to the weighted Sobolev spaces of Folland-Stein [FS1], as we have:

Proposition 1.7. *1) For $k = 1, 2, \dots$ the weighted Sobolev spaces $W_H^k(M)$ and $S_k^2(M)$ agree as spaces and bear the same topology.*

2) For $s \in \mathbb{R}$ the following continuous embedding hold:

$$(1.4) \quad \begin{aligned} L_s^2(M) &\hookrightarrow W_H^s(M) \hookrightarrow L_{s/2}^2(M) && \text{if } s \geq 0, \\ L_{s/2}^2(M) &\hookrightarrow W_H^s(M) \hookrightarrow L_s^2(M) && \text{if } s < 0. \end{aligned}$$

We can similarly define weighted Sobolev spaces $W_H^s(M, \mathcal{E})$ of distributional sections of $\mathcal{E}$ and the embedding (1.4) hold *verbatim* in this setting. In addition, we can localize the Sobolev spaces $W_H^s(M, \mathcal{E})$, that is, we can speak about distributional sections which are W_H^s near a point (see Definition 7.10 and Lemma 7.11).

The main interest of the weighted Sobolev spaces $W_H^s(M, \mathcal{E})$, $s \in \mathbb{R}$, with respect to the standard Sobolev spaces is that they allow us to get sharper regularity estimates for $\Psi_H DO$'s, for we have:

Proposition 1.8. *Let $P \in \Psi_H^m(M, \mathcal{E})$, set $k = \Re m$ and let $s \in \mathbb{R}$.*

1) The operator P extends to a continuous linear mapping from $W_H^{s+k}(M, \mathcal{E})$ to $W_H^s(M, \mathcal{E})$.

2) Assume that the principal symbol of P is invertible. Then, for any $u \in \mathcal{D}'(M, \mathcal{E})$ and any $x_0 \in M$, we have

$$(1.5) \quad Pu \in W_H^s(M, \mathcal{E}) \implies u \in W_H^{s+k}(M, \mathcal{E}),$$

$$(1.6) \quad Pu \text{ is } W_H^s \text{ near } x_0 \implies u \text{ is } W_H^{s+k} \text{ near } x_0.$$

In fact, for any $s' \in \mathbb{R}$ we have the estimate,

$$(1.7) \quad \|u\|_{W_H^{s+k}} \leq C_{ss'} (\|Pu\|_{W_H^{s'}} + \|u\|_{W_H^{s'}}), \quad u \in W_H^{s+k}(M, \mathcal{E}).$$

We can also obtain a version of Proposition 1.8 for holomorphic families and complex powers of hypoelliptic operators as follows.

Proposition 1.9. 1) Let $\Omega \subset \mathbb{C}$ be open and let $(P_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi_H^*(M, \mathcal{E}))$. Assume there exists a real m such that $\Re \text{ord} P_z \leq m < \infty$. Then for any $s \in \mathbb{R}$ the family $(P_z)_{z \in \Omega}$ defines a holomorphic family with values in $\mathcal{L}(W_H^{s+m}(M, \mathcal{E}), W_H^s(M, \mathcal{E}))$.

2) Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a positive differential operator of Heisenberg order m such that the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol. Then, for any reals a and s , the complex powers of P satisfy

$$(1.8) \quad (P^z)_{\Re z < a} \in \text{Hol}(\Re z < a, \mathcal{L}(W_H^{s+ma}(M, \mathcal{E}), W_H^s(M, \mathcal{E}))).$$

1.4. Rockland condition and the heat equation. Since Theorem 1.4 and the main results of [BGS] hold under the condition that the principal symbol of the heat operator $P + \partial_t$ is an invertible Volterra-Heisenberg symbol, it stands to reason to find criterions for the invertibility of the latter. We show here that, to a large extent, the Rockland condition and the positivity of the principal symbol of P are enough to imply the invertibility of the principal symbol of $P + \partial_t$. In particular, the results of [BGS] and Theorem 1.5 are valid for a very wide class of hypoelliptic operators.

Here we say that a homogeneous symbol p_m of degree m is positive if it can be put into the form $\bar{q}_{\frac{m}{2}} * q_{\frac{m}{2}}$ for some symbol $q_{\frac{m}{2}}$ homogeneous of degree $\frac{m}{2}$. It may be difficult in practice to check whether an operator has a positive principal symbol only by looking at this principal symbol. Nevertheless, for hypoelliptic operators we have:

Proposition 1.10. Assume that the Levi form of (M, H) has constant rank and let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of even Heisenberg order m with an invertible principal symbol. Then the following are equivalent:

- (i) The operator P is bounded from below.
- (ii) The principal symbol $\sigma_m(P)$ of P is positive.

In particular, for any lower order selfadjoint perturbation R the operator $P + R$ remains bounded from below.

Next, we prove:

Theorem 1.11. Assume that the Levi form of (M, H) has constant rank and let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of even Heisenberg order m which is bounded from below and satisfies the Rockland condition at every point. Then:

- 1) The principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol.
- 2) The heat operator $P + \partial_t$ admits a Volterra- Ψ_H DO inverse;
- 3) The heat kernel $k_t(x, y)$ of P has an asymptotics in $C^\infty(M, (\text{End } \mathcal{E}) \otimes |\Lambda|(M))$ of the form

$$(1.9) \quad k_t(x, x) \sim_{t \rightarrow 0^+} t^{-\frac{d+2}{m}} \sum t^{\frac{2j}{m}} a_j(P)(x),$$

where the density $a_j(P)(x)$ is locally computable in terms of the symbol $q_{-m-2j}(x, \xi, \tau)$ of degree $-m-2j$ of any Volterra- Ψ_H DO parametrix for $P + \partial_t$.

Notice that we only have to prove the first statement since the results of [BGS] allows us to deduce the last two statements from the first one. This is done by making use of Theorem 1.4 and of results of Christ-Geller-Głowacki-Polin [CGGP] and Folland-Stein [FS2] for Rockland operators on nilpotent graded Lie groups.

The above theorem is valid for the following operators:

- (a) The conformal powers $\square_\theta^{(k)}$ of the (scalar) Tanaka Laplacian acting on functions on a strictly pseudoconvex CR manifold;
- (b) The contact Laplacian on a contact manifold.

These examples are not covered by the results of [BGS].

On the other hand, if Δ is a sublaplacian then, when the Levi form of (M, H) is nondegenerate, it is shown in [BG] that the invertibility of the principal symbol of Δ is equivalent to the Rockland condition, but is weaker than the condition considered in [BGS, 5.23] (see [BG, 18.4] and Section 4 for the precise statement of the condition). Anyway, using Theorem 1.11 we get:

Theorem 1.12. *Assume that the Levi form of (M, H) is nondegenerate and let $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint sublaplacian which is bounded from below and satisfies the weaker condition of [BG, 18.4]. Then the conclusions of Theorem 1.11 are valid for $\Delta + \partial_t$ and the heat kernel of Δ .*

1.5. Spectral asymptotics for hypoelliptic operators. One interesting application of the heat kernel asymptotics in [BGS] and in (1.9) is to allow us to derive spectral asymptotics for hypoelliptic operators as follows.

Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of Heisenberg order m which is bounded from below and such that the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol.

Let $\lambda_0(P) \leq \lambda_1(P) \leq \dots$ denote the eigenvalues of P counted with multiplicity and let $N(P; \lambda)$ denote its counting function, that is,

$$(1.10) \quad N(P; \lambda) = \#\{k \in \mathbb{N}; \lambda_k(P) \leq \lambda\}, \quad \lambda \geq 0.$$

Then we prove:

Theorem 1.13. *Under the above assumptions the following hold.*

1) As $t \rightarrow 0^+$ we have

$$(1.11) \quad \text{Tr } e^{-tP} \sim t^{-\frac{d+2}{m}} \sum t_m^{2j} A_j(P), \quad A_j(P) = \int_M \text{tr}_{\mathcal{E}} a_j(P)(x).$$

where the density $a_j(P)(x)$ is the coefficient of $t^{\frac{j-d+2}{m}}$ in the asymptotics (8.10).

2) We have $A_0(P) > 0$.

2) As $\lambda \rightarrow \infty$ we have

$$(1.12) \quad N(P; \lambda) \sim \nu_0(P) \lambda^{\frac{d+2}{m}}, \quad \nu_0(P) = \Gamma(1 + \frac{d+2}{m})^{-1} A_0(P).$$

3) As $k \rightarrow \infty$ we have

$$(1.13) \quad \lambda_k(P) \sim \left(\frac{k}{\nu_0(P)} \right)^{\frac{m}{d+2}}.$$

The first asymptotics is an immediate consequence of (1.9). Moreover, once proved that we have $A_0(P) > 0$ the asymptotics (1.12) and (1.13) follows from (1.11) and Karamata's Tauberian theorem. The bulk of the proof is thus to prove the second assertion, which is obtained by making use of spectral theoretic arguments.

By relying on other pseudodifferential calculi several authors have obtained Weyl asymptotics closely related to (1.12) in the more general setting of hypoelliptic with multicharacteristics (see [II], [Me1], [MS], [Mo1], [Mo2]).

As far as the Heisenberg setting is concerned, the approach using the Volterra-Heisenberg calculus has two main advantages.

First, the pseudodifferential analysis is significantly simpler. In particular, the Volterra-Heisenberg calculus yields for free the heat kernel asymptotics once the principal symbol of the heat operator is shown to be invertible, for which it is enough to use the Rockland condition in many cases.

Second, as the Volterra-Heisenberg calculus take fully into account the underlying Heisenberg geometry of the manifold and is invariant by change of Heisenberg coordinates, we can very effectively deal with operators admitting normal forms (see Proposition 9.4 on this point). In particular, as

explained below, we can reformulate in geometric terms the Weyl asymptotics (1.12) for the main geometric operators on CR and contact manifolds.

1.6. Weyl asymptotics and CR geometry. Let M^{2n+1} be a compact κ -strictly pseudoconvex CR manifold, together with a pseudohermitian contact form θ , that is, a contact form such that the associated Levi form has signature $(n - \kappa, \kappa, 0)$. We also endow M with a Levi metric compatible with θ . Then the volume of M with respect to this Levi metric is independent of the choice of the Levi form and is equal to

$$(1.14) \quad \text{vol}_\theta M = \frac{(-1)^\kappa}{n!2^n} \int_M \theta \wedge d\theta^n.$$

We call $\text{vol}_\theta M$ the pseudohermitian volume of (M, θ) . We can relate the Weyl asymptotics (1.12) for the Kohn Laplacian and the horizontal sublaplacian to $\text{vol}_\theta M$ as follows.

First, for $\mu \in (-n, n)$ we let

$$(1.15) \quad \nu(\mu) = (2\pi)^{-(n+1)} \int_{-\infty}^{\infty} e^{-\mu\xi_0} \left(\frac{\xi_0}{\sinh \xi_0} \right)^n d\xi_0.$$

Second, in the case of the Kohn Laplacian we obtain:

Theorem 1.14. *Let $\square_b : C^\infty(M, \Lambda^{*,*}) \rightarrow C^\infty(M, \Lambda^{*,*})$ denote the Kohn Laplacian associated to the Levi metric on M . Then, for $p, q = 0, \dots, n$ with $q \neq \kappa$ and $q \neq n - \kappa$, as $\lambda \rightarrow \infty$ we have*

$$(1.16) \quad N(\square_{b|\Lambda^{p,q}}; \lambda) \sim \alpha_{n\kappa pq} (\text{vol}_\theta M) \lambda^{n+1},$$

$$(1.17) \quad \alpha_{n\kappa pq} = \frac{1}{2^{n+1}} \binom{n}{p} \sum_{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa)} \binom{n-\kappa}{k} \binom{\kappa}{q-k} \nu(n-2(\kappa-q+2k)).$$

In the strictly pseudoconvex case, i.e. $\kappa = 0$, the result is an easy consequence of [BGS, Thm. 8.31], but in the case $\kappa \geq 1$ this seems to be new.

Next, in the CR setting the horizontal sublaplacian preserves the bidegree and, in the same way as with the Kohn Laplacian, we get:

Theorem 1.15. *Let Δ_b be the horizontal sublaplacian associated to the Levi metric on M . Then for $p, q = 0, \dots, n$, with $(p, q) \neq (\kappa, n - \kappa)$ and $(p, q) \neq (n - \kappa, \kappa)$, as $\lambda \rightarrow \infty$ we have*

$$(1.18) \quad N(\Delta_{b|\Lambda^{p,q}}; \lambda) \sim \beta_{npq} (\text{vol}_\theta M) \lambda^{n+1},$$

$$(1.19) \quad \beta_{n\kappa pq} = \sum_{\substack{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa) \\ \max(0, p-\kappa) \leq l \leq \min(p, n-\kappa)}} \binom{n-\kappa}{l} \binom{\kappa}{p-l} \binom{n-\kappa}{k} \binom{\kappa}{q-k} \nu(2(q-p) + 4(l-k)).$$

Finally, as the Weyl asymptotics (1.12) depends only on the principal symbol of the operator P , in the strictly pseudoconvex case, i.e. $\kappa = 0$, we can also deal with the conformal powers of the horizontal Laplacian.

Theorem 1.16. *Assume that (M^{2n+1}, θ) is a strictly pseudoconvex pseudohermitian manifold and for $k = 1, \dots, n+1$ let $\square_\theta^{(k)} : C^\infty(M) \rightarrow C^\infty(M)$ be a k 'th conformal power of Δ_b associated to θ . Then as $\lambda \rightarrow \infty$ we have*

$$(1.20) \quad N(\square_\theta^{(k)}; \lambda) \sim \nu(0) (\text{vol}_\theta M) \lambda^{\frac{n+1}{k}}.$$

1.7. Weyl asymptotics and contact geometry. Let (M^{2n+1}, θ) be a compact orientable contact manifold. We assume that the Heisenberg bundle $H = \ker \theta$ has a calibrated almost complex structure J so that $d\theta(X, JX) = -d\theta(JX, X) > 0$ for any section X of $H \setminus 0$. We then endow M with the Riemannian metric $g_\theta = d\theta(\cdot, J\cdot) + \theta^2$. The volume of M with respect to g_θ depends only on θ and is equal to:

$$(1.21) \quad \text{vol}_\theta M = \frac{1}{n!} \int_M d\theta^n \wedge \theta.$$

We call $\text{vol}_\theta M$ the contact volume of M .

We can relate the Weyl asymptotics for the horizontal sublaplacian to the contact volume to get:

Theorem 1.17. *Let $\Delta_b : C^\infty(M, \Lambda_{\mathbb{C}}^* H^*) \rightarrow C^\infty(M, \Lambda_{\mathbb{C}}^* H^*)$ be the horizontal sublaplacian associated to the metric g_θ . Then, for $k = 0, \dots, 2n$ with $k \neq n$, as $\lambda \rightarrow \infty$ we have*

$$(1.22) \quad N(\Delta_b|_{\Lambda_{\mathbb{C}}^k H^*}; \lambda) \sim \gamma_{nk} (\text{vol}_\theta M) \lambda^{n+1}, \quad \gamma_{nk} = 2^{-n} \sum_{p+q=k} \binom{n}{p} \binom{n}{q} \nu(p-q).$$

Note that when (M, θ) is a strictly pseudoconvex pseudohermitian manifold the asymptotics (1.22) is compatible with (1.18) because the contact volume differs from the pseudohermitian volume by a factor of 2^{-n} .

Finally, we can also deal with the contact Laplacian as follows.

Theorem 1.18. *Let $\Delta_R : C^\infty(M, \Lambda^* \oplus \Lambda_*^n) \rightarrow C^\infty(M, \Lambda^* \oplus \Lambda_*^n)$ be the contact Laplacian on M .*

1) *For $k = 0, \dots, 2n$ with $k \neq n$ there exists a universal constant $\nu_{nk} > 0$ depending only on n and k such that as $\lambda \rightarrow \infty$ we have*

$$(1.23) \quad N(\Delta_R|_{\Lambda^k}) \sim \nu_{nk} (\text{vol}_\theta M) \lambda^{n+1}.$$

2) *For $j = 1, 2$ there exists a universal constant $\nu_n^{(j)} > 0$ depending only on n and j such that as $\lambda \rightarrow \infty$ we have*

$$(1.24) \quad N(\Delta_R|_{\Lambda_n^j}) \sim \nu_n^{(j)} (\text{vol}_\theta M) \lambda^{\frac{n+1}{2}}.$$

1.8. Organization of the paper. The paper is organized as follows. We start by reviewing the background needed in this paper. In Section 2 we recall the main definitions and examples concerning Heisenberg manifolds and their tangent Lie group bundles following the point of view of [Po6]. In Section 3 we recall the definitions of main operators on Heisenberg manifolds: sum of squares, Kohn Laplacian, horizontal sublaplacian and its conformal powers and the contact Laplacian. In Section 4 we give a detailed overview of the Heisenberg calculus of [BG] and [Tay], following closely the expositions of [BG] and [Po8].

In Section 5 we define holomorphic families of $\Psi_H\text{DO}$'s and check their main properties. In Section 6, after having recalled the pseudodifferential representation of the heat kernel of [BGS], we make use of the latter to construct complex powers of positive hypoelliptic operators as holomorphic families of $\Psi_H\text{DO}$'s. In Section 7 we define the weighted Sobolev spaces $W_H(M, \mathcal{E})$ and show that they give sharp regularity results for $\Psi_H\text{DO}$'s. In Section 8 we show that if P is selfadjoint differential operator bounded from below, then the fact that P satisfies the Rockland condition at every point is enough to insure us the invertibility of the principal symbol of $P + \partial_t$. This shows that the results of [BGS] and Section 6 are valid for a wide class of hypoelliptic operators.

In the last three sections we deal with spectral asymptotics for hypoelliptic operators on Heisenberg manifolds. In Section 9 we derive general spectral asymptotics for such operators on a general Heisenberg manifold. We specialize them in Section 10 to the Kohn Laplacian and the horizontal sublaplacian and its conformal powers on a CR manifold. Finally, in Section 11 we look at these asymptotics in the special cases of the horizontal sublaplacian and of the contact Laplacian on a contact manifold.

Acknowledgements. The author is grateful to Alain Connes, Charles Epstein, Bernard Helffer, Henri Moscovici and Michel Rumin for helpful and stimulating discussions. He wish also to thank for its hospitality the Mathematics Department of Princeton University, where this paper was finally completed.

In addition, the research of the paper was partially supported by a graduate fellowships from the French Ministry of National Education and Research, by a fellowship from the RT Network geometric analysis HPCRN-CT-1999-00118, and by the NSF grant DMS 0409005.

Finally, some of the results of this paper were announced in [Po2] and presented as part of the author's PhD thesis at University of Paris-Sud (Orsay, France), made under the supervision of Alain Connes.

2. HEISENBERG MANIFOLDS AND THEIR TANGENT LIE GROUP BUNDLES

In this section we recall the main facts about Heisenberg manifolds and their tangent Lie group bundle. The exposition here follows closely that of [Po6].

Definition 2.1. 1) A Heisenberg manifold is a smooth manifold M equipped with a distinguished hyperplane bundle $H \subset TM$.

2) A Heisenberg diffeomorphism ϕ from a Heisenberg manifold (M, H) onto another Heisenberg manifold (M', H') is a diffeomorphism $\phi: M \rightarrow M'$ such that $\phi^*H = H'$.

Definition 2.2. Let (M^{d+1}, H) be a Heisenberg manifold. Then:

- 1) A (local) H -frame for TM is a (local) frame $X_0, X_1, \dots, X_d$ of TM so that $X_1, \dots, X_d$ span H .
- 2) A local Heisenberg chart is a local chart with a local H -frame of TM over its domain.

The main examples of Heisenberg manifolds are the following.

a) *Heisenberg group.* The $(2n + 1)$ -dimensional Heisenberg group $\mathbb{H}^{2n+1}$ is $\mathbb{R}^{2n+1} = \mathbb{R} \times \mathbb{R}^n$ equipped with the group law,

$$(2.1) \quad x.y = (x_0 + y_0 + \sum_{1 \leq j \leq n} (x_{n+j}y_j - x_jy_{n+j}), x_1 + y_1, \dots, x_{2n} + y_{2n}).$$

A left-invariant basis for its Lie algebra $\mathfrak{h}^{2n+1}$ is then provided by the vector-fields,

$$(2.2) \quad X_0 = \frac{\partial}{\partial x_0}, \quad X_j = \frac{\partial}{\partial x_j} + x_{n+j} \frac{\partial}{\partial x_0}, \quad X_{n+j} = \frac{\partial}{\partial x_{n+j}} - x_j \frac{\partial}{\partial x_0}, \quad 1 \leq j \leq n,$$

which for $j, k = 1, \dots, n$ and $k \neq j$ satisfy the relations,

$$(2.3) \quad [X_j, X_{n+k}] = -2\delta_{jk}X_0, \quad [X_0, X_j] = [X_j, X_k] = [X_{n+j}, X_{n+k}] = 0.$$

In particular, the subbundle spanned by the vector fields $X_1, \dots, X_{2n}$ yields a left-invariant Heisenberg structure on $\mathbb{H}^{2n+1}$.

b) *Foliations.* Recall that a (smooth) foliation is a manifold M together with a subbundle $\mathcal{F} \subset TM$ which is integrable in the Froebenius' sense, i.e. so that $[\mathcal{F}, \mathcal{F}] \subset \mathcal{F}$. Therefore, any codimension 1 foliation is a Heisenberg manifold.

c) *Contact manifolds.* Opposite to foliations are contact manifolds: a *contact structure* on a manifold M^{2n+1} is given by a global non-vanishing 1-form θ on M such that $d\theta$ is non-degenerate on $H = \ker \theta$. In particular, (M, H) is a Heisenberg manifold. In fact, by Darboux's theorem any contact manifold (M^{2n+1}, θ) is locally contact-diffeomorphic to the Heisenberg group $\mathbb{H}^{2n+1}$ equipped with its standard contact form $\theta^0 = dx_0 + \sum_{j=1}^n (x_j dx_{n+j} - x_{n+j} dx_j)$.

d) *Confoliations.* According to Elyashberg-Thurston [ET] a *confoliation structure* on an oriented manifold M^{2n+1} is given by a global non-vanishing 1-form θ on M such that $(d\theta)^n \wedge \theta \geq 0$. In particular, when $d\theta \wedge \theta = 0$ (resp. $(d\theta)^n \wedge \theta > 0$) we are in presence of a foliation (resp. a contact structure). In any case the hyperplane bundle $H = \ker \theta$ defines a Heisenberg structure on M .

e) CR manifolds. A CR structure on an orientable manifold M^{2n+1} is given by a rank n complex subbundle $T_{1,0} \subset T_{\mathbb{C}}M$ which is integrable in Froebenius' sense and such that $T_{1,0} \cap T_{0,1} = \{0\}$, where $T_{0,1} = \overline{T_{1,0}}$. Equivalently, the subbundle $H = \Re(T_{1,0} \oplus T_{0,1})$ has the structure of a complex bundle of (real) dimension $2n$. In particular, the pair (M, H) forms a Heisenberg manifold.

Moreover, since M and H is orientable by means of its complex structure the normal bundle TM/H is orientable, hence admits a global nonvanishing section T . Let θ be a global section of T^*M/H^* such that $\theta(T) = 1$ and θ annihilates H . Then Kohn's Levi form is the form L_θ on $T_{1,0}$ such that, for sections Z and W of $T_{1,0}$, we have

$$(2.4) \quad L_\theta(Z, W) = -id\theta(Z, \bar{W}) = i\theta([Z, \bar{W}]).$$

We say that M is strictly pseudoconvex (resp. nondegenerate, κ -strictly pseudoconvex) when for some choice of θ the Levi form L_θ is everywhere positive definite (resp. is everywhere nondegenerate, has everywhere signature $(n - \kappa, \kappa, 0)$). In particular, when M is nondegenerate the 1-form θ is a contact form on M .

The main example of a CR manifold is that of the (smooth) boundary $M = \partial D$ of a complex domain $D \subset \mathbb{C}^n$. In particular, when D is strongly pseudoconvex (or strongly pseudoconcave) then M is strictly pseudoconvex.

2.1. The tangent Lie group bundle. A simple description of the tangent Lie group bundle of a Heisenberg manifold (M^{d+1}, H) is given as follows.

Lemma 2.3 ([Po6]). *The Lie bracket of vector fields induces on H a 2-form with values in TM/H ,*

$$(2.5) \quad \mathcal{L} : H \times H \longrightarrow TM/H,$$

so that for any sections X and Y of H near a point $a \in M$ we have

$$(2.6) \quad \mathcal{L}_a(X(a), Y(a)) = [X, Y](a) \mod H_a.$$

Definition 2.4. *The 2-form $\mathcal{L}$ is called the Levi form of (M, H) .*

The Levi form $\mathcal{L}$ allows us to define a bundle $\mathfrak{g}M$ of graded Lie algebras by endowing $(TM/H) \oplus H$ with the smooth fields of Lie Brackets and gradings such that

$$(2.7) \quad [X_0 + X', Y_0 + Y']_a = \mathcal{L}_a(X', Y') \quad \text{and} \quad t.(X_0 + X') = t^2 X_0 + tX' \quad t \in \mathbb{R},$$

for $a \in M$ and X_0, Y_0 in $T_a M/H_a$ and X', Y' in H_a .

Definition 2.5. *The bundle $\mathfrak{g}M$ is called the tangent Lie algebra bundle of M .*

As we can easily check $\mathfrak{g}M$ is a bundle of 2-step nilpotent Lie algebras which contains the normal bundle TM/H in its center. Therefore, its associated graded Lie group bundle GM can be described as follows. As a bundle GM is $(TM/H) \oplus H$ and the exponential map is merely the identity. In particular, the grading of GM is as in (2.7). Moreover, since $\mathfrak{g}M$ is 2-step nilpotent the Campbell-Hausdorff formula gives

$$(2.8) \quad (\exp X)(\exp Y) = \exp(X + Y + \frac{1}{2}[X, Y]) \quad \text{for sections } X, Y \text{ of } \mathfrak{g}M.$$

From this we deduce that the product on GM is such that

$$(2.9) \quad (X_0 + X').(Y_0 + X') = X_0 + Y_0 + \frac{1}{2}\mathcal{L}(X', Y') + X' + Y',$$

for sections X_0, Y_0 of TM/H and sections X', Y' of H .

Definition 2.6. *The bundle GM is called the tangent Lie group bundle of M .*

In fact, the fibers of GM as classified by the Levi form $\mathcal{L}$ as follows.

Proposition 2.7 ([Po6]). 1) Let $a \in M$. Then $\mathcal{L}_a$ has rank $2n$ if, and only if, as a graded Lie group $G_a M$ is isomorphic to $\mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$.

2) The Levi form $\mathcal{L}$ has constant rank $2n$ if, and only if, GM is a fiber bundle with typical fiber $\mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$.

Now, let $\phi : (M, H) \rightarrow (M', H')$ be a Heisenberg diffeomorphism from (M, H) onto another Heisenberg manifold (M', H') . Since $\phi_* H = H'$ we see that ϕ' induces a smooth vector bundle isomorphism $\bar{\phi} : TM/H \rightarrow TM'/H'$.

Definition 2.8. We let $\phi'_H : (TM/H) \oplus H \rightarrow (TM'/H') \oplus H'$ denote the vector bundle isomorphism such that

$$(2.10) \quad \phi'_H(a)(X_0 + X') = \bar{\phi}'(a)X_0 + \phi'(a)X',$$

for any $a \in M$ and any $X_0 \in T_a/H_a$ and $X' \in H_a$.

Proposition 2.9 ([Po6]). The vector bundle isomorphism ϕ'_H is an isomorphism of graded Lie group bundles from GM onto GM' . In particular, the Lie group bundle isomorphism class of GM depends only on the Heisenberg diffeomorphism class of (M, H) .

2.2. Heisenberg coordinates and nilpotent approximation of vector fields. It is interesting to relate the intrinsic description of GM above with the more extrinsic description of [BG] (see also [Be], [EMM], [FS1], [Gro], [Ro2]) in terms of the Lie group associated to a nilpotent Lie algebra of model vector fields.

First, let $a \in M$ and let us describe $\mathfrak{g}_a M$ as the graded Lie algebra of left-invariant vector fields on $G_a M$ by identifying any $X \in \mathfrak{g}_a M$ with the left-invariant vector fields L_X on $G_a M$ given by

$$(2.11) \quad L_X f(x) = \frac{d}{dt} f[(t \exp X) \cdot x]_{|t=0} = \frac{d}{dt} f[(tX) \cdot x]_{|t=0}, \quad f \in C^\infty(G_a M).$$

This allows us to associate to any vector fields X near a a unique left-invariant vector fields X^a on $G_a M$ such that

$$(2.12) \quad X^a = \begin{cases} L_{X_0(a)} & \text{if } X(a) \notin H_a, \\ L_{X(a)} & \text{otherwise,} \end{cases}$$

where $X_0(a)$ denotes the class of $X(a)$ modulo H_a .

Definition 2.10. The left-invariant vector fields X^a is called the model vector fields of X at a .

Let us look at the above construction in terms of a H -frame $X_0, \dots, X_d$ near a , i.e. of a local trivialization of the vector bundle $(TM/H) \oplus H$. For $j, k = 1, \dots, d$ we let

$$(2.13) \quad \mathcal{L}(X_j, X_k) = [X_j, X_k]X_0 = L_{jk}X_0 \mod H.$$

With respect to the coordinate system $(x_0, \dots, x_d) \rightarrow x_0 X_0(a) + \dots + x_d X_d(a)$ we can write the product law of $G_a M$ as

$$(2.14) \quad x \cdot y = (x_0 + \frac{1}{2} \sum_{j,k=1}^d L_{jk} x_j x_k, x_1, \dots, x_d).$$

Then the vector fields X_j^a , $j = 1, \dots, d$, in (2.12) is just the left-invariant vector fields corresponding to the vector e_j of the canonical basis of $\mathbb{R}^{d+1}$, that is, we have

$$(2.15) \quad X_0^a = \frac{\partial}{\partial x_0} \quad \text{and} \quad X_j^a = \frac{\partial}{\partial x_j} - \frac{1}{2} \sum_{k=1}^d L_{jk} x_k \frac{\partial}{\partial x_0}, \quad 1 \leq j \leq d.$$

In particular, for $j, k = 1, \dots, d$ we have the relations,

$$(2.16) \quad [X_j^a, X_k^a] = L_{jk}(a)X_0^a, \quad [X_j^a, X_0^a] = 0.$$

Now, let $\kappa : \text{dom } \kappa \rightarrow U$ be a Heisenberg chart near $a = \kappa^{-1}(u)$ and let $X_0, \dots, X_d$ be the associated H -frame of TU . Then there exists a unique affine coordinate change $x \rightarrow \psi_u(x)$ such that $\psi_u(u) = 0$ and $\psi_{u*}X_j(0) = \frac{\partial}{\partial x_j}$ for $j = 0, 1, \dots, d$. Indeed, if for $j = 1, \dots, d$ we set $X_j(x) = \sum_{k=0}^d B_{jk}(x) \frac{\partial}{\partial x_k}$ then we have

$$(2.17) \quad \psi_u(x) = A(u)(x - u), \quad A(u) = (B(u)^t)^{-1}.$$

Definition 2.11. 1) The coordinates provided by ψ_u are called the privileged coordinates at u with respect to the H -frame $X_0, \dots, X_d$.

2) The map ψ_u is called the privileged-coordinate map with respect to the H -frame $X_0, \dots, X_d$.

Remark 2.12. The privileged coordinates at u are called u -coordinates in [BG], but they correspond to the privileged coordinates of [Be] and [Gro] in the special case of a Heisenberg manifold.

Next, on $\mathbb{R}^{d+1}$ we consider the dilations

$$(2.18) \quad \delta_t(x) = t.x = (t^2x_0, tx_1, \dots, tx_d), \quad t \in \mathbb{R},$$

with respect to which $\frac{\partial}{\partial x_0}$ is homogeneous of degree -2 and $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_d}$ is homogeneous of degree -1 .

Since in the privileged coordinates at u we have $X_j(0) = \frac{\partial}{\partial x_j}$ we can write

$$(2.19) \quad X_j = \frac{\partial}{\partial x_j} + \sum_{k=0}^d a_{jk}(x) \frac{\partial}{\partial x_k}, \quad j = 0, 1, \dots, d,$$

where the a_{jk} 's are smooth functions such that $a_{jk}(0) = 0$. Therefore, we may define

$$(2.20) \quad X_0^{(u)} = \lim_{t \rightarrow 0} t^2 \delta_t^* X_0 = \frac{\partial}{\partial x_0},$$

$$(2.21) \quad X_j^{(u)} = \lim_{t \rightarrow 0} t^{-1} \delta_t^* X_j = \frac{\partial}{\partial x_j} + \sum_{k=1}^d b_{jk} x_k \frac{\partial}{\partial x_0}, \quad j = 1, \dots, d,$$

where for $j, k = 1, \dots, d$ we have set $b_{jk} = \partial_{x_k} a_{j0}(0)$.

Observe that $X_0^{(u)}$ is homogeneous of degree -2 and $X_1^{(u)}, \dots, X_d^{(u)}$ are homogeneous of degree -1 . Moreover, for $j, k = 1, \dots, d$ we have

$$(2.22) \quad [X_j^{(u)}, X_0^{(u)}] = 0 \quad \text{and} \quad [X_j^{(u)}, X_0^{(u)}] = (b_{kj} - b_{jk})X_0^{(u)}.$$

Thus, the linear space spanned by $X_0^{(u)}, X_1^{(u)}, \dots, X_d^{(u)}$ is a graded 2-step nilpotent Lie algebra $\mathfrak{g}^{(u)}$. In particular, $\mathfrak{g}^{(u)}$ is the Lie algebra of left-invariant vector fields over the graded Lie group $G^{(u)}$ consisting of $\mathbb{R}^{d+1}$ equipped with the grading (2.18) and the group law,

$$(2.23) \quad x.y = (x_0 + \sum_{j,k=1}^d b_{kj} x_j x_k, x_1, \dots, x_d).$$

Now, if near a we let $\mathcal{L}(X_j, X_k) = [X_j, X_k] = L_{jk}(x)X_0 \text{ mod } H$ then we have

$$(2.24) \quad [X_j^{(u)}, X_k^{(u)}] = \lim_{t \rightarrow 0} [t\delta_t^* X_j, t\delta_t^* X_k] = \lim_{t \rightarrow 0} t^2 \delta_t^* (L_{jk}(\circ \kappa^{-1}(x))X_0) = L_{jk}(a)X_0^{(u)}.$$

Comparing this with (2.16) and (2.22) then shows that $\mathfrak{g}^{(u)}$ has the same the constant structures as those of $\mathfrak{g}_a M$, hence is isomorphic to $\mathfrak{g}_a M$. Consequently, the Lie groups $G^{(u)}$ and $G_a M$ are isomorphic. In fact, as it follows from [BG] and [Po6] an explicit isomorphism is given by

$$(2.25) \quad \phi_u(x_0, \dots, x_d) = (x_0 - \frac{1}{4} \sum_{j,k=1}^d (b_{jk} + b_{kj}) x_j x_k, x_1, \dots, x_d).$$

Definition 2.13. Let $\varepsilon_u = \phi_u \circ \psi_u$. Then:

1) The new coordinates provided by ε_u are called *Heisenberg coordinates at u with respect to the H -frame $X_0, \dots, X_d$* .

2) The map ε_u is called the *u -Heisenberg coordinate map*.

Remark 2.14. The Heisenberg coordinates at u have been also considered in [BG] as a technical tool for inverting the principal symbol of a hypoelliptic sublaplacian.

Next, as it follows from [Po6, Lem. 1.17] we also have

$$(2.26) \quad \phi_* X_0^{(u)} = \frac{\partial}{\partial x_0} = X_0^a \quad \text{and} \quad \phi_* X_j^{(u)} = \frac{\partial}{\partial x_j} - \frac{1}{2} \sum_{k=1}^d L_{jk} x_k \frac{\partial}{\partial x_0} = X_j^a, \quad j = 1, \dots, d.$$

Since ϕ_u commutes with the Heisenberg dilations (2.18) using (2.20)–(2.21) we get

$$(2.27) \quad \lim_{t \rightarrow 0} t^2 \delta_t^* \phi_{u*} X_0^{(u)} = X_0^a \quad \text{and} \quad \lim_{t \rightarrow 0} t \delta_t^* \phi_{u*} X_j^{(u)} = X_j^a, \quad j = 1, \dots, d.$$

In fact, as shown in [Po6] for any vector fields X near a , as $t \rightarrow 0$ and in Heisenberg coordinates at a , we have

$$(2.28) \quad \delta_t^* X = \begin{cases} t^{-2} X^a + O(t^{-1}) & \text{if } X(a) \in H_a, \\ t^{-1} X^a + O(1) & \text{otherwise.} \end{cases}$$

Therefore, we obtain:

Proposition 2.15 ([Po6]). *In the Heisenberg coordinates centered at $m = \kappa^{-1}(u)$ the tangent Lie group $G_a M$ coincides with $G^{(u)}$ and for any vector fields X the model vector fields X^a approximates X near a in the sense of (2.28).*

One consequence of the equivalence between the two approaches to GM is a tangent approximation for Heisenberg diffeomorphisms as follows.

Let $\phi : (M, H) \rightarrow (M', H')$ be a Heisenberg diffeomorphism from (M, H) to another Heisenberg manifold (M', H') . We also endow $\mathbb{R}^{d+1}$ with the pseudo-norm,

$$(2.29) \quad \|x\| = (x_0^2 + (x_1^2 + \dots + x_d^2)^2)^{1/4}, \quad x \in \mathbb{R}^{d+1},$$

so that for any $x \in \mathbb{R}^{d+1}$ and any $t \in \mathbb{R}$ we have

$$(2.30) \quad \|t \cdot x\| = |t| \|x\|.$$

Proposition 2.16 ([Po6, Prop. 2.21]). *Let $a \in M$ and set $m' = \phi(a)$. Then, in Heisenberg coordinates at a and at a' the diffeomorphism $\phi(x)$ has a behavior near $x = 0$ of the form*

$$(2.31) \quad \phi(x) = \phi'_H(0)x + (O(\|x\|^3), O(\|x\|^2), \dots, O(\|x\|^2)).$$

In particular, there is no term of the form $x_j x_k$, $1 \leq j, k \leq d$, in the Taylor expansion of $\phi_0(x)$ at $x = 0$.

Remark 2.17. An asymptotics similar to (2.31) is given in [Be, Prop. 5.20] in privileged coordinates at u and $u' = \kappa_1(a')$, but the leading term there is only a Lie algebra isomorphism from $\mathfrak{g}^{(u)}$ onto $\mathfrak{g}^{(u')}$. This is only in Heisenberg coordinates that we recover the Lie group isomorphism $\phi'_H(a)$ as the leading term of the asymptotics.

Remark 2.18. An interesting application of Proposition 2.16 in [Po6] is the construction of the tangent groupoid $\mathcal{G}_H M$ of (M, H) as the differentiable groupoid encoding the smooth deformation of $M \times M$ to GM . This groupoid is the analogue in the Heisenberg setting of Connes' tangent groupoid of a manifold ([Co, II.5], [HS]) and it shows that GM is tangent to a in a differentiable fashion (compare [Be], [Gro]).

3. MOST COMMON OPERATORS ON HEISENBERG MANIFOLDS

In this section we recall the constructions of the most common operators on a Heisenberg manifold. At the exception of the contact Laplacian, all these operators are sublaplacians or coincides with an integer power of a sublaplacian up to a lower order term.

Recall that a sublaplacian on a Heisenberg manifold (M^{d+1}, H) acting on the sections of a vector bundle $\mathcal{E}$ over M is differential operator $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ such that, near any $a \in M$, there exists a H -frame $X_0, X_1, \dots, X_d$ of TM so that Δ takes the form

$$(3.1) \quad \Delta = - \sum_{j=1}^d X_j^2 + \text{lower order terms.}$$

3.1. Hörmander's sum of squares. Let $X_1, \dots, X_m$ be (real) vector fields on a manifold M^{d+1} and consider the sum of squares,

$$(3.2) \quad \Delta = -(X_1^2 + \dots + X_m^2).$$

By a celebrated theorem of Hörmander [Hö2] the operator Δ is hypoelliptic provided that the following bracket condition is satisfied: the vector fields $X_0, X_1, \dots, X_m$ together with their successive Lie brackets $[X_{j_1}, [X_{j_2}, \dots, X_{j_l}]]$ span the tangent bundle TM at every point.

When $X_1, \dots, X_m$ span a hyperplane bundle H then Δ is a sublaplacian with *real* coefficients and the bracket condition reduces to $H + [H, H] = TM$ or, equivalently, the Levi form of (M, H) is nonvanishing.

In fact, given a vector bundle $\mathcal{E}$, the theorem of Hörmander holds for more general sublaplacians $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ of the form

$$(3.3) \quad \Delta = -(\nabla_{X_1}^2 + \dots + \nabla_{X_m}^2) + L,$$

where ∇ is a connection on $\mathcal{E}$ and $L = L(X_1, \dots, X_m)$ is a first order polynomial with *real* coefficients in the vector fields $X_1, \dots, X_m$. In particular, if M is endowed with a positive density and $\mathcal{E}$ with a Hermitian metric, this includes the selfadjoint sum of squares,

$$(3.4) \quad \Delta = \nabla_{X_1}^* \nabla_{X_1} + \dots + \nabla_{X_m}^* \nabla_{X_m}^2.$$

3.2. The Kohn Laplacian on a CR manifold. In [KR] Kohn-Rossi showed that the Dolbeault complex on a bounded complex domain induces on its boundary a horizontal complex of differential forms. This was later extended by Kohn [Koh1] to the general setting of a CR manifold M^{2n+1} as follows.

Let M^{2n+1} be a CR manifold with CR bundle $T_{1,0} \subset T_{\mathbb{C}}M$ and set $T_{0,1} = \overline{T_{1,0}}$. Then the subbundle $H = \Re(T_{1,0} \oplus T_{0,1}) \subset TM$ admits an integrable complex structure and the splitting $H \otimes \mathbb{C} = T_{1,0} \oplus T_{0,1}$ gives rise to a decomposition $\Lambda H^* \otimes \mathbb{C} = \oplus_{0 \leq p, q \leq n} \Lambda^{p,q}$ into forms of bidegree (p, q) .

Assume that $T_{\mathbb{C}}M$ is endowed with a Hermitian metric such that $T_{1,0}$ and $T_{0,1}$ are orthogonal subspaces and complex conjugation is an (antilinear) isometry. This Hermitian metric gives rise to a Hermitian metric on $\Lambda^* T_{\mathbb{C}}^* M$ with respect to which the decomposition $\Lambda H^* \otimes \mathbb{C} = \oplus_{0 \leq p, q \leq n} \Lambda^{p,q}$ becomes orthogonal. Let $\Pi_{p,q} : \Lambda^* T_{\mathbb{C}}^* M \rightarrow \Lambda^{p,q}$ be the orthogonal projection onto $\Lambda^{p,q}$. Then the Kohn-Rossi operator $\bar{\partial}_b : C^\infty(M, \Lambda^{p,q}) \rightarrow C^\infty(M, \Lambda^{p,q+1})$ is given by

$$(3.5) \quad \bar{\partial}_b \eta = \Pi_{p,q+1}(d\eta), \quad \eta \in C^\infty(M, \Lambda^{p,q}).$$

Since the integrability of $T_{1,0}$ implies that $\bar{\partial}_b^2 = 0$, this yields chain complexes $\bar{\partial}_b : C^\infty(M, \Lambda^{p,*}) \rightarrow C^\infty(M, \Lambda^{p,*+1})$.

Endowing M with a smooth density $\rho > 0$ we let $\bar{\partial}_b^*$ denote the formal adjoint of $\bar{\partial}_b$. Then the Kohn Laplacian is

$$(3.6) \quad \square_b = (\bar{\partial}_b + \bar{\partial}_b^*)^2 = \bar{\partial}_b^* \bar{\partial}_b + \bar{\partial}_b \bar{\partial}_b^*.$$

The Kohn Laplacian is a sublaplacian (see [FS1, Sect. 13], [BG, Sect. 20]), so is not elliptic. Nevertheless, Kohn [Koh1] proved that under a geometric condition on the Levi form (2.4), the so-called condition $Y(q)$, the operator $\square_b$ acting on (p, q) -forms is hypoelliptic with loss of one derivative, i.e. for any compact $K \subset M$ we have

$$(3.7) \quad \|u\|_{s+1} \leq C_{Ks}(\|\square_b u\|_s + \|u\|_0) \quad \forall C_K^\infty(M, \Lambda^{p,q}).$$

where $\|\cdot\|_s$ denotes the norm of the Sobolev space $L_s^2(M, \Lambda^{p,q})$.

The condition $Y(q)$ at point $x \in M$ means that if we let $(r(x) - \kappa(x), \kappa(x), n - r(x))$ be the signature of L_θ at x , so that $r(x)$ is the rank of L_θ and $\kappa(x)$ the number of its negative eigenvalues, then we must have

$$(3.8) \quad q \notin \{\kappa(x), \kappa(x) + 1, \dots, \kappa(x) + n - r(x)\} \cup \{r(x) - \kappa(x), r(x) - \kappa(x) + 1, \dots, n - \kappa(x)\}.$$

Kohn's theorem then tells us that when the $Y(q)$ -condition holds everywhere $\square_b$ is hypoelliptic on (p, q) -forms.

For instance, when M is κ -strictly pseudoconvex, the $Y(q)$ -condition exactly means that we must have $q \neq \kappa$ and $q \neq n - \kappa$. In general this condition is equivalent to the existence of a parametrix within the Heisenberg calculus (cf. [BG], [Po8]; see also [Bo], [FS1]), from which we recover the hypoellipticity of $\square_b$.

Finally, let us mention that the condition $Y(q)$ is a sufficient, but not a necessary condition for the hypoellipticity of the Kohn Laplacian (see, e.g., [Koh2], [Ko], [Ni]).

3.3. The horizontal Laplacian on a Heisenberg manifold. Let (M^{d+1}, H) be a Heisenberg manifold endowed with a Riemannian metric. Then the horizontal sublaplacian is the differential operator $\Delta_b : C^\infty(M, \Lambda_{\mathbb{C}}^* H^*) \rightarrow C^\infty(M, \Lambda_{\mathbb{C}}^* H^*)$ is given by

$$(3.9) \quad \Delta_b = d_b^* d_b + d_b d_b^*, \quad d_b \alpha = \pi_b(d\alpha),$$

where $\pi_b : \Lambda_{\mathbb{C}}^* T^* M \rightarrow \Lambda_{\mathbb{C}}^* H^*$ denotes the orthogonal projection onto $\Lambda_{\mathbb{C}}^* H^*$.

This operator was first introduced by Tanaka [Ta] in the CR setting, but versions of this operator acting on functions were independently defined by Greenleaf [Gr] and Lee [Le]. Moreover, it can be shown that $d_b^2 = 0$ if, and only if, the subbundle H is integrable, so in general Δ_b is not the Laplacian of a chain complex.

On functions Δ_b is a sum of squares modulo a lower order term, hence is hypoelliptic by Hörmander's theorem. When acting on horizontal forms of higher degree, that is, on sections of $\Lambda_{\mathbb{C}}^k H^*$ with $k \geq 1$, it is was shown, in [Ta] and [Ru] in the contact case and in [Po8] in the general case, that the operator Δ_b is hypoelliptic when the condition $X(k)$ holds everywhere.

In the terminology of [Po8] the condition $X(k)$ at a point $x \in M$ means that, if we let $2r(x)$ denote the rank of the Levi form $\mathcal{L}$ at x , then we must have

$$(3.10) \quad k \notin \{r(x), r(x) + 1, \dots, d - r(x)\}.$$

For instance, if M^{2n+1} is a contact manifold or a nondegenerate CR manifold then the Levi form is everywhere nondegenerate, so $r(x) = 2n$ and the $X(k)$ -condition becomes $k \neq n$.

Assume now that M is a CR manifold with Heisenberg structure $H = \Re(T_{1,0} \oplus T_{0,1})$ and assume that $T_{\mathbb{C}} M$ is endowed with a Hermitian metric with respect to which $T_{1,0}$ and $T_{0,1}$ are orthogonal subspaces and complex conjugation is an isometry. Then we have $d_b = \bar{\partial}_b + \partial_b$, where ∂_b denotes the conjugate of $\bar{\partial}_b$, that is, $\partial_b \alpha = \overline{\bar{\partial}_b \omega}$ for any $\omega \in C^\infty(M, \Lambda_{\mathbb{C}}^* H^*)$. Moreover, as $\bar{\partial}_b \partial_b^* + \partial_b^* \bar{\partial}_b = \bar{\partial}_b^* \partial_b + \partial_b \bar{\partial}_b^* = 0$ (see [Ta]) we get

$$(3.11) \quad \Delta_b = \square_b + \bar{\square}_b,$$

where $\bar{\square}_b$ is the conjugate of $\square_b$. In particular, we see that the horizontal sublaplacian preserves the bidegree, i.e., it acts on (p, q) -forms.

In fact, as explained in [Po8, p. 34], one can show that the operator Δ_b acting on (p, q) -forms is hypoelliptic when at any point $x \in M$ the condition $Y(p, q)$ is satisfied. This means that at any point $x \in M$ we have

$$(3.12) \quad (p, q) \notin \{(\kappa(x)+j, r(x)-\kappa(x)+k); \max(j, k) \leq n-r(x)\} \cup \{(r(x)-\kappa(x)+j, \kappa(x)+k); \max(j, k) \leq n-r(x)\},$$

where $(r(x) - \kappa(x), \kappa(x), n - r(x))$ is the signature at x of the Levi form L_θ associated to some non-vanishing real 1-form θ annihilating $T_{1,0} \oplus T_{0,1}$. In particular, when M is κ -strictly pseudoconvex the $Y(p, q)$ reduces to $(p, q) \neq (\kappa, n - \kappa)$ and $(p, q) \neq (n - \kappa, \kappa)$.

3.4. The conformal powers of the horizontal sublaplacian on a strictly pseudoconvex CR manifold. Let (M^{2n+1}, θ) be a strictly pseudoconvex pseudohermitian manifold. Let $T_{1,0} \subset T_{\mathbb{C}}M$ be the CR bundle of M and define $T_{0,1} = \overline{T_{1,0}}$ and $H = \Re(T_{1,0} \oplus T_{0,1})$.

According to Webster [We] a pseudohermitian structure on M is given a choice of a 1-form θ annihilating H such that the associated Levi form L_θ is positive definite on $T_{1,0}$, so that θ is a contact form with respect to which the complex structure of H is calibrated, i.e. we have $d\theta(X, JX) > 0$ for any non-zero section of H .

We extend the L_θ into a Hermitian metric h_θ on $T_{\mathbb{C}}M$ that $T_{1,0}$ and $T_{0,1}$, are orthogonal subspaces, complex conjugation is an (antilinear) isometry and $h_{\theta|_{H^\perp}} = \theta^2$. Then as shown by Tanaka [Ta] and Webster [We] there is a unique unitary connection on $T_{\mathbb{C}}M$ preserving the pseudohermitian structure. Note that the contact form θ is unique up to a conformal change $f \rightarrow e^{2f}\theta$, $f \in C^\infty(M, \mathbb{R})$. In order to study the analogue in CR geometry of the Yamabe problem Jerison-Lee [JL1] (see also [JL2], [JL3]) introduced a conformal version of the scalar horizontal sublaplacian acting on functions,

$$(3.13) \quad \square_\theta : C^\infty(M) \longrightarrow C^\infty(M), \quad \square_\theta = \Delta_b + \frac{n}{n+2}R_n,$$

where R_n denotes the scalar curvature of the Tanaka-Webster connection. This is a conformal operator in the sense that

$$(3.14) \quad \square_{e^{2f}\theta} = e^{-(n+2)f} \square_\theta e^{nf} \quad \forall f \in C^\infty(M, \mathbb{R}).$$

Assume now that M is strictly pseudoconvex. Then the construction of Jerison-Lee has been recently generalized by Gover-Graham [GG]. Assuming the existence of a $(n+2)$ 'th root for the canonical bundle $\Lambda^{n,1}$ their constructions yield for $k = 1, 2, \dots, n+1$ a selfadjoint differential operator,

$$(3.15) \quad \square_\theta^{(k)} : C^\infty(M) \longrightarrow C^\infty(M),$$

such that the leading part (in the Heisenberg sense) of $\square_\theta^{(k)}$ agrees with that of Δ_b^k and we have

$$(3.16) \quad \square_{e^{2f}\theta}^{(k)} = e^{-(n+1+k)f} \square_\theta^{(k)} e^{(n+1-k)f}, \quad \forall f \in C^\infty(M, \mathbb{R}).$$

This is essentially the operator $P_{w,w'}$ of Theorem 1.1 of [GG] with $w = w' = \frac{k-n-1}{2}$, noticing that when $w' = w$ the density bundles $\mathcal{E}(w, w)$, on which the Gover-Graham's operators act, are in fact trivializable. For $k = 1$ we recover the operator $\square_\theta$ of Jerison-Lee.

3.5. Contact complex and contact Laplacian. Given an orientable contact manifold (M^{2n+1}, θ) the contact complex of Rumin [Ru] can be seen as an attempt to get on M a complex of horizontal differential forms by forcing out the equalities $d_b^2 = 0$ and $(d_b^*)^2$ as follows.

Let $H = \ker \theta$ and assume that H is endowed a calibrated almost complex structure $J \in \text{End}_{\mathbb{R}} H$, $J^2 = -1$, so that $d\theta(X, JX) = -d\theta(JX, X) > 0$ for any section X of H . We then get a Riemannian metric on M by letting

$$(3.17) \quad g_\theta = d\theta(., J.) + \theta^2.$$

Let T be the Reeb fields of θ . Then we have

$$(3.18) \quad d_b^2 = -\mathcal{L}_T \varepsilon(d\theta) = \varepsilon(d\theta) \mathcal{L}_T,$$

where $\varepsilon(d\theta)$ denotes the exterior multiplication by $d\theta$.

There are two ways of modifying the space $\Lambda_{\mathbb{C}}^* H^*$ of horizontal forms to get a complex. The first one is to force the equality $d_b^2 = 0$ by restricting the operator d_b to $\Lambda_2^* := \ker \varepsilon(d\theta) \cap \Lambda_{\mathbb{C}}^* H^*$ since this bundle is stable under d_b and on there d_b^2 vanishes.

The second way by similarly forcing the equality $(d_b^*)^2$ via the restriction of d_b^* to $\Lambda_1^* := \ker \iota(d\theta) \cap \Lambda_{\mathbb{C}}^* H^* = (\text{im } \varepsilon(d\theta))^\perp \cap \Lambda_{\mathbb{C}}^* H^*$, where $\iota(d\theta)$ denotes the interior product with $d\theta$. This means that we replace d_b by the operator $\pi_1 \circ d_b$, where π_1 is the orthogonal projection onto Λ_1^* .

On the other hand, since $d\theta$ is nondegenerate on H the operator $\varepsilon(d\theta) : \Lambda_{\mathbb{C}}^k H^* \rightarrow \Lambda_{\mathbb{C}}^{k+2} H^*$ is injective for $k \leq n-1$ and surjective for $k \geq n+1$. This implies that $\Lambda_2^k = 0$ for $k \leq n$ and $\Lambda_1^k = 0$ for $k \geq n+1$. Therefore, we only have two halves of complexes. As observed by Rumin we get a full complex by connecting the two halves by means of the (second order) differential operator,

$$(3.19) \quad D_R : \Lambda_1^n \longrightarrow \Lambda_2^{n+1}, \quad D_R = \mathcal{L}_T - d_b \varepsilon(d\theta)^{-1} d_b,$$

where $\varepsilon(d\theta)^{-1}$ is the inverse of $\varepsilon(d\theta) : \Lambda_{\mathbb{C}}^{n-1} H^* \rightarrow \Lambda_{\mathbb{C}}^{n+1} H^*$. In other words, letting $\Lambda^k = \Lambda_1^k$ for $k = 0, \dots, n-1$ and $\Lambda^k = \Lambda_2^k$ for $k = n+1, \dots, 2n$, we have the chain complex:

$$(3.20) \quad C^\infty(M) \xrightarrow{d_R} \dots \xrightarrow{d_R} C^\infty(M, \Lambda_1^n) \xrightarrow{D_R} C^\infty(M, \Lambda_2^n) \xrightarrow{d_R} \dots \xrightarrow{d_R} C^\infty(M, \Lambda^{2n}),$$

where $d_R : C^\infty(M, \Lambda^k) \rightarrow C^\infty(M, \Lambda^{k+1})$ is equal to $\pi_1 \circ d_b$ for $k = 0, \dots, n-1$ and to d_b for $k = n+1, \dots, 2n$. This complex is called the contact complex of M .

The contact Laplacian $\Delta_R : C^\infty(M, \Lambda^* \oplus \Lambda_*^n) \rightarrow C^\infty(M, \Lambda^* \oplus \Lambda_*^n)$ is given by the formulas,

$$(3.21) \quad \Delta_R = \begin{cases} (n-k)d_R d_R^* + (n-k+1)d_R^* d_R & \text{on } \Lambda^k, k = 0, \dots, n-1, \\ (d_R^* d_R)^2 + D_R^* D_R & \text{on } \Lambda_1^n, \\ D_R D_R^* + (d_R d_R^*) & \text{on } \Lambda_2^n, \\ (n-k+1)d_R d_R^* + (n-k)d_R^* d_R & \text{on } \Lambda^k, k = n+1, \dots, 2n. \end{cases}$$

Next, the almost complex structure J of H defines a bigrading on $\Lambda_{\mathbb{C}}^* H^*$. More precisely, we have an orthogonal splitting $H \otimes \mathbb{C} = T_{1,0} \oplus T_{0,1}$ with $T_{1,0} = \ker(J+i)$ and $T_{0,1} = \overline{T_{1,0}} = \ker(J-i)$. Therefore, if we consider the subbundles $\Lambda^{1,0} = T_{1,0}^*$ and $\Lambda^{0,1} = T_{0,1}^*$ of $H^* \otimes \mathbb{C} \subset T_{\mathbb{C}}^* M$ then we have the orthogonal decomposition $\Lambda_{\mathbb{C}}^* H^* = \bigoplus_{0 \leq p, q \leq n} \Lambda^{p,q}$ with $\Lambda^{p,q} = (\Lambda^{1,0})^p \wedge (\Lambda^{0,1})^q$. We then get a bigrading on Λ_j^n , $j = 1, 2$, by letting

$$(3.22) \quad \Lambda_j^n = \bigoplus_{p+q=n} \Lambda_j^{p,q}, \quad \Lambda_j^{p,q} = \Lambda_j^n \cap \Lambda^{p,q}.$$

For $k = 0, \dots, 2n$ with $k \neq n$ there exists a first order differential operator P_k such that

$$(3.23) \quad (n-k+2)\Delta_R = (n-k)(n-k+1)\Delta_b + P_k^* P_k \quad \text{on } \Lambda^k, k = 0, \dots, n-1,$$

$$(3.24) \quad (l-n+2)\Delta_R = (k-n)(k-n+1)\Delta_b + P_l^* P_l \quad \text{on } \Lambda^l, k = n+1, \dots, 2n.$$

For $j = 1, 2$ and $p+q = n$ there exist second order differential operators $P_{p,q}^{(j)}$ so that we have

$$(3.25) \quad 4\Delta_R = \Delta_b^2 + (P_{p,q}^{(j)})^* P_{p,q}^{(j)} \quad \text{on } \Lambda_{\pm}^{p,q} \text{ with } \sup(p, q) \geq 1,$$

$$(3.26) \quad \Delta_R = (\Delta_b + iT)^2 \quad \text{on } \Lambda_j^{n,0},$$

$$(3.27) \quad \Delta_R = (\Delta_b - iT)^2 \quad \text{on } \Lambda_j^{0,n}.$$

These formulas enabled Rumin [Ru] to show that Δ_R satisfied at every point a condition, the Rockland condition [Ro1], which then allowed him to apply results of Helffer-Nourrigat ([HN3]) to show that Δ_R was maximal hypoelliptic. Alternatively, the facts that the Rockland condition is

satisfied at every point and that the manifold is contact insure us that Δ_R admits a parametrix within the Heisenberg calculus, hence is hypoelliptic (see [Po8]).

4. HEISENBERG CALCULUS

In this section we gather the main facts about the Heisenberg calculus, following closely the expositions in [BG] and [Po8] (see also [EMM], [Tay]).

4.1. Left-invariant pseudodifferential operators. Let (M^{d+1}, H) be a Heisenberg manifold and let $G = G_a M$ be the tangent Lie group of M at a point $a \in M$. We review here the main results about left-invariant pseudodifferential operators on G (see also [BG], [CGGP], [Tay]).

Recall that for any finite dimensional vector space E the Schwartz class $\mathcal{S}(E)$ is a Fréchet space and the Fourier transform is the continuous isomorphism of $\mathcal{S}(E)$ onto $\mathcal{S}(E^*)$ given by

$$(4.1) \quad \hat{f}(\xi) = \int_E e^{i\langle \xi, x \rangle} f(x) dx, \quad f \in \mathcal{S}(E), \quad \xi \in E^*,$$

where dx denotes the Lebesgue measure of E .

Definition 4.1. $\mathcal{S}_0(E)$ is the closed subspace of $\mathcal{S}(E)$ consisting of $f \in \mathcal{S}(E)$ such that for any differential operator P on E^* we have $(P\hat{f})(0) = 0$.

Since G has the same underlying set as that of its Lie algebra $\mathfrak{g} = \mathfrak{g}_x M$ we can let $\mathcal{S}(G)$ and $\mathcal{S}_0(G)$ denote the Fréchet spaces $\mathcal{S}(E)$ and $\mathcal{S}_0(E)$ associated to the underlying linear space E of $\mathfrak{g}$ (notice that the Lebesgue measure of E coincides with the Haar measure of G since G is nilpotent).

Next, for $\lambda \in \mathbb{R}$ and $\xi = \xi_0 + \xi'$ in $\mathfrak{g}^* = (T_a^* M / H_a^*) \oplus H_a$ we let

$$(4.2) \quad \lambda \cdot \xi = \lambda \cdot (\xi_0 + \xi') = \lambda^2 \xi_0 + \lambda \xi'.$$

Definition 4.2. $S_m(\mathfrak{g}^*)$, $m \in \mathbb{C}$, is the space of functions $p \in C^\infty(\mathfrak{g}^* \setminus 0)$ which are homogeneous of degree m , in the sense that, for any $\lambda > 0$ we have

$$(4.3) \quad p(\lambda \cdot \xi) = \lambda^m p(\xi) \quad \forall \xi \in \mathfrak{g}^* \setminus 0.$$

In addition $S_m(\mathfrak{g}^*)$ is endowed with the Fréchet space topology induced from that of $C^\infty(\mathfrak{g}^* \setminus 0)$.

Note that the image $\hat{\mathcal{S}}_0(G)$ of $\mathcal{S}(G)$ under the Fourier transform consists of functions $v \in \mathcal{S}(\mathfrak{g}^*)$ such that, given any norm $|\cdot|$ on G , near $\xi = 0$ we have $|g(\xi)| = O(|\xi|^N)$ for any integer $N \geq 0$. Thus, any $p \in S_m(\mathfrak{g}^*)$ defines an element of $\hat{\mathcal{S}}_0(\mathfrak{g}^*)'$ by letting

$$(4.4) \quad \langle p, g \rangle = \int_{\mathfrak{g}^*} p(\xi) g(\xi) d\xi, \quad g \in \hat{\mathcal{S}}_0(\mathfrak{g}^*).$$

This allows us to define the inverse Fourier transform of p as the element $\check{p} \in \mathcal{S}_0(G)'$ such that

$$(4.5) \quad \langle \check{p}, f \rangle = \langle p, \check{f} \rangle \quad \forall f \in \mathcal{S}_0(G).$$

Proposition 4.3 ([BG], [CGGP]). 1) For any $p \in S_m(\mathfrak{g}^*)$ the left-convolution by $\check{p}$,

$$(4.6) \quad \check{p} * f(x) := \langle \check{p}(y), f(x \cdot y^{-1}) \rangle, \quad f \in \mathcal{S}_0(G),$$

defines a continuous endomorphism of $\mathcal{S}_0(G)$.

2) There is a continuous bilinear product,

$$(4.7) \quad * : S_{m_1}(\mathfrak{g}^*) \times S_{m_2}(\mathfrak{g}^*) \longrightarrow S_{m_1+m_2}(\mathfrak{g}^*),$$

such that, for any $p_1 \in S_{m_1}(\mathfrak{g}^*)$ and $p_2 \in S_{m_2}(\mathfrak{g}^*)$, the composition of the left-convolution operators by $\check{p}_1$ and $\check{p}_2$ is the left-convolution operator by $(p_1 * p_2)^\vee$, that is, we have

$$(4.8) \quad \check{p}_1 * (\check{p}_2 * f) = (p_1 * p_2)^\vee * f \quad \forall f \in \mathcal{S}_0(G).$$

Let us also mention that if $p \in S_m(\mathfrak{g}^*)$ then the convolution operator $Pu = \check{p} * f$ is a pseudodifferential operator. Indeed, let $X_0(a), \dots, X_d(a)$ be a (linear) basis of $\mathfrak{g}$ so that $X_0(a)$ is in $T_a M/H_a$ and $X_1(a), \dots, X_d(a)$ span H_a . For $j = 0, \dots, d$ let X_j^a be the left-invariant vector fields on G such that $X_j^w|_{x=0} = X_j(a)$. The basis $X_0(a), \dots, X_d(a)$ yields a linear isomorphism $\mathfrak{g} \simeq \mathbb{R}^{d+1}$, hence a global chart of G . In this chart p is a homogeneous symbol on $\mathbb{R}^{d+1} \setminus 0$ with respect to the dilations

$$(4.9) \quad \lambda.x = (\lambda^2 x_0, \lambda x_1, \dots, \lambda x_d), \quad x \in \mathbb{R}^{d+1}, \quad \lambda > 0.$$

Similarly, each vector fields $\frac{1}{i}X_j^a$, $j = 0, \dots, d$, corresponds to a vector fields on $\mathbb{R}^{d+1}$ whose symbol is denoted $\sigma_j^a(x, \xi)$. Then, setting $\sigma = (\sigma_0, \dots, \sigma_d)$, it can be shown that in the above chart P is the operator

$$(4.10) \quad Pf(x) = \int_{\mathbb{R}^{d+1}} e^{ix \cdot \xi} p(\sigma^a(x, \xi)) \hat{f}(\xi), \quad f \in \mathcal{S}_0(\mathbb{R}^{d+1}).$$

In other words P is the pseudodifferential operator $p(-iX^a) := p(\sigma^a(x, D))$ acting on $\mathcal{S}_0(\mathbb{R}^{d+1})$.

4.2. Ψ_H DO's on an open subset of $\mathbb{R}^{d+1}$. Let U be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $H \subset TU$ and a H -frame $X_0, X_1, \dots, X_d$ of TU . Then the class of Ψ_H DO's on U is a class of pseudodifferential operators modelled on that of homogeneous convolution operators on the fibers of GU .

Definition 4.4. $S_m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, is the space of symbols $p(x, \xi) \in C^\infty(U \times \mathbb{R}^{d+1} \setminus 0)$ that are homogeneous of degree m with respect to the ξ -variable, that is,

$$(4.11) \quad p(x, \lambda.\xi) = \lambda^m p(x, \xi) \quad \text{for any } \lambda > 0,$$

where $\xi \rightarrow \lambda.\xi$ denotes the Heisenberg dilation (4.9).

Observe that the homogeneity of $p \in S_m(U \times \mathbb{R}^{d+1})$ implies that, for any compact $K \subset U$, it satisfies estimates

$$(4.12) \quad |\partial_x^\alpha \partial_\xi^\beta p(x, \xi)| \leq C_{K\alpha\beta} \|\xi\|^{\Re m - \langle \beta \rangle}, \quad x \in K, \quad \xi \neq 0,$$

where $\|\xi\| = (|\xi_0|^2 + |\xi_1|^4 + \dots + |\xi_d|^4)^{1/4}$ and $\langle \alpha \rangle = 2\alpha_0 + \alpha_1 + \dots + \alpha_d$.

Definition 4.5. $S^m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, consists of symbols $p \in C^\infty(U \times \mathbb{R}^{d+1})$ with an asymptotic expansion $p \sim \sum_{j \geq 0} p_{m-j}$, $p_k \in S_k(U \times \mathbb{R}^{d+1})$, in the sense that, for any integer N and for any compact $K \subset U$, we have

$$(4.13) \quad |\partial_x^\alpha \partial_\xi^\beta (p - \sum_{j < N} p_{m-j})(x, \xi)| \leq C_{\alpha\beta NK} \|\xi\|^{\Re m - \langle \beta \rangle - N}, \quad x \in K, \quad \|\xi\| \geq 1.$$

Next, for $j = 0, \dots, d$ let $\sigma_j(x, \xi)$ denote the symbol of $\frac{1}{i}X_j$ (in the classical sense) and set $\sigma = (\sigma_0, \dots, \sigma_d)$. For any $p \in S^m(U \times \mathbb{R}^{d+1})$ it can be shown that the symbol $p_\sigma(x, \xi) := p(x, \sigma(x, \xi))$ is in the Hörmander class of symbols of type $(\frac{1}{2}, \frac{1}{2})$ (see [BG, Prop. 10.22]). Therefore, we define a continuous linear operator from $C_c^\infty(U)$ to $C^\infty(U)$ by letting

$$(4.14) \quad p(x, -iX)f(x) = (2\pi)^{-(d+1)} \int e^{ix \cdot \xi} p(x, \sigma(x, \xi)) \hat{f}(\xi) d\xi, \quad f \in C_c^\infty(U).$$

In the sequel we let $\Psi^{-\infty}(U)$ denotes the class of smoothing operators, i.e. of operators given by smooth kernels.

Definition 4.6. $\Psi_H^m(U)$, $m \in \mathbb{C}$, consists of operators $P : C_c^\infty(U) \rightarrow C^\infty(U)$ of the form

$$(4.15) \quad P = p(x, -iX) + R,$$

with p in $S^m(U \times \mathbb{R}^{d+1})$, called the symbol of P , and R is smoothing.

The above definition of the symbol of P differs from that of [BG], since there the authors defined it to be $p_\sigma(x, \xi) = p(x, \sigma(x, \xi))$. Note also that p is unique modulo $S^{-\infty}(U \times \mathbb{R}^{d+1})$.

Lemma 4.7. *For $j = 0, 1, \dots$ let $p_{m-j} \in S_{m-j}(U \times \mathbb{R}^{d+1})$. Then there exists $P \in \Psi_H^m(U)$ with symbol $p \sim \sum_{j \geq 0} p_{m-j}$. Moreover, the operator P is unique modulo smoothing operators.*

The class $\Psi_H^m(U)$ does not depend on the choice of the H -frame $X_0, \dots, X_d$ (see [BG, Prop. 10.46]). Moreover, since it is contained in the class of Ψ DO's of type $(\frac{1}{2}, \frac{1}{2})$ we get:

Proposition 4.8. *Let P be a Ψ_H DO of order m on U .*

1) P extends to a continuous mapping from $\mathcal{E}'(U)$ to $\mathcal{D}'(U)$ and has a distribution kernel which is smooth off the diagonal.

3) Let $k = \Re m$ if $\Re m \geq 0$ and $k = \frac{1}{2}\Re m$ otherwise. Then for any $s \in \mathbb{R}$ the operator $P \in \Psi_H^m(U)$ extends to a continuous mapping from $L_{s, \text{comp}}^2(U)$ to $L_{s-k, \text{loc}}^2(U)$.

4.3. Composition of Ψ_H DO's. Recall that there is no symbolic calculus for Ψ DO's of type $(\frac{1}{2}, \frac{1}{2})$ since the product of two such Ψ DO's needs not be again a Ψ DO of type $(\frac{1}{2}, \frac{1}{2})$. However, using the fact that the Ψ_H DO's are modelled on left-invariant pseudodifferential operators allows us to construct a symbolic calculus for Ψ_H DO's.

First, for $j = 0, \dots, d$ let $X_j^{(x)}$ be the leading homogeneous part of X_j in privileged coordinates centered at x defined according to (2.20) and (2.21). These vectors span a nilpotent Lie algebra of left-invariant vector fields on a nilpotent graded Lie group G^x which corresponds to $G_x U$ by pulling back the latter from the Heisenberg coordinates at x to the privileged coordinates at x .

As alluded to above the product law of $G^{(x)}$ defines a convolution product for symbols,

$$(4.16) \quad *^{(x)} : S_{m_1}(\mathbb{R}^{d+1}) \times S_{m_2}(\mathbb{R}^{d+1}) \longrightarrow S_{m_1+m_2}(\mathbb{R}^{d+1}).$$

such that, with the notations of (4.10), on $\mathcal{L}(\mathcal{S}_0(\mathbb{R}^{d+1}))$ we have

$$(4.17) \quad p_1(-iX^{(x)})p_2(-iX^{(x)}) = (p_1 *^{(x)} p_2)(-iX^{(x)}) \quad \forall p_j \in S_{m_j}(\mathbb{R}^{d+1}).$$

As it turns out the product $*^{(x)}$ depends smoothly on x (see [BG, Prop. 13.33]). Therefore, we get a continuous bilinear product,

$$(4.18) \quad * : S_{m_1}(U \times \mathbb{R}^{d+1}) \times S_{m_2}(U \times \mathbb{R}^{d+1}) \rightarrow S_{m_1+m_2}(U \times \mathbb{R}^{d+1}),$$

$$(4.19) \quad p_1 * p_2(x, \xi) = (p_1(x, \cdot) *^{(x)} p_2(x, \cdot))(\xi), \quad p_j \in S_{m_j}(U \times \mathbb{R}^{d+1}).$$

Proposition 4.9 ([BG, Thm. 14.7]). *For $j = 1, 2$ let $P_j \in \Psi_H^{m_j}(U)$ have symbol $p_j \sim \sum_{k \geq 0} p_{j, m_j - k}$ and assume that one of these operators is properly supported. Then the operator $P = P_1 P_2$ is a Ψ_H DO of order $m_1 + m_2$ and has symbol $p \sim \sum_{k \geq 0} p_{m_1+m_2-k}$, with*

$$(4.20) \quad p_{m_1+m_2-k}(x, \xi) = \sum_{k_1+k_2 \leq k} \sum_{\alpha, \beta, \gamma, \delta}^{(k-k_1-k_2)} h_{\alpha\beta\gamma\delta}(x) (D_\xi^\delta p_{1, m_1-k_1}) * (\xi^\gamma \partial_x^\alpha \partial_\xi^\beta p_{2, m_2-k_2})(x, \xi),$$

where $\sum_{\alpha, \beta, \gamma, \delta}^{(l)}$ denotes the sum over the indices such that $|\beta| = |\gamma|$ and $|\alpha| + |\beta| \leq \langle \beta \rangle - \langle \gamma \rangle + \langle \delta \rangle = l$, and the functions $h_{\alpha\beta\gamma\delta}(x)$'s are polynomials in the derivatives of the coefficients of the vector fields $X_0, \dots, X_d$.

Remark 4.10. It follows from (4.19) that for any $x \in U$ the x -symbol $p_{1, m_1} * p_{2, m_2}(x, \cdot)$ depends only on $p_{m_1}(x, \cdot)$ and $p_{m_2}(x, \cdot)$. However, the value of $(p_{1, m_1} * p_{2, m_2})(x, \xi)$ at $(x, \xi) \in U \times \mathbb{R}^{d+1}$ depends on all the η -values $p_{m_1}(x, \eta)$ and $p_{m_2}(x, \eta)$ as η ranges over $\mathbb{R}^{d+1}$. Thus we may localize the product of Heisenberg symbols with respect to x , but not respect to (x, ξ) , that is, the product of Ψ_H DO's is local, but is not microlocal.

4.4. The distribution kernels of Ψ_{HDO} 's. An important fact about ΨDO 's is their characterization in terms of their distribution kernels.

First, we extend the notion of homogeneity of functions to distributions. For $K \in \mathcal{S}'(\mathbb{R}^{d+1})$ and $\lambda > 0$ we let K_λ denote the element of $\mathcal{S}'(\mathbb{R}^{d+1})$ such that

$$(4.21) \quad \langle K_\lambda, f \rangle = \lambda^{-(d+2)} \langle K(x), f(\lambda^{-1} \cdot x) \rangle \quad \forall f \in \mathcal{S}(\mathbb{R}^{d+1}).$$

In the sequel we will also use the notation $K(\lambda \cdot x)$ for denoting $K_\lambda(x)$. We then say that K is homogeneous of degree m , $m \in \mathbb{C}$, when $K_\lambda = \lambda^m K$ for any $\lambda > 0$.

Definition 4.11. $\mathcal{S}'_{\text{reg}}(\mathbb{R}^{d+1})$ consists of tempered distributions on $\mathbb{R}^{d+1}$ which are smooth outside the origin. We equip it with the weakest topology such that the inclusions of $\mathcal{S}'_{\text{reg}}(\mathbb{R}^{d+1})$ into $\mathcal{S}'(\mathbb{R}^{d+1})$ and $C^\infty(\mathbb{R}^{d+1} \setminus 0)$ are continuous.

Definition 4.12. $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, consists of distributions $K(x, y)$ in $C^\infty(U) \hat{\otimes} \mathcal{S}'_{\text{reg}}(\mathbb{R}^{d+1})$ such that for some functions $c_\alpha(x) \in C^\infty(U)$, $\langle \alpha \rangle = m$, we have

$$(4.22) \quad K(x, \lambda \cdot y) = \lambda^m K(x, y) + \lambda^m \log \lambda \sum_{\langle \alpha \rangle = m} c_\alpha(x) y^\alpha \quad \text{for any } \lambda > 0.$$

The interest of considering the distribution class $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$ stems from:

Lemma 4.13 ([BG, Prop. 15.24], [CM, Lem. I.4]). 1) Any $p \in S_m(U \times \mathbb{R}^{d+1})$ agrees on $U \times (\mathbb{R}^{d+1} \setminus 0)$ with a distribution $\tau(x, \xi)$ in $C^\infty(U) \hat{\otimes} \mathcal{S}'(\mathbb{R}^{d+1})$ such that $\tilde{\tau}_{\xi \rightarrow y}$ is in $\mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m + d + 2)$.

2) If $K(x, y)$ is in $\mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$ then the restriction of $\hat{K}_{y \rightarrow \xi}(x, \xi)$ to $U \times (\mathbb{R}^{d+1} \setminus 0)$ belongs to $S_m(U \times \mathbb{R}^{d+1})$.

This result is a consequence of the solution to the problem of extending a homogeneous function $p \in C^\infty(\mathbb{R}^{d+1} \setminus 0)$ into a homogeneous distribution on $\mathbb{R}^{d+1}$ and of the fact that for $\tau \in \mathcal{S}'(\mathbb{R}^{d+1})$ we have

$$(4.23) \quad (\hat{\tau})_\lambda = |\lambda|^{-(d+2)} (\tau_{\lambda^{-1}})^\wedge \quad \forall \lambda \in \mathbb{R} \setminus 0.$$

In particular, if τ is homogeneous of degree m then $\hat{\tau}$ is homogeneous of $-(m + d + 2)$.

The relevant class of kernels for the Heisenberg calculus is the following.

Definition 4.14. $\mathcal{K}^m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, consists of distributions $K \in \mathcal{D}'(U \times \mathbb{R}^{d+1})$ with an asymptotic expansion $K \sim \sum_{j \geq 0} K_{m+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, in the sense that, for any integer N , as soon as J is large enough we have

$$(4.24) \quad K - \sum_{j \leq J} K_{m+j} \in C^N(U \times \mathbb{R}^{d+1}).$$

Since under the Fourier transform the asymptotic expansion (4.13) for symbols corresponds to that for distributions in (4.24), using Lemma 4.13 we get:

Lemma 4.15 ([BG, pp. 133–134]). Let $K \in \mathcal{D}'(U \times \mathbb{R}^{d+1})$. Then the following are equivalent:

- (i) The distribution K belongs to $\mathcal{K}^m(U \times \mathbb{R}^{d+1})$;
- (ii) We can put K into the form

$$(4.25) \quad K(x, y) = \check{p}_{\xi \rightarrow y}(x, y) + R(x, y),$$

for some $p \in S^{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m + d + 2)$, and some $R \in C^\infty(U \times \mathbb{R}^{d+1})$.

Moreover, if (i) and (ii) holds and we expand $K \sim \sum_{j \geq 0} K_{m+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, then we have $p \sim \sum_{j \geq 0} p_{\hat{m}-j}$ where $p_{\hat{m}-j} \in S_{\hat{m}-j}(U \times \mathbb{R}^{d+1})$ is the restriction to $U \times (\mathbb{R}^{d+1} \setminus 0)$ of $(K_{m+j})_{y \rightarrow \xi}^\wedge$.

Next, for $x \in U$ let ψ_x denote the affine change to the privileged coordinates at x and let us write $(A_x^t)^{-1}\xi = \sigma(x, \xi)$ with $A_x \in \text{GL}_{d+1}(\mathbb{R})$. Since $\psi_x(x) = 0$ and $\psi_{x*}X_j = \frac{\partial}{\partial y_j}$ at $y = 0$ for $j = 0, \dots, d$, one checks that $\psi_x(y) = A_x(y - x)$.

Let $p \in S^m(U \times \mathbb{R}^{d+1})$. As $p(x, -iX) = p^\sigma(x, D)$ with $p^\sigma(x, \xi) = p(x, \sigma(x, \xi)) = p(x, (A_x^t)^{-1}\xi)$ the distribution kernel $k_{p(x, -iX)}(x, y)$ of $p(x, -iX)$ is represented by the oscillating integrals

$$(4.26) \quad (2\pi)^{-(d+1)} \int e^{i(x-y) \cdot \xi} p(x, (A_x^t)^{-1}\xi) d\xi = (2\pi)^{-(d+1)} |A_x| \int e^{iA_x(x-y) \cdot \xi} p(x, \xi) d\xi.$$

Since $\psi_x(y) = A_x(y - x)$ we deduce that

$$(4.27) \quad k_{p(x, -iX)}(x, y) = |\psi'_x| \check{p}_{\xi \rightarrow y}(x, -\psi_x(y)).$$

Combining this with Lemma 4.15 then gives:

Proposition 4.16 ([BG, Thms. 15.39, 15.49]). *Let $P : C_c^\infty(U) \rightarrow C^\infty(U)$ be a continuous linear operator with distribution kernel $k_P(x, y)$. Then the following are equivalent:*

- (i) P is a Ψ_H DO of order m , $m \in \mathbb{C}$.
- (ii) There exist $K \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m + d + 2)$, and $R \in C^\infty(U \times U)$ such that

$$(4.28) \quad k_P(x, y) = |\psi'_x| K(x, -\psi_x(y)) + R(x, y).$$

Furthermore, if (i) and (ii) hold and $K \sim \sum_{j \geq 0} K_{\hat{m}+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, then P has symbol $p \sim \sum_{j \geq 0} p_{m-j}$, $S_l \in S_l(U \times \mathbb{R}^{d+1})$, where $p_{\hat{m}-j}$ is the restriction to $U \times (\mathbb{R}^{d+1} \setminus 0)$ of $(K_{m+j})_{y \rightarrow \xi}^\wedge$.

In the sequel we will need a version of Proposition 4.16 in Heisenberg coordinates. To this end let ε_x denote the coordinate change to the Heisenberg coordinates at x and set $\phi_x = \varepsilon_x \circ \psi_x^{-1}$. Recall that ϕ_x is a Lie group isomorphism from $G^{(x)}$ to $G_x U$ such that $\phi_x(\lambda \cdot y) = \lambda \cdot \phi_x(y)$ for any $\lambda \in \mathbb{R}$. Moreover, using (2.25) one can check that $|\phi'_x| = 1$ and $\phi_x^{-1}(y) = -\phi_x(-y)$. Therefore, from (4.27) we see that we can put $k_{p(x, -iX)}(x, y)$ into the form

$$(4.29) \quad k_{p(x, -iX)}(x, y) = |\varepsilon'_x| K_P(x, -\varepsilon_x(y)), \quad K_P(x, y) = \check{p}_{\xi \rightarrow y}(x, -\phi_x(-y)) = \check{p}_{\xi \rightarrow y}(x, \phi_x^{-1}(y)).$$

In fact, the coordinate changes ϕ_x give rise to an action on distributions on $U \times \mathbb{R}^{d+1}$ given by

$$(4.30) \quad K(x, y) \longrightarrow \phi_x^* K(x, y), \quad \phi_x^* K(x, y) = K(x, \phi_x^{-1}(y)).$$

Since ϕ_x depends smoothly on x , this action induces a continuous linear isomorphisms of $C^N(U \times \mathbb{R}^{d+1})$, $N \geq 0$, and $C^\infty(U \times \mathbb{R}^{d+1})$ onto themselves. As $\phi_x(y)$ is polynomial in y in such way that $\phi_x(0) = 0$ and $\phi_x(\lambda \cdot y) = \lambda \cdot \phi_x(y)$ for every $\lambda \in \mathbb{R}$, we deduce that the above action also yields a continuous linear isomorphism of $C^\infty(U) \hat{\otimes} S'_{\text{reg}}(\mathbb{R}^{d+1})$ onto itself and, for every $\lambda > 0$, we have

$$(4.31) \quad (\phi_x^* K)(x, \lambda \cdot y) = \phi_x^*[K(x, \lambda \cdot y)], \quad K \in \mathcal{D}'(U \times \mathbb{R}^{d+1}).$$

Furthermore, as $\phi_x(y)$ is polynomial in y we see that for every $\alpha \in \mathbb{N}^{d+1}$ we can write $\phi_x(y)^\alpha$ in the form $\phi_x(y)^\alpha = \sum_{\langle \beta \rangle = \langle \alpha \rangle} d_{\alpha\beta}(x) y^\beta$ with $d_{\alpha\beta} \in C^\infty(U \times \mathbb{R}^{d+1})$. It then follows that, for every $m \in \mathbb{C}$, the map $K(x, y) \rightarrow \phi_x^* K(x, y)$ induces a linear isomorphisms of $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$ and $\mathcal{K}^m(U \times \mathbb{R}^{d+1})$ onto themselves. Combining this with (4.29) and Proposition 4.16 then gives:

Proposition 4.17. *Let $P : C_c^\infty(U) \rightarrow C^\infty(U)$ be a continuous linear operator with distribution kernel $k_P(x, y)$. Then the following are equivalent:*

- (i) P is a Ψ_H DO of order m , $m \in \mathbb{C}$.
- (ii) There exist $K_P \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m + d + 2)$, and $R \in C^\infty(U \times U)$ such that

$$(4.32) \quad k_P(x, y) = |\varepsilon'_x| K_P(x, -\varepsilon_x(y)) + R(x, y).$$

Furthermore, if (i) and (ii) hold and $K_P \sim \sum_{j \geq 0} K_{P, \hat{m}+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, then P has symbol $p \sim \sum_{j \geq 0} p_{m-j}$, $S_l \in S_l(U \times \mathbb{R}^{d+1})$, where $p_{\hat{m}-j}$ is the restriction to $U \times (\mathbb{R}^{d+1} \setminus 0)$ of $[K_{P, \hat{m}+j}(x, \phi_x^{-1}(y))]_{y \rightarrow \xi}^\wedge$.

Remark 4.18. Let $a \in U$. Then from (4.32) we see that the distribution kernel of $\tilde{P} = (\varepsilon_a)_* P$ at $x = 0$ is

$$(4.33) \quad k_{\tilde{P}}(0, y) = |\varepsilon'_a|^{-1} k_P(\varepsilon_a^{-1}(0), \varepsilon_a^{-1}(y)) = K_P(a, -y).$$

On the other hand, as we are in Heisenberg coordinates already, we have $\psi_0 = \varepsilon_0 = \phi_0 = \text{id}$. Thus, in the form (4.32) for $\tilde{P}$ we have $K_{\tilde{P}}(0, y) = K_P(a, y)$. Therefore, if we let $p_m(x, \xi)$ denote the principal symbol of P and let $K_{P, \hat{m}} \in \mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$ denote the leading kernel of K_P then by Proposition 4.17 we have

$$(4.34) \quad p_m(0, \xi) = [K_{P, \hat{m}}]_{y \rightarrow \xi}^\wedge(a, \xi),$$

This shows that $[K_{P, \hat{m}}]_{y \rightarrow \xi}^\wedge(a, \xi)$ is the principal symbol of P at $x = 0$ in Heisenberg coordinates centered at a .

4.5. Ψ_H DO's on a general Heisenberg manifold. Let (M^{d+1}, H) be a Heisenberg manifold. As alluded to before the Ψ_H DO's on an subset of $\mathbb{R}^{d+1}$ are Ψ DO's of type $(\frac{1}{2}, \frac{1}{2})$. However, the latter don't make sense on a general manifold, for their class is not preserved by an arbitrary change of chart. Nevertheless, when dealing with Ψ_H DO's this issue is resolved if we restrict ourselves to changes of Heisenberg charts. Indeed, we have:

Proposition 4.19 ([BG], [Po8]). *Let U (resp. $\tilde{U}$) be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $H \subset TU$ (resp. $\tilde{H} \subset T\tilde{U}$) and a H -frame of TU (resp. a $\tilde{H}$ -frame of $T\tilde{U}$). Let $\phi : (U, H) \rightarrow (\tilde{U}, \tilde{H})$ be a Heisenberg diffeomorphism and let $\tilde{P} \in \Psi_{\tilde{H}}^m(\tilde{U})$.*

1) *The operator $P = \phi^* \tilde{P}$ is a Ψ_H DO of order m on U .*

2) *If the distribution kernel of $\tilde{P}$ is of the form (4.32) with $K_{\tilde{P}}(\tilde{x}, \tilde{y}) \in \mathcal{K}^{\hat{m}}(\tilde{U} \times \mathbb{R}^{d+1})$ then the distribution kernel of P can be written in the form (4.32) with $K_P(x, y) \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ such that*

$$(4.35) \quad K_P(x, y) \sim \sum_{\langle \beta \rangle \geq \frac{3}{2} \langle \alpha \rangle} \frac{1}{\alpha! \beta!} a_{\alpha\beta}(x) y^\beta (\partial_y^\alpha K_{\tilde{P}})(\phi(x), \phi'_H(x)y),$$

where we have let $a_{\alpha\beta}(x) = \partial_y^\beta [|\partial_y(\tilde{\varepsilon}_{\phi(x)} \circ \phi \circ \varepsilon_x^{-1})(y)|(\tilde{\varepsilon}_{\phi(x)} \circ \phi \circ \varepsilon_x^{-1}(y) - \phi'_H(x)y)^\alpha]_{|y=0}$ and $\tilde{\varepsilon}_{\tilde{x}}$ denote the change to the Heisenberg coordinates at $\tilde{x} \in \tilde{f}$. In particular, we have

$$(4.36) \quad K_P(x, y) = |\phi'_H(x)| K_{\tilde{P}}(\phi(x), \phi'_H(x)y) \mod \mathcal{K}^{\hat{m}+1}(U \times \mathbb{R}^{d+1}).$$

As a consequence of Proposition 4.19 we can define Ψ_H DO's on M acting on the sections of a vector bundle $\mathcal{E}$ over M .

Definition 4.20. $\Psi_H^m(M, \mathcal{E})$, $m \in \mathbb{C}$, consists of continuous operators $P : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ such that:

(i) *The distribution kernel of P is smooth off the diagonal;*

(ii) *For any trivialization $\tau : \mathcal{E}|_U \rightarrow U \times \mathbb{C}^r$ over a local Heisenberg chart $\kappa : U \rightarrow V \subset \mathbb{R}^{d+1}$ the operator $\kappa_* \tau_*(P|_f)$ belongs to $\Psi_H^m(V, \mathbb{C}^r) := \Psi_H^m(V) \otimes \text{End } \mathbb{C}^r$.*

All the aforementioned properties of Ψ DO's on an open subset of $\mathbb{R}^{d+1}$ hold *mutatis standis* in this setting.

4.6. Transposes and adjoints of Ψ_H DO's. Let us now look at the transpose and adjoints of Ψ_H DO's. First, given a Heisenberg chart $U \subset \mathbb{R}^{d+1}$ we have:

Proposition 4.21 ([BG], [Po8]). *Let $P \in \Psi_H^m(U)$. Then:*

- 1) *The transpose operator P^t is a Ψ_H DO of order m on U .*
- 2) *If we write the distribution kernel of P in the form (4.32) with $K_P \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ then P^t can be written in the form (4.32) with $K_{P^t} \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ such that*

$$(4.37) \quad K_{P^t}(x, y) \sim \sum_{\frac{3}{2} \langle \alpha \rangle \leq \langle \beta \rangle} \sum_{|\gamma| \leq |\delta| \leq 2|\gamma|} a_{\alpha\beta\gamma\delta}(x) y^{\beta+\delta} (\partial_x^\gamma \partial_y^\alpha K_P)(x, -y),$$

where $a_{\alpha\beta\gamma\delta}(x) = \frac{|\varepsilon_x^{-1}|}{\alpha! \beta! \gamma! \delta!} [\partial_y^\beta (|\varepsilon_x^{-1}(-y)| (y - \varepsilon_x^{-1}(y)(x))^\alpha) \partial_y^\delta (\varepsilon_x^{-1}(-y) - x)^\gamma](x, 0)$. In particular,

$$(4.38) \quad K_{P^t}(x, y) = K_P(x, -y) \quad \text{mod } \mathcal{K}^{\hat{m}+1}(U \times \mathbb{R}^{d+1}).$$

Thanks to this result we can prove:

Proposition 4.22 ([BG], [Po8]). *Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ_H DO of order m . Then:*

- 1) *The transpose operator $P^t : \mathcal{E}'(M, \mathcal{E}^*) \rightarrow \mathcal{D}'(M, \mathcal{E}^*)$ is a Ψ_H DO of order m ;*
- 2) *If M is endowed with a smooth positive density and $\mathcal{E}$ with a Hermitian metric then the adjoint $P^* : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ is a Ψ_H DO of order $\bar{m}$.*

4.7. Principal symbol and model operators. Let us now give an intrinsic definition of the principal symbol of a Ψ_H DO and of its model operator at a point.

Let $\pi : \mathfrak{g}^*M \rightarrow M$ be the canonical projection of the bundle $\mathfrak{g}^*M$ onto M .

Definition 4.23. $S_m(\mathfrak{g}^*M, \mathcal{E})$, $m \in \mathbb{C}$, is the space of sections $p \in C^\infty(\mathfrak{g}^*M \setminus 0, \text{End } \pi^*\mathcal{E})$ which are homogeneous of degree m in the sense that, for any $\lambda > 0$, we have

$$(4.39) \quad p(x, \lambda\xi) = \lambda^m p(x, \xi) \quad \forall (x, \xi) \in \mathfrak{g}^*M \setminus 0,$$

where $\xi \rightarrow \lambda\xi$ denotes the dilation (4.2).

Let $P \in \Psi_H^m(M)$ and for $j = 1, 2$ let κ_j be a Heisenberg chart with domain $V_j \subset M$. We let $\phi : U_1 \rightarrow U_2$ be the corresponding transition map, with $U_j = \kappa_j(V_1 \cap V_2) \subset \mathbb{R}^{d+1}$, and for $j = 1, 2$ we define $P_j := \kappa_{j*}(P|_{V_1 \cap V_2})$, so that $P_1 = \phi^*P_2$. Since P_j belongs to $\Psi_H^m(U_j)$ its distribution kernel is of the form (4.32) with $K_{P_j} \in \mathcal{K}^{\hat{m}}(U_j \times \mathbb{R}^{d+1})$. Let $K_{P_j, \hat{m}} \in \mathcal{K}_{\hat{m}}(U_j \times \mathbb{R}^{d+1})$ be the leading kernel of K_{P_j} and define

$$(4.40) \quad p_{j,m}(x, \xi) = [K_{P_j, \hat{m}}]_{y \rightarrow \xi}^\wedge(x, \xi), \quad (x, \xi) \in U_j \times \mathbb{R}^{d+1} \setminus 0.$$

Recall that by Remark 4.18 for any $a \in U_j$ the symbol $p_j(a, \cdot)$ yields in Heisenberg coordinates centered at a the principal symbol of P_j at $x = 0$. Moreover, as shown in [Po8], it follows from Proposition 4.19 that we have

$$(4.41) \quad p_{1,m}(x, \xi) = p_{2,m}(\phi(x), [\phi'_H(x)]^{-1t}\xi).$$

This shows that $p_m := \kappa_1^* p_{1,m}$ is an element of $S_m(\mathfrak{g}^*(V_1 \cap V_2))$ which is independent of the choice of the chart κ_1 . Since $S_m(\mathfrak{g}^*M)$ is a sheaf this gives rise to a uniquely defined symbol $\sigma_m(P) \in S_m(\mathfrak{g}^*M)$. As we can similarly deal with Ψ_H DO's acting on $\mathcal{E}$ we obtain:

Proposition 4.24 ([Po8]). *For any $P \in \Psi_H^m(M, \mathcal{E})$ there is a unique symbol $\sigma_m(P) \in S_m(\mathfrak{g}^*M, \mathcal{E})$ such that if in a local trivializing Heisenberg chart $U \subset \mathbb{R}^{d+1}$ we let $K_{P, \hat{m}}(x, y) \in \mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$ be the leading kernel for the kernel $K_P(x, y)$ in the form (4.32) for P , then we have*

$$(4.42) \quad \sigma_m(P)(x, \xi) = [K_{P, \hat{m}}]_{y \rightarrow \xi}^\wedge(x, \xi), \quad (x, \xi) \in U \times \mathbb{R}^{d+1} \setminus 0.$$

Equivalently, for any $x_0 \in M$ the symbol $\sigma_m(P)(x_0, \cdot)$ agrees in trivializing Heisenberg coordinates centered at x_0 with the principal symbol of P at $x = 0$.

Definition 4.25. For $P \in \Psi_H^m(M, \mathcal{E})$ the symbol $\sigma_m(P) \in S_m(\mathfrak{g}^*M, \mathcal{E})$ provided by Proposition 4.24 is called the principal symbol of P .

Since we have two notions of principal symbol we shall distinguish between them by saying that $\sigma_m(P)$ is the global principal symbol of P and that in a local trivializing chart the principal symbol p_m of P in the sense of (4.13) is the local principal symbol of P in this chart.

In fact, as shown in [Po8] the symbols $p_m(x, \xi)$ and $\sigma_m(P)(x, \xi)$ are related by

$$(4.43) \quad p_m(x, \xi) = (\hat{\phi}_x^* \sigma_m(P))(x, \xi),$$

$$(4.44) \quad (\hat{\phi}_x^* \sigma_m(P)) = [[\sigma_m(P)]_{\xi \rightarrow y}^\vee(x, \phi_x^{-1}(y))]_{y \rightarrow \xi}^\wedge = [\phi_x^* [\sigma_m(P)]_{\xi \rightarrow y}^\vee]_{y \rightarrow \xi}^\wedge,$$

where ϕ_x^* is the isomorphism map (4.30). In particular, since the latter is a linear isomorphism of $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$ onto itself the map $p \rightarrow \hat{\phi}_x^* p$ is a linear isomorphism of $S_m(U \times \mathbb{R}^{d+1})$ onto itself.

On the other hand, the principal map is surjective. More precisely, we have:

Proposition 4.26 ([Po8]). For every $m \in \mathbb{C}$ the principal symbol map $\sigma_m : \Psi_H^m(M, \mathcal{E}) \rightarrow S_m(\mathfrak{g}^*M, \mathcal{E})$ gives rise to a linear isomorphism $\Psi_H^m(M, \mathcal{E}) / \Psi_H^{m-1}(M, \mathcal{E}) \xrightarrow{\sim} S_m(\mathfrak{g}^*M, \mathcal{E})$.

Now, granted the above definition of the principal symbol, we can define the model operator at a point as follows.

Definition 4.27. Let $P \in \Psi_H^m(M, \mathcal{E})$ have (global) principal symbol $\sigma_m(P)$. Then the model operator of P at $a \in M$ is the left-invariant Ψ_H DO-operator $P^a : \mathcal{S}_0(G_a M, \mathcal{E}_a) \rightarrow \mathcal{S}_0(G_a M, \mathcal{E}_a)$ with symbol $\sigma_m(P)_{\xi \rightarrow y}^\vee(a, \cdot)$, so that we have

$$(4.45) \quad P^a f(x) = \langle \sigma_m(P)_{\xi \rightarrow y}^\vee(a, y), f(x \cdot y^{-1}) \rangle, \quad f \in \mathcal{S}_0(G_a M, \mathcal{E}_a).$$

For $a \in M$ we let $*^a : S_{m_1}(G_a M) \times S_{m_2}(G_a M) \rightarrow S_{m_1+m_2}(G_a M)$ be the convolution product for symbols defined by the product law of $G_a M$. Then it is proved in [Po8] that in a local trivializing chart we have

$$(4.46) \quad p_{m_1} *^a p_{m_2} = (\hat{\phi}_a)_* [(\hat{\phi}_a^* p_{m_1}) *^{(a)} (\hat{\phi}_a^* p_{m_2})] \quad \forall p_{m_j} \in S_{m_j}(\mathbb{R}^{d+1}),$$

where $(\hat{\phi}_a)_*$ denotes the inverse of $\hat{\phi}_a^*$. Since $\hat{\phi}_a^*$, its inverse and $*^{(a)}$ depend smoothly on a , we deduce that so does $*^a$. This allows us to get:

Proposition 4.28 ([Po8]). 1) The group laws on the fibers of GM give rise to a convolution product,

$$(4.47) \quad * : S_{m_1}(\mathfrak{g}^*M, \mathcal{E}) \times S_{m_2}(\mathfrak{g}^*M, \mathcal{E}) \longrightarrow S_{m_1+m_2}(\mathfrak{g}^*M, \mathcal{E}),$$

such that for symbols $p_{m_j} \in S_{m_j}(\mathfrak{g}^*M, \mathcal{E})$, $j = 1, 2$, we have

$$(4.48) \quad p_{m_1} * p_{m_2}(x, \xi) = [p_{m_1}(x, \cdot) *^x p_{m_2}(x, \cdot)](\xi), \quad (x, \xi) \in \mathfrak{g}^*M \setminus 0,$$

where $*^x$ denote the convolution product for symbols on $G_x M$.

Proposition 4.29 ([Po8]). For $j = 1, 2$ let $P_j \in \Psi_H^{m_j}(M, \mathcal{E})$ and suppose that P_1 or P_2 is properly supported. Then:

- 1) We have $\sigma_{m_1+m_2}(P_1 P_2) = \sigma_{m_1}(P) * \sigma_{m_2}(P)$.
- 2) At every point $a \in M$ the model operator of $P_1 P_2$ is $(P_1 P_2)^a = P_1^a P_2^a$.

Using Proposition 4.21 we can also deal with the principal symbols and model operators of the adjoints and transposes of Ψ_H DO's.

Proposition 4.30 ([BG], [Po8]). Let $P \in \Psi_H^m(M, \mathcal{E})$ have principal symbol $\sigma_m(P)$. Then:

- 1) The principal symbol of the transpose P^t is $\sigma_m(P^t)(x, \xi) = \sigma_m(x, -\xi)^t$;
- 2) If P^a is the model operator of P at a , then the model operator of P^t at a is the transpose operator $(P^a)^t : \mathcal{S}_0(G_x M, \mathcal{E}_x^*) \rightarrow \mathcal{S}_0(G_x M, \mathcal{E}_x^*)$.

Assume now that M and $\mathcal{E}$ are endowed with a positive density and a Hermitian metric respectively and let $L^2(M, \mathcal{E})$ be the associated L^2 -Hilbert space.

Proposition 4.31 ([Po8]). *Let $P \in \Psi_H^m(M, \mathcal{E})$ have principal symbol $\sigma_m(P)$. Then:*

- 1) *The principal symbol of the adjoint P^* is $\sigma_m(P^*)(x, \xi) = \sigma_m(P)(x, \xi)^*$.*
- 2) *If P^a denotes the model operator of P at $a \in M$ then the model operator of P^* at a is the adjoint $(P^a)^*$ of P^a .*

4.8. Hypoellipticity, parametrices and Rockland condition. Let $P : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ_H DO of order m such that $k := \Re m > 0$. First, we have:

Proposition 4.32 ([BG], [Po8]). *The following are equivalent:*

- 1) *The principal symbol $\sigma_m(P)$ of P is invertible with respect to the convolution product for homogeneous symbols;*
 - 2) *The operator P admits a parametrix Q in $\Psi_H^{-m}(M, \mathcal{E})$, i.e. $PQ = QP = 1 \bmod \Psi^{-\infty}(M, \mathcal{E})$.*
- Furthermore, if 1) and 2) hold then P is hypoelliptic with loss of $\frac{k}{2}$ -derivatives, i.e., for any $s \in \mathbb{R}$ and any compact $K \subset M$ we have estimates*

$$(4.49) \quad \|f\|_{L_{s+k/2}^2} \leq C_K (\|Pf\|_{H^s} + \|f\|_{L_s^2}) \quad \forall f \in C_K^\infty(M, \mathcal{E}).$$

When M is compact, combining this with the compactness of the embedding of $H^{k/2}(M, \mathcal{E})$ into $L^2(M, \mathcal{E})$ we get:

Proposition 4.33. *Suppose M compact and assume that P has an invertible principal symbol and a spectrum different from $\mathbb{C}$. Then:*

- 1) *The spectrum of P consists of isolated eigenvalues with finite multiplicities.*
- 2) *For any $\lambda \in \text{Sp } P$ the eigenspace $\ker(P - \lambda)$ is a finite dimensional subspace of $C^\infty(M, \mathcal{E})$.*

Now, assume that M is endowed with a positive density and $\mathcal{E}$ with a Hermitian metric. Let P^a be the model operator of P at a point $a \in M$ and let $\pi : G \rightarrow \mathcal{H}_\pi$ be a (nontrivial) unitary representation of $G = G_a M$. We define the symbol π_{P^a} as follows (see also [Ro1], [G1], [CGGP]).

Let $\mathcal{H}_\pi^0(\mathcal{E}_a)$ be the subspace of $\mathcal{H}_\pi(\mathcal{E}_a) := \mathcal{H}_\pi \otimes \mathcal{E}_a$ spanned by the vectors of the form

$$(4.50) \quad \pi_f \xi = \int_G (\pi_x \otimes 1_{\mathcal{E}_a})(\xi \otimes f(x)) dx, \quad f \in \mathcal{S}_0(G), \quad \eta \in \mathcal{H}_\pi(\mathcal{E}_a).$$

with f in $\mathcal{S}_0(G, \mathcal{E}_a) = \mathcal{S}_0(G)(\mathcal{E}_a)$ and $\xi \in \mathcal{H}_\pi$. Then we let π_{P^a} denote the (unbounded) operator of $\mathcal{H}_\pi(\mathcal{E}_a)$ with domain $\mathcal{H}_\pi^0(\mathcal{E}_a)$ such that

$$(4.51) \quad \pi_{P^a}(\pi_f \xi) = \pi_{P^a f} \xi \quad \forall f \in \mathcal{S}_0(G, \mathcal{E}_a) \quad \forall \xi \in \mathcal{H}_\pi.$$

One can check that $\pi_{P^a}^*$ is the adjoint of π_{P^a} on $\mathcal{H}_\pi^0$, hence is densely defined. Thus π_{P^a} is closable and we can let $\overline{\pi_{P^a}}$ denotes its closure.

In the sequel we let $C_\pi^\infty(\mathcal{E}_a) = C_\pi^\infty \otimes \mathcal{E}_a$, where $C_\pi^\infty \subset \mathcal{H}_\pi$ denotes the space of smooth vectors of π (i.e. the subspace of vectors $\xi \in \mathcal{H}_\pi$ so that $x \rightarrow \pi(x)\xi$ is smooth from G to $\mathcal{H}_\pi$).

Proposition 4.34 ([CGGP]). 1) *The domain of $\overline{\pi_{P^a}}$ always contains $C_\pi^\infty(\mathcal{E}_a)$.*

2) *If $\Re m \leq 0$ then the operator $\overline{\pi_{P^a}}$ is bounded.*

3) *We have $\overline{(\pi_{P^a})^*} = (\overline{\pi_{P^a}})^*$.*

4) *If P_1 and P_2 are Ψ DO's on M then $\overline{\pi_{(P_1 P_2)^a}} = \overline{\pi_{P_1^a}} \overline{\pi_{P_2^a}}$.*

Remark 4.35. If $\mathcal{E}_a = \mathbb{C}$ and P^a is a differentiable operator then, as it is left-invariant, P^a belongs to the enveloping algebra $\mathcal{U}(\mathfrak{g})$ of the Lie algebra $\mathfrak{g} = \mathfrak{g}_a M$ of G . In this case $\overline{\pi_{P^a}}$ coincides on C_π^∞ with the operator $d\pi(P^a)$, where $d\pi$ is the representation of $\mathcal{U}(\mathfrak{g})$ induced by π .

Definition 4.36. We say that P satisfies the Rockland condition at a if for any nontrivial unitary irreducible representation π of $G_a M$ the operator $\overline{\pi P^a}$ is injective on $C_\pi^\infty(\mathcal{E}_a)$.

Proposition 4.37 ([Po8]). Suppose that the Levi form of (M, H) has constant rank. Then the following are equivalent:

- (i) P and P^t satisfy the Rockland condition at every point of M ;
- (ii) P and P^* satisfy the Rockland condition at every point of M ;
- (iii) The principal symbol of P is invertible.

In particular, if P is selfadjoint then principal symbol of P is invertible if, and only if, P satisfies the Rockland condition at every point of M . In any case, if the conditions (i), (ii), (iii) hold P admits a parametrix in $\Psi_H^{-m}(M, \mathcal{E})$ and is hypoelliptic in the sense of (4.49).

Finally, for a sublaplacian $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ the Rockland condition can be reformulated as follows. Near a point $a \in M$ let $X_0, X_1, \dots, X_d$ be a local H -frame of TM with respect to which Δ takes the form,

$$(4.52) \quad \Delta = -(X_1^2 + \dots + X_d^2) - i\mu(x)X_0 + O_H(1),$$

where $\mu(x)$ is a local section of $\text{End } \mathcal{E}$ and the notation $O_H(1)$ means a differential operator of Heisenberg order ≤ 1 .

Let $L(x) = (L_{jk}(x))$ be the matrix of $\mathcal{L}$ with respect to the H -frame $X_0, X_1, \dots, X_d$, so that for $j, k = 1, \dots, d$ we have

$$(4.53) \quad \mathcal{L}(X_j, X_k) = [X_j, X_k] = L_{jk}(x)X_0 \mod H.$$

Let $2n$ be the rank of $\mathcal{L}_a$ and $L(a)$, let $\lambda_1, \dots, \lambda_d$ denote the eigenvalues of $L(a)$ and consider the condition,

$$(4.54) \quad \text{Sp } \mu(a) \cap \Lambda_a = \emptyset,$$

where the singular set Λ_a is defined as follows,

$$(4.55) \quad \Lambda_a = (-\infty, -\frac{1}{2} \text{Trace } |L(a)|] \cup [\frac{1}{2} \text{Trace } |L(a)|, \infty) \quad \text{if } 2n < d,$$

$$(4.56) \quad \Lambda_a = \{\pm(\frac{1}{2} \text{Trace } |L(a)| + \sum_{j=1}^d \alpha_j |\lambda_j|); \alpha_j \in \mathbb{N}^d\} \quad \text{if } 2n = d.$$

As it turns out this condition makes sense independently of the choice of the H -frame and is the relevant condition to look at in the case of a sublaplacian. More precisely, we have:

Proposition 4.38 ([BG, Thm. 18.4], [Po8]). 1) The condition (4.54) makes sense intrinsically for any $a \in M$.

2) At every point $a \in M$ the Rockland conditions for Δ and Δ^t are equivalent to (4.54).

3) The principal symbol of Δ is invertible if, and only if, the condition (4.54) holds at every point of M , so when the latter occurs Δ admits a parametrix in $\Psi_H^{-2}(M, \mathcal{E})$ and is hypoelliptic with loss of one derivative.

Remark 4.39. In the sequel we will also make use of the weaker condition,

$$(4.57) \quad \text{Sp } \mu(a) \cap \Lambda'_a = \emptyset,$$

$$(4.58) \quad \Lambda'_a = (-\infty, -\frac{1}{2} \text{Trace } |L(a)|] \cup [\frac{1}{2} \text{Trace } |L(a)|, \infty).$$

This condition was considered in [BGS] and implies the existence of a parametrix for the heat operator associated to a sublaplacian in the Volterra-Heisenberg calculus of [BGS] (see Section 6).

4.9. Examples. We now briefly explained how Proposition 4.37 and Proposition 4.38 can be used in the case of the examples described in Section 3.

(a) *Generalized sums of squares.* If a sublaplacian Δ is a generalized sum of squares of the form (3.3) then in (4.52) the matrix $\mu(x)$ is diagonal with purely imaginary values, so that the condition (4.54) is satisfied if, and only if, the Levi form does not vanish at x . Thus Δ has an invertible principal symbol if, and only if, the Levi form of (M, H) is nonvanishing or, equivalently, the bracket condition $H + [H, H] = TM$ is satisfied. In this case Δ admits a Ψ_H DO parametrix and is hypoelliptic, so in the special case of a Heisenberg manifold we recover the hypoellipticity result of [Hö2].

(b) *Kohn Laplacian.* Let $\square_b : C^\infty(M, \Lambda^{p,q}) \rightarrow C^\infty(M, \Lambda^{p,q})$ be the Kohn Laplacian on a CR manifold M^{2n+1} acting on (p, q) -forms, $0 \leq p, q \leq n$. Then at point x the condition (4.54) for $\square_b$ is equivalent to the $Y(q)$ -condition (see [BG], [Po8]). Thus $\square_b$ has an invertible principal symbol if, and only if, the condition $Y(q)$ holds everywhere. In this case $\square_b$ admits a parametrix in $\Psi_H^{-2}(M, \Lambda^{p,q})$ and we recover the theorem of Kohn [Koh1] on the hypoellipticity of $\square_b$.

(c) *Horizontal sublaplacian.* Let $\Delta_b : C^\infty(M, \Lambda_{\mathbb{C}}^k H^*) \rightarrow C^\infty(M, \Lambda_{\mathbb{C}}^k H^*)$ be the horizontal sublaplacian on a Heisenberg manifold (M^{d+1}, H) acting on horizontal forms of degree k , $0 \leq k \leq d$. Then at a point $x \in M$ the condition (4.54) for Δ_b are equivalent to the condition $X(k)$ (see [Po8]). Therefore, Δ_b has an invertible principal symbol if, and only if, the condition $X(k)$ holds at every point. In particular, in case of a non-vanishing Levi form Δ_b has always an invertible principal symbol when acting on functions. In any case, when the condition $X(k)$ holds everywhere Δ_b admits a parametrix in $\Psi_H^{-2}(M, \Lambda_{\mathbb{C}}^k H^*)$ and is hypoelliptic with loss of one derivative.

(d) *Conformal powers of the horizontal sublaplacian.* Let (M^{2n+1}, θ) be a strictly pseudoconvex CR manifold equipped with a pseudohermitian structure and for $k = 1, \dots, n$ let $\square_\theta^{(k)}$ be a conformal power of Δ_b acting on functions as in [JL1] and [GG]. As on functions Δ_b satisfies the Rockland condition at every point and has invertible principal symbol, the same is true for all its integer powers. As $\square_\theta^{(k)}$ and Δ_b^k has same principal symbol, it follows that $\square_\theta^{(k)}$ satisfies the Rockland condition at every point and has an invertible principal symbol. In particular, the operator $\square_\theta^{(k)}$ admits a parametrix in $\Psi_H^{-2k}(M)$ and is hypoelliptic with loss of k derivatives.

(e) *Contact Laplacian.* Let (M^{2n+1}, θ) be an orientable contact manifold. For $k = 0, 1, \dots, 2n$ with $k \neq n$ let $\Delta_R : C^\infty(M, \Lambda^k) \rightarrow C^\infty(M, \Lambda^k)$ be the contact Laplacian on M acting on contact forms of degree k . Then Δ_b satisfies the Rockland condition at every point (see [Ru, p. 100]). Since in the contact case the Levi form of (M, H) has constant rank $2n$, we then may apply Proposition 4.37 to deduce that Δ_R has an invertible principal symbol, hence admits a parametrix in $\Psi_H^{-2}(M, \Lambda^k)$ and is hypoelliptic with loss of one derivative.

Likewise, the contact Laplacian $\Delta_R : C^\infty(M, \Lambda_*^n) \rightarrow C^\infty(M, \Lambda_*^n)$ acting on contact forms of degree n satisfies the Rockland condition at every point and has an invertible principal symbol, so it admits a parametrix in $\Psi_H^{-4}(M, \Lambda_*^n)$ and is hypoelliptic with loss of two derivatives.

5. HOLOMORPHIC FAMILIES OF Ψ DOs

In this section we define holomorphic families of Ψ_H DO's and check their main properties. To this end we make use of an "almost homogeneous" approach to the Heisenberg calculus, described in the first subsection.

5.1. Almost homogeneous approach to the Heisenberg calculus. In this subsection we explain how the Ψ DO's can be described in terms of symbols and kernels which are almost homogeneous, in the sense that there are homogeneous modulo infinite order terms.

Definition 5.1. A symbol $p(x, \xi) \in C^\infty(U \times \mathbb{R}^{d+1})$ is almost homogeneous of degree m , $m \in \mathbb{C}$, when we have

$$(5.1) \quad p(x, \lambda \xi) - \lambda^m p(x, \xi) \in S^{-\infty}(U \times \mathbb{R}^{d+1}) \quad \text{for any } \lambda > 0.$$

We let $S_{ah}^m(U \times \mathbb{R}^{d+1})$ denote the space of almost homogeneous symbols of degree m .

Lemma 5.2 ([BG, Prop. 12.72]). Let $q(x, \xi) \in C^\infty(U \times \mathbb{R}^{d+1})$. Then we have equivalence:

- (i) The symbol $q(x, \xi)$ is almost homogeneous of degree m ;
- (ii) The symbol $q(x, \xi)$ is in $S^m(U \times \mathbb{R}^{d+1})$ and we have $q \sim p_m$ with $p_m \in S_m(U \times \mathbb{R}^{d+1})$ (i.e. the only nonzero homogeneous term in the asymptotic expansion (4.13) for q is p_m).

Granted this we shall now prove:

Lemma 5.3. Let $p \in C^\infty(U \times \mathbb{R}^{d+1})$. Then we have equivalence:

- (i) The function p belongs to $S^m(U \times \mathbb{R}^{d+1})$.
- (ii) For $j = 0, 1, \dots$ there exists $q_{m-j} \in S_{ah}^{m-j}(U \times \mathbb{R}^{d+1})$ such that $p \sim \sum_{j \geq 0} q_{m-j}$.

Proof. Suppose that for $j = 0, 1, \dots$ there exists $q_{m-j} \in S_{ah}^{m-j}(U \times \mathbb{R}^{d+1})$ such that $p \sim \sum_{j \geq 0} q_{m-j}$. By Lemma 5.2 there exists $p_{m-j} \in S_{m-j}(U \times \mathbb{R}^{d+1})$ such that $q_{m-j} \sim p_{m-j}$. Then we have $p \sim \sum_{j \geq 0} p_{m-j}$ and so p belongs to $S^m(U \times \mathbb{R}^{d+1})$. Thus (ii) implies (i).

Conversely, assume that p belongs to $S^m(U \times \mathbb{R}^{d+1})$ and let us write $p \sim \sum_{j \geq 0} p_{m-j}$, $p_l \in S_l(U \times \mathbb{R}^{d+1})$. Let $\varphi \in C_c^\infty(\mathbb{R}^{d+1})$ be such that $\varphi(\xi) = 1$ near $\xi = 1$ and $\varphi(\xi) = 0$ for $\|\xi\| \leq 1$. For $j = 0, 1, \dots$ set $q_{m-j}(x, \xi) = (1 - \varphi(\xi))p_{m-j}(x, \xi)$. As for any $t > 0$ the function $q_{m-j}(x, t\xi) - t^{m-j}q_{m-j}(x, \xi)$ is equal to $(\varphi(t\xi) - \varphi(\xi))p_{m-j}(x, \xi)$, so belongs to $S^{-\infty}(U \times \mathbb{R}^{d+1})$, we see that q_{m-j} lies in $S_{ah}^{m-j}(U \times \mathbb{R}^{d+1})$. Moreover, as we have $q_{m-j}(x, \xi) = p_{m-j}(x, \xi)$ for $\|\xi\| > 1$ we see that $p \sim \sum_{j \geq 0} \tilde{p}_{m-j}$. Hence (i) implies (ii). $\square$

The almost homogeneous symbols have been considered in [BG, Sect. 12] already. In the sequel it will be important to have a "dual" notion of almost homogeneity for distributions as follows.

Definition 5.4. The space $\mathcal{D}'_{reg}(\mathbb{R}^{d+1})$ consists of the distributions on $\mathbb{R}^{d+1}$ that are smooth outside the origin. It is endowed with the weakest topology that makes continuous the inclusions of $\mathcal{D}'_{reg}(\mathbb{R}^{d+1})$ into $\mathcal{D}'(\mathbb{R}^{d+1})$ and $C^\infty(\mathbb{R}^{d+1} \setminus \{0\})$.

Definition 5.5. A distribution $K(x, y) \in C^\infty(U) \hat{\otimes} \mathcal{D}'_{reg}(\mathbb{R}^{d+1})$ is said to be almost homogeneous of degree m , $m \in \mathbb{C}$, when

$$(5.2) \quad K(x, \lambda y) - \lambda^m K(x, y) \in C^\infty(U \times \mathbb{R}^{d+1}) \quad \text{for any } \lambda > 0.$$

We let $\mathcal{K}_{ah}^m(U \times \mathbb{R}^{d+1})$ denote the space of almost homogeneous distributions of degree m .

Proposition 5.6 (Compare [Tay, pp. 18-21]). Let $K(x, y) \in C^\infty(U) \hat{\otimes} \mathcal{D}'_{reg}(\mathbb{R}^{d+1})$ and set $\hat{m} = -(m + d + 2)$. Then we have equivalence:

- (i) The distribution K belongs to $\mathcal{K}_{ah}^m(U \times \mathbb{R}^{d+1})$.
- (ii) We can put $K(x, y)$ into the form,

$$(5.3) \quad K(x, y) = K_m(x, y) + R(x, y),$$

for some $K_m \in \mathcal{K}_m(U \times \mathbb{R}^{d+1})$ and $R \in C^\infty(U \times \mathbb{R}^{d+1})$.

- (iii) We can put $K(x, y)$ into the form,

$$(5.4) \quad K(x, y) = \check{p}_{\xi \rightarrow y}(x, y) + R(x, y),$$

for some $p \in S_{\text{ah}}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ and $R \in C^\infty(U \times \mathbb{R}^{d+1})$.

Proof. First, if $K_m \in \mathcal{K}_m(U \times \mathbb{R}^{d+1})$ then (4.21) implies that, for any $\lambda > 0$, the distribution $K(x, \lambda \cdot y) - \lambda^m K(x, y)$ is in $C^\infty(U \times \mathbb{R}^{d+1})$. Thus (ii) implies (i).

Second, let $p \in S_{\text{ah}}^{\hat{m}}(U \times \mathbb{R}^{d+1})$. By Lemma 5.2 there is $p_{\hat{m}} \in S_m(U \times \mathbb{R}^{d+1})$ such that $p \sim p_{\hat{m}}$. Thanks to Lemma 4.13 we extend $p_{\hat{m}}$ into a distribution $\tau(x, \xi)$ in $C^\infty(U) \hat{\otimes} \mathcal{S}'(\mathbb{R}^{d+1})$ such that $\tilde{\tau}_{\xi \rightarrow y}(x, y)$ is in $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$. Let $\varphi \in C_c^\infty(\mathbb{R}^{d+1})$ be such that $\varphi = 1$ near the origin. Then we can write

$$(5.5) \quad p = \tau + \varphi(p - \tau) + (1 - \varphi)(p - p_{\hat{m}}).$$

Here $\varphi(\xi)(p(x, \xi) - \tau(x, \xi))$ belongs to $C^\infty(U) \otimes \mathcal{D}'(\mathbb{R}^{d+1})$ and is supported on a fixed compact set with respect to ξ , so $[\varphi(p - \tau)]_{\xi \rightarrow y}^\vee$ is smooth. Moreover, as $p \sim p_{\hat{m}}$ both $(1 - \varphi)(p - p_m)$ and $[(1 - \varphi)(p - p_m)]_{\xi \rightarrow y}^\vee$ are in $S^{-\infty}(U \times \mathbb{R}^{d+1})$. It then follows that $\check{p}_{\xi \rightarrow y}$ coincides with $\tilde{\tau}_{\xi \rightarrow y}$ up to an element of $C^\infty(U \times \mathbb{R}^{d+1})$. Since $\tilde{\tau}_{\xi \rightarrow y}(x, y)$ is in $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$ we deduce from this that (iii) implies (ii).

To complete the proof it remains to show that (i) implies (iii). Assume that $K(x, y)$ belongs to $\mathcal{K}_{\text{ah}}^m(U \times \mathbb{R}^{d+1})$. Let $\varphi(y) \in C_c^\infty(\mathbb{R}^{d+1})$ be so that $\varphi(y) = 1$ near $y = 0$ and set $p = (\varphi K)_{y \rightarrow \xi}^\wedge(x, y)$. Then p is a smooth and has slow growth with respect to ξ . Moreover $\check{p}_{\xi \rightarrow y}(x, y)$ differs from $K(x, y)$ by the smooth function $(1 - \varphi(y))K(x, y)$.

Next, using (4.23) we see that, for any $\lambda > 0$, the function $p(x, \lambda \cdot \xi) - \lambda^{\hat{m}} p(x, \xi)$ is equal to

$$(5.6) \quad \lambda^{-(d+2)} [\varphi(\lambda^{-1} \cdot y) K(x, \lambda^{-1} \cdot y) - \lambda^{-m} \varphi(y) K(x, y)]_{y \rightarrow \xi}^\wedge \\ = \lambda^{-(d+2)} [(\varphi(\lambda^{-1} \cdot y) - \varphi(y)) K(x, \lambda^{-1} \cdot y) + \varphi(y) (K(x, \lambda^{-1} \cdot y) - \lambda^{-m} K(x, y))]_{y \rightarrow \xi}^\wedge.$$

Note that $(\varphi(\lambda^{-1} \cdot y) - \varphi(y)) K(x, \lambda^{-1} \cdot y) + \varphi(y) (K(x, \lambda^{-1} \cdot y) - \lambda^{-m} K(x, y))$ belongs $C^\infty(U \times \mathbb{R}^{d+1})$ and is compactly supported with respect to y , so it belongs to $S^{-\infty}(U \times \mathbb{R}^{d+1})$. Since the latter is also true for Fourier transform with respect to y we see that $p(x, \lambda \cdot \xi) - \lambda^{\hat{m}} p(x, \xi)$ is in $S^{-\infty}(U \times \mathbb{R}^{d+1})$ for any $\lambda > 0$, that is, the symbol p is almost homogeneous of degree $\hat{m}$. It then follows that the distribution K satisfies (iii). This proves that (i) implies (iii). The proof is thus achieved. $\square$

5.2. Holomorphic families of Ψ_H DO's. From now on we let Ω denote an open subset of $\mathbb{C}$.

Definition 5.7. A family $(p_z)_{z \in \Omega} \subset S^*(U \times \mathbb{R}^{d+1})$ is said to be holomorphic when:

- (i) The order $m(z)$ of p_z depends analytically on z ;
- (ii) For any $(x, \xi) \in U \times \mathbb{R}^{d+1}$ the function $z \rightarrow p_z(x, \xi)$ is holomorphic on Ω ;
- (iii) The bounds of the asymptotic expansion $p_z \sim \sum_{j \geq 0} p_{z, m(z) - j}$, $p_{z, l} \in S_l(U \times \mathbb{R}^{d+1})$, are locally uniform with respect to z , that is, for any integer N and any compacts $K \subset U$ and $L \subset \Omega$ we have

$$(5.7) \quad |\partial_x^\alpha \partial_\xi^\beta (p_z - \sum_{j < N} p_{z, m(z) - j})(x, \xi)| \leq C_{NKL\alpha\beta} \|\xi\|^{\Re m(z) - N - \langle \beta \rangle}, \quad x \in K, \|\xi\| \geq 1, z \in L.$$

We let $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ denote the set of the families $(p_z)_{z \in \Omega} \subset S^*(U \times \mathbb{R}^{d+1})$ that are holomorphic.

Remark 5.8. If $(p_z)_{z \in \Omega}$ is a holomorphic family of symbols then the homogeneous symbols $p_{z, m(z) - j}$ depend analytically on z . Indeed, for $\xi \neq 0$ we have

$$(5.8) \quad p_{z, m(z)}(x, \xi) = \lim_{\lambda \rightarrow \infty} \lambda^{-m(z)} p_z(x, \lambda \cdot \xi).$$

Since the above axioms imply that the family $(\lambda^{-m(z)} p_z(x, \lambda \cdot \xi))_{\lambda \geq 1}$ is bounded in the Fréchet-Montel space $\text{Hol}(\Omega, C^\infty(U \times (\mathbb{R}^{d+1} \setminus \{0\})))$ the convergence actually holds in $\text{Hol}(\Omega, C^\infty(U \times (\mathbb{R}^{d+1} \setminus \{0\})))$.

Hence $p_{z,m(z)}$ depends analytically on z . Moreover, as for $\xi \neq 0$ we also have

$$(5.9) \quad p_{j,z}(x, \xi) = \lim_{\lambda \rightarrow \infty} \lambda^{j-m(z)} (p_z(x, \lambda \xi) - \sum_{l < j} \lambda^{m(z)-l} p_{z,m(z)-l}(x, \xi)), \quad j = 1, 2, \dots,$$

an easy induction shows that all the symbols p_{z,m_z-j} depend analytically on z .

Recall that $\Psi^{-\infty}(U) = \mathcal{L}(\mathcal{E}'(U), C^\infty(U))$ is naturally a Fréchet space which is isomorphic to $C^\infty(U \times U)$ by Schwartz's kernel theorem. Therefore holomorphic families of smoothing operators makes well sense and we may define holomorphic families of $\Psi_H DO$'s as follows.

Definition 5.9. *A family $(P_z)_{z \in \Omega} \subset \Psi_H^m(U)$ is holomorphic when it can be put into the form*

$$(5.10) \quad P_z = p_z(x, -iX) + R_z \quad z \in \Omega,$$

for some family $(p_z)_{z \in \Omega} \in \text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ and some family $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi^{-\infty}(U))$. We let $\text{Hol}(\Omega, \Psi_H^*(U))$ denote the set of holomorphic families of $\Psi_H DO$'s.

For technical sake it will be useful to consider the symbol class below.

Definition 5.10. $S_{||}^k(U \times \mathbb{R}^{d+1})$, $k \in \mathbb{R}$, consists of symbols $p(x, \xi) \in C^\infty(U \times \mathbb{R}^{d+1})$ such that, for any compact $K \subset U$, we have

$$(5.11) \quad |\partial_x^\alpha \partial_\xi^\beta p(x, \xi)| \leq C_{K\alpha\beta} (1 + \|\xi\|)^{k-\langle \beta \rangle}, \quad (x, \xi) \in K \times \mathbb{R}^{d+1}.$$

Its topology is given by the sharpest constants $C_{K\alpha\beta}$'s in the estimates (5.11).

Note that the estimates (4.13) imply that $S^m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, is contained in $S_{||}^k(U \times \mathbb{R}^{d+1})$ for any $k \geq \Re m$.

Proposition 5.11. *Let $(P_z)_{z \in \Omega}$ be a holomorphic family of $\Psi_H DO$'s. Then:*

- 1) $(P_z)_{z \in \Omega}$ gives rise to families in $\text{Hol}(\Omega, \mathcal{L}(C_c^\infty(U), C^\infty(U)))$ and $\text{Hol}(\Omega, \mathcal{L}(\mathcal{E}'(U)), \mathcal{D}'(U))$.
- 2) Off the diagonal of $U \times U$ the distribution kernel of P_z is represented by a holomorphic family of smooth functions.

Proof. Without any loss of generality we may suppose that $P_z = p_z(x, -iX)$, with $(p_z)_{z \in \Omega}$ in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$. Moreover, shrinking Ω if necessary, we may also assume that the degree m_z of p_z stays bounded, as much so $(p_z)_{z \in \Omega}$ is contained in $S_{||}^k(U \times \mathbb{R}^{d+1})$ for some real $k \geq 0$. Let $\sigma_j(x, \xi)$ denote the classical symbol of $-iX_j$ and set $\sigma = (\sigma_0, \dots, \sigma_d)$. Then the proof of [BG, Prop. 10.22] shows that the map $p(x, \xi) \rightarrow p^\sigma(x, \xi) := p(x, \sigma(x, \xi))$ is continuous from $S_{||}^k(U \times \mathbb{R}^{d+1})$ to $S_{\frac{1}{2}, \frac{1}{2}}^k(U \times \mathbb{R}^{d+1})$. Thus, the family $(p_z^\sigma)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S_{\frac{1}{2}, \frac{1}{2}}^k(U \times \mathbb{R}^{d+1}))$.

Next, it follows from the proof of [Hö1, Thm. 2.2] that:

- (i) The quantization map $q \rightarrow q(x, D)$ induces continuous $\mathbb{C}$ -linear maps from $S_{\frac{1}{2}, \frac{1}{2}}^k(U \times \mathbb{R}^{d+1})$ to $\mathcal{L}(C_c^\infty(U), C^\infty(U))$ and to $\mathcal{L}(\mathcal{E}'(U), \mathcal{D}'(U))$;
- (ii) The linear map $q(x, \xi) \rightarrow \check{q}_{\xi \rightarrow y}(x, y)$ is continuous from $S_{\frac{1}{2}, \frac{1}{2}}^k(U \times \mathbb{R}^{d+1})$ to $C^\infty(U) \otimes \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1})$, in such way that for any $q \in S_{\frac{1}{2}, \frac{1}{2}}^k(U \times \mathbb{R}^{d+1})$ the distribution kernel $\check{q}_{\xi \rightarrow y}(x, x - y)$ of $q(x, D)$ is represented off the diagonal by a smooth function depending continuously on q .

As a continuous $\mathbb{C}$ -linear map is analytic it follows that, on the one hand, $(p_z(x, -iX))_{z \in \Omega}$ gives rise to elements of $\text{Hol}(\Omega, \mathcal{L}(C_c^\infty(U), C^\infty(U)))$ and $\text{Hol}(\Omega, \mathcal{L}(\mathcal{E}'(U), \mathcal{D}'(U)))$ and, on the other hand, the distribution kernel of P_z is represented outside the diagonal of $U \times U$ by a holomorphic family of smooth functions. $\square$

Definition 5.12. Let $(P_z)_{z \in \Omega} \subset \mathcal{L}(C_c^\infty(U), C^\infty(U))$ and for $z \in \Omega$ let $k_{P_z}(x, y)$ denote the distribution kernel of P_z . Then the family $(P_z)_{z \in \Omega}$ is said to be uniformly properly supported when, for any compact $K \subset U$ there exist compacts $L_1 \subset U$ and $L_2 \subset K$ such that for any $z \in \Omega$ we have

$$(5.12) \quad \text{supp } k_{P_z}(x, y) \cap (U \times K) \subset L_1 \quad \text{and} \quad \text{supp } k_{P_z}(x, y) \cap (K \times U) \subset L_2.$$

Proposition 5.13. Let $(P_z)_{z \in \Omega}$ be a holomorphic family of Ψ_H DO's.

1) We can write P_z in the form $P_z = Q_z + R_z$ with $(Q_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi_H^*(U))$ uniformly properly supported and $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi^{-\infty}(U))$.

2) If the family $(P_z)_{z \in \Omega}$ is uniformly properly supported then it gives rise to holomorphic families of continuous endomorphisms of $C_c^\infty(U)$ and $C^\infty(U)$ and of $\mathcal{E}'(U)$ and $\mathcal{D}'(U)$.

Proof. Let $(\varphi_i)_{i \geq 0} \subset C_c^\infty(U)$ be a partition of unity which is subordinated to a locally finite covering $(U_i)_{i \geq 0}$ of U by relatively compact open subsets. For each $i \geq 0$ let $\psi_i \in C_c^\infty(U)$ be such that $\psi_i = 1$ near $\text{supp } \varphi_i$ and set $\chi(x, y) = \sum \varphi_i(x) \psi_i(y)$. Then χ is a smooth function on $U \times U$ which is properly supported and such that $\chi(x, y) = 1$ near the diagonal of $U \times U$.

For $z \in \Omega$ let $k_{P_z}(x, y)$ denote the distribution kernel of P_z and let Q_z and R_z be the elements of $\mathcal{L}(C_c^\infty(U), C^\infty(U))$ with respective distribution kernels

$$(5.13) \quad k_{Q_z}(x, y) = \chi(x, y) k_{P_z}(x, y) \quad \text{and} \quad k_{R_z}(x, y) = (1 - \chi(x, y)) k_{P_z}(x, y).$$

Notice that since χ is properly supported the family $(Q_z)_{z \in \Omega}$ is uniformly properly supported. As by Proposition 5.11 the distribution $k_{P_z}(x, y)$ is represented outside the diagonal of $U \times U$ by a holomorphic family of smooth functions, we see that $(k_{R_z}(x, y))$ is a holomorphic family of smooth kernels, i.e. $(R_z)_{z \in \Omega}$ is a holomorphic family of smoothing operators. Since $Q_z = P_z - R_z$ it follows that $(Q_z)_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's. Hence the first assertion.

Assume now that $(P_z)_{z \in \Omega}$ is uniformly properly supported. Thanks to Proposition 5.11 we already know that $(P_z)_{z \in \Omega}$ gives rise to holomorphic families with values in $\mathcal{L}(C_c^\infty(U), C^\infty(U))$ and $\mathcal{L}(\mathcal{E}'(U), \mathcal{D}'(U))$. Let K be a compact subset of U . Then (5.12) implies that there exists a compact $L \subset U$ such that for every $z \in \Omega$ the operator P_z maps $C_K^\infty(U)$ to $C_L^\infty(U)$ and $\mathcal{E}'_K(U)$ to $\mathcal{E}'_L(U)$, in such way that $(P_z)_{z \in \Omega}$ gives rise to holomorphic families with values in $\mathcal{L}(C_K^\infty(U), C_L^\infty(U))$ and $\mathcal{L}(\mathcal{E}'_K(U), \mathcal{E}'_L(U))$. In view of the definitions of the topologies of $C_c^\infty(U)$ and $\mathcal{E}'(U)$ as the inductive limit topologies of $C_K^\infty(U)$ and $\mathcal{E}_K(U)$ as K ranges over compacts of U , this shows that the family $(P_z)_{z \in \Omega}$ gives rise to elements of $\text{Hol}(\Omega, C_c^\infty(U))$ and $\text{Hol}(\Omega, \mathcal{E}'(U))$.

Next, let $(\varphi_i)_{i \geq 0} \subset C_c^\infty(U)$ be a partition of unity. For each index i let K_i be a compact neighborhood of $\text{supp } \varphi_i$. Then (5.12) implies that there exists a compact $L_i \subset U$ such that $\text{supp } k_{P_z}(x, y) \cap (K_i \times U) \subset L_i$ for every $z \in \Omega$. Let $\psi_i \in C^\infty(U)$ be such that $\psi_i = 1$ near K_i . Then we have

$$(5.14) \quad P_z = \sum_{i \geq 0} \varphi_i P_z = \sum_{i \geq 0} \varphi_i P_z \psi_i.$$

Since each family $(\varphi_i P_z \psi_i)_{z \in \Omega}$ is holomorphic with values in $\mathcal{L}(C^\infty(U))$ and $\mathcal{L}(\mathcal{D}'(U))$ and the sums are locally finite this shows that $(P_z)_{z \in \Omega}$ gives rise to elements of $\text{Hol}(\Omega, C^\infty(U))$ and $\text{Hol}(\Omega, \mathcal{D}'(U))$. $\square$

5.3. Composition of holomorphic families of Ψ_H DO's. Let us now look at the analyticity of the composition of Ψ_H DO's. To this end we need to deal with holomorphic families of almost homogeneous symbols as follows.

Definition 5.14. A family $(q_z)_{z \in \Omega} \subset S_{ah}^*(U \times \mathbb{R}^{d+1})$ is holomorphic when:

- (i) The degree $m(z)$ of q_z is a holomorphic function on Ω ;
- (ii) The family $(q_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$;
- (iii) For any $t > 0$ the family $(q_z(x, t, \xi) - t^{m(z)} q_z(x, \xi))_{z \in \Omega}$ is in $\text{Hol}(\Omega, S^{-\infty}(U \times \mathbb{R}^{d+1}))$.

We let $\text{Hol}(\Omega, S_{ah}^*(U \times \mathbb{R}^{d+1}))$ denote the set of holomorphic families of almost homogeneous symbols.

Lemma 5.15. *Let $(q_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$. Then we have equivalence:*

- (i) *The family $(q_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S_{ah}^*(U \times \mathbb{R}^{d+1}))$ and has degree $m(z)$;*
- (ii) *The family $(q_z)_{z \in \Omega}$ lies in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ and, in the sense of (5.7), we have $q_z \sim p_z$ where, for every $z \in \Omega$, the symbol p_z belongs to $S_{m(z)}(U \times \mathbb{R}^{d+1})$.*

Proof. Assume that $(q_z)_{z \in \Omega}$ is in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ and, in the sense of (5.7), we have $q_z \sim p_z$ where, for every $z \in \Omega$, the symbol p_z belongs to $S_{m(z)}(U \times \mathbb{R}^{d+1})$. Then the order $m(z)$ of q_z is a holomorphic function on Ω and, for any compact subset $K \subset U$, any integer N and any open $\Omega' \subset\subset \Omega$, we have

$$(5.15) \quad |\partial_x^\alpha \partial_\xi^\beta (q_z - p_z)(x, \xi)| \leq C_{NK\Omega' \alpha \beta} \|\xi\|^{-N},$$

for $x \in K$, $\|\xi\| \geq 1$ and $z \in \Omega'$. It follows that, for any $t > 0$, the family $\{q_z(x, t\xi) - t^{m(z)} q_z(x, \xi)\}_{z \in \Omega} = \{(q_z(x, t\xi) - q_z(x, \xi)) - t^{m(z)}(q_z(x, \xi) - q_z(x, \xi))\}_{z \in \Omega}$ is contained in $\text{Hol}(\Omega, S^{-\infty}(U \times \mathbb{R}^{d+1}))$. Thus $(q_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S_{ah}^*(U \times \mathbb{R}^{d+1}))$.

Conversely, suppose that $(q_z)_{z \in \Omega}$ is contained in $\text{Hol}(\Omega, S_{ah}^*(U \times \mathbb{R}^{d+1}))$ and has degree $m(z)$. Then, for any $t > 0$, any compact $K \subset U$, any integer N and any open $\Omega' \subset\subset \Omega$, we have

$$(5.16) \quad |\partial_x^\alpha \partial_\xi^\beta (q_z(x, t\xi) - t^{m(z)} q_z(x, \xi))| \leq C_{tNK\Omega' \alpha \beta} (1 + \|\xi\|)^{-N},$$

for $(x, \xi) \in K \times \mathbb{R}^{d+1}$ and $z \in \Omega'$. Then replacing ξ by $s\xi$, $s > 0$, in (5.16) shows that when $N \geq \sup_{z \in \Omega'} \Re m(z)$ we have

$$(5.17) \quad |\partial_x^\alpha \partial_\xi^\beta (s^{m(z)} q_z(x, st\xi) - (st)^{m(z)} q_z(x, s\xi))| \leq C_{tNK\Omega' \alpha \beta} s^{\Re m(z) - N} \|\xi\|^{-N} \\ \leq C_{tNK\Omega' \alpha \beta} s^{-1} \|\xi\|^{-N}.$$

for $(x, \xi) \in K \times \mathbb{R}^{d+1} \setminus 0$ and $z \in \Omega'$.

Next, for $k \in \mathbb{N}$ let $q_{z,k}(x, \xi) = (2^k)^{-m} q_z(x, 2^k \xi)$. Then, for any compact $K \subset U$, any open $\Omega' \subset\subset \Omega$ and any integer $N \geq \sup_{z \in \Omega'} \Re m(z)$, we have

$$(5.18) \quad |\partial_x^\alpha \partial_\xi^\beta (q_{z,k+1}(x, \xi) - q_{z,k}(x, \xi))| \leq C_{2NK\Omega' \alpha \beta} 2^{-k} \|\xi\|^{-N},$$

for $(x, \xi) \in K \times \mathbb{R}^{d+1} \setminus 0$ and $z \in \Omega'$. This shows that the series $\sum_{k \geq 0} (q_{z,k+1} - q_{z,k})$ is convergent in $\text{Hol}(\Omega, C^\infty(U \times (\mathbb{R}^{d+1} \setminus 0)))$. Hence the sequence $(q_{z,k})_{k \geq 0}$ converges in $\text{Hol}(\Omega, C^\infty(U \times (\mathbb{R}^{d+1} \setminus 0)))$ to some family $(p_z)_{z \in \Omega}$. In fact, taking $s = 2^k$ in (5.17) and letting $k \rightarrow \infty$ with t fixed shows that q_z is homogeneous of degree $m(z)$ with respect to the ξ -variable. Moreover, for any compact $K \subset U$, any open $\Omega' \subset\subset \Omega$ and any integer $N \geq \sup_{z \in \Omega'} \Re m(z)$, we have

$$(5.19) \quad |\partial_x^\alpha \partial_\xi^\beta (q_z - p_z)(x, \xi)| \leq \sum |\partial_x^\alpha \partial_\xi^\beta (q_{z,k+1} - q_{z,k})(x, \xi)| \leq C_{2NK\Omega' \alpha \beta} \|\xi\|^{-N},$$

for $(x, \xi) \in K \times (\mathbb{R}^{d+1} \setminus 0)$ and $z \in \Omega'$, i.e. we have $q_z \sim p_z$ in the sense of (5.7). $\square$

Using Lemma 5.15 and arguing as in the proof of Lemma 5.3 we get the following characterization of holomorphic families of symbols.

Lemma 5.16. *Let $(p_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$. Then we have equivalence:*

- (i) *The family $(p_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ and has holomorphic order $m(z)$.*
- (ii) *For $j = 0, 1, \dots$ there exists $(q_{j,z})_{z \in \Omega} \in \text{Hol}(\Omega, S_{ah}(U \times \mathbb{R}^{d+1}))$ almost homogeneous of degree $m(z) - j$ so that we have $p_z \sim \sum_{j \geq 0} q_{j,z}$ in the sense of (5.7).*

Next, it is shown in [BG, Sect. 12] that, as for homogeneous symbols in (4.16)–(4.19), there is a continuous bilinear product,

$$(5.20) \quad * : S_{||}^{k_1}(U \times \mathbb{R}^{d+1}) \times S_{||}^{k_2}(U \times \mathbb{R}^{d+1}) \longrightarrow S_{||}^{k_1+k_2}(U \times \mathbb{R}^{d+1}),$$

which is homogeneous in the sense that, for any $\lambda \in \mathbb{R}$, we have

$$(5.21) \quad (p_1 * p_2)_\lambda = p_{1,\lambda} * p_{2,\lambda}, \quad p_j \in S_{||}^{k_j}(U \times \mathbb{R}^{d+1}).$$

This product is related to the product of homogeneous symbols as follows.

Lemma 5.17 ([BG, Sect. 13]). *For $j = 1, 2$ let $p_j \in S^{m_j}(U \times \mathbb{R}^{d+1})$ have principal symbol $p_{m_j} \in S_{m_j}(U \times \mathbb{R}^{d+1})$. Then $p_1 * p_2$ lies in $S^{m_1+m_2}(U \times \mathbb{R}^{d+1})$ and has principal symbol $p_{m_1} * p_{m_2}$.*

Furthermore, this product is holomorphic, for we have:

Lemma 5.18. *For $j = 1, 2$ let $(p_{j,z})_{z \in \Omega} \subset S^*(U \times \mathbb{R}^{d+1})$ be a holomorphic family of symbols. Then $(p_{1,z} * p_{2,z})_{z \in \Omega}$ is a holomorphic family of symbols as well.*

Proof. For $j = 1, 2$ let $m_j(z)$ be the order of $p_{j,z}$. Since $m_1(z)$ and $m_2(z)$ are holomorphic functions, possibly by shrinking Ω , we may assume that $\sup_{z \in \Omega} m_j(z) \leq k < \infty$. Then each family $(p_{j,z})_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S_{||}^k(U \times \mathbb{R}^{d+1}))$. Since $*$ is a continuous $\mathbb{C}$ -bilinear map from $S_{||}^k(U \times \mathbb{R}^{d+1}) \times S_{||}^k(U \times \mathbb{R}^{d+1})$ to $S_{||}^{2k}(U \times \mathbb{R}^{d+1})$ we see that $p_{1,z} * p_{2,z}$ is in $\text{Hol}(\Omega, S_{||}^{2k}(U \times \mathbb{R}^{d+1}))$, hence in $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$.

Now, assume that $p_{j,z}$, $j = 1, 2$, is almost homogeneous of degree $m_j(z)$. Then using (5.21) we see that, for any $\lambda > 0$, the symbol $(p_{1,z} * p_{2,z})_\lambda - \lambda^{m_1(z)+m_2(z)} p_{1,z} * p_{2,z}$ is equal to

$$(5.22) \quad p_{1,z,\lambda} * p_{1,z,\lambda} - \lambda^{m_1(z)+m_2(z)} p_{1,z} * p_{2,z} \\ = (p_{1,z,\lambda} - \lambda^{m_1(z)} p_{1,z}) * p_{2,z} + \lambda^{m_1(z)} p_{1,z,\lambda} * (p_{2,z,\lambda} - \lambda^{m_2(z)} p_{2,z}).$$

Since $(p_{1,z,\lambda} - \lambda^{m_1(z)} p_{1,z})_{z \in \Omega}$ and $(p_{2,z,\lambda} - \lambda^{m_2(z)} p_{2,z})_{z \in \Omega}$ both belong to $\text{Hol}(\Omega, S^{-\infty}(U \times \mathbb{R}^{d+1}))$ combining this with the analyticity of $*$ on $S_{||}^*(U \times \mathbb{R}^{d+1})$ shows that, for any $\lambda > 0$, the family $(p_{1,z} * p_{2,z})_\lambda - \lambda^{m_1(z)+m_2(z)} p_{1,z} * p_{2,z}$ belongs to $\text{Hol}(\Omega, S^{-\infty}(U \times \mathbb{R}^{d+1}))$. Then Lemma 5.16 implies that $p_{1,z} * p_{2,z}$ is a holomorphic family of almost homogeneous of symbols of degree $m_1(z) + m_2(z)$.

In general, by Lemma 5.16 we have $p_{j,z} \sim \sum_{l \geq 0} p_{j,z,l}$, with $(p_{j,z,l})_{z \in \Omega} \in \text{Hol}(\Omega, S_{\text{ah}}^*(U \times \mathbb{R}^{d+1}))$ of degree $m_j(z) - l$ and $\sim$ taken in the sense of (5.7). In particular, for any integer N we have $p_{j,z} = \sum_{l < N} p_{j,z,l} \text{ mod } \text{Hol}(\Omega, S_{||}^{k-N}(U \times \mathbb{R}^{d+1}))$. Thus,

$$(5.23) \quad p_{1,z} * p_{2,z} = \sum_{l+p < N} p_{1,z,l} * p_{2,z,p} \quad \text{mod } \text{Hol}(\Omega, S_{||}^{2k-N}(U \times \mathbb{R}^{d+1})).$$

As explained above $(p_{1,z,l} * p_{2,z,p})_{z \in \Omega}$ is a holomorphic family of almost homogeneous of symbols of degree $m_1(z) + m_2(z) - l - p$. It then follows from Lemma 5.16 that $(p_{1,z} * p_{2,z})_{z \in \Omega}$ is a holomorphic family of symbols. $\square$

We are now ready to prove:

Proposition 5.19. *For $j = 1, 2$ let $(P_{j,z})_{z \in \Omega}$ be in $\text{Hol}(\Omega, \Psi_H^*(U))$ and suppose that at least one the families $(P_{1,z})_{z \in \Omega}$ or $(P_{2,z})_{z \in \Omega}$ is uniformly properly supported. Then the family $(P_{1,z} P_{2,z})_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's.*

Proof. By assumption $(P_{1,z})_{z \in \Omega}$ or $(P_{2,z})_{z \in \Omega}$ is a uniformly properly supported holomorphic family of Ψ_H DO's, hence gives rise to elements of $\text{Hol}(\Omega, \mathcal{L}(C^\infty(U)))$ and $\text{Hol}(\Omega, \mathcal{L}(\mathcal{E}'(U)))$ by Proposition 5.13. Moreover, Proposition 5.13 tells us that the other family at least coincides with a uniformly properly supported holomorphic family of Ψ_H DO's up to a holomorphic family of smoothing

operators. It thus follows that $(P_{1,z}P_{2,z})_{z \in \Omega}$ is the product of two uniformly properly supported holomorphic families of Ψ_H DO's up to a holomorphic families of smoothing operators.

As a consequence we may assume that the families $(P_{1,z})_{z \in \Omega}$ and $(P_{2,z})_{z \in \Omega}$ are both uniformly properly supported. Thanks to (5.14) this allows us to write

$$(5.24) \quad P_{1,z}P_{2,z} = \sum_{i \geq 0} \varphi_i P_{1,z} \psi_i P_{2,z},$$

where $(\varphi_i)_{i \geq 0} \subset C_c^\infty(U)$ and $(\psi_i)_{i \geq 0} \subset C_c^\infty(U)$ are locally finite families such that (φ_i) is a partition of the unity and $\psi_i = 1$ near $\text{supp } \varphi_i$.

Next, for $j = 1, 2$ let us write $P_{j,z} = p_{j,z}(x, -iX) + R_{j,z}$, with $(p_{j,z})_{z \in \Omega}$ in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ and $(R_{j,z})_{z \in \Omega}$ in $\text{Hol}(\Omega, \Psi^{-\infty}(U))$. Since by Proposition 5.11 each family $(p_{j,z}(x, -iX))_{z \in \Omega}$ is holomorphic with values in $\mathcal{L}(C_c^\infty(U), C^\infty(U))$ and $\mathcal{L}(\mathcal{E}'(U), \mathcal{D}'(U))$ using (5.24) we see that

$$(5.25) \quad P_{1,z}P_{2,z} = \sum \varphi_i p_{1,z}(x, -iX) \psi_i p_{2,z}(x, -iX) \mod \text{Hol}(\Omega, \Psi^{-\infty}(U)).$$

At this stage we make appeal to:

Lemma 5.20 ([BG, Prop. 14.45]). *For $j = 1, 2$ let $p_j \in S_{||}^{k_j}(U \times \mathbb{R}^{d+1})$ and let $\psi \in C_c^\infty(U)$. Then:*

$$(5.26) \quad p_1(x, -iX) \psi p_2(x, -iX) = p_1 \#_\psi p_2(x, -iX),$$

where $\#_\psi$ is a continuous bilinear map from $S_{||}^{k_1}(U \times \mathbb{R}^{d+1}) \times S_{||}^{k_2}(U \times \mathbb{R}^{d+1})$ to $S_{||}^{k_1+k_2}(U \times \mathbb{R}^{d+1})$. Moreover, for any integer $N \geq 1$ we have

$$(5.27) \quad p_1 \#_\psi p_2 = \sum_{j < N} \sum_{\alpha \beta \gamma \delta}^{(j)} h_{\alpha \beta \gamma \delta} \psi (D_\xi^\delta p_1) * (\xi^\gamma \partial_x^\alpha \partial_\xi^\beta p_2) + R_{N,\psi}(p_1, p_2),$$

where the notations are the same as in Proposition 4.9 and $R_{N,\psi}$ is a continuous bilinear map from $S_{||}^{k_1}(U \times \mathbb{R}^{d+1}) \times S_{||}^{k_2}(U \times \mathbb{R}^{d+1})$ to $S_{||}^{k_1+k_2-N}(U \times \mathbb{R}^{d+1})$.

Remark 5.21. The continuity contents of Lemma 5.20 is explicitly stated in Proposition 14.45 of [BG], but they follow from its proof or from a standard use of the closed graph theorem.

Now, thanks to Lemma 5.20 we have

$$(5.28) \quad P_{1,z}P_{2,z} = p_z(x, -iX) + R_z, \quad p_z = \sum \varphi_i p_{1,z} \#_{\psi_i} p_{2,z}.$$

Furthermore, possibly by shrinking Ω , we may assume that there is a real k such that for $j = 1, 2$ we have $\Re \text{ord } p_{j,z} \leq k$ for any $z \in \Omega$. Then the continuity contents of Lemma 5.20 imply that $(p_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S_{||}^{2k}(U \times \mathbb{R}^{d+1}))$ and for any integer $N \geq 1$ we can write

$$(5.29) \quad p_z = \sum_{r < N} q_{r,z} + R_{N,z}, \quad q_{j,z} = \sum_{\alpha \beta \gamma \delta}^{(r-s-t)} h_{\alpha \beta \gamma \delta} (D_\xi^\delta p_{1,z}) * (\xi^\gamma \partial_x^\alpha \partial_\xi^\beta p_{2,z}),$$

where $(R_{N,z})_{z \in \Omega} := (\sum_{i \geq 0} \varphi_i R_{N,\psi_i}(p_{1,z}, p_{2,z}))_{z \in \Omega}$ is in $\text{Hol}(\Omega, S_{||}^{2k-N}(U \times \mathbb{R}^{d+1}))$. Thanks to Lemma 5.18 the family $(q_{r,z})_{z \in \Omega}$ is in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ and has order $m_1(z) + m_2(z) - r$. Therefore, we have $p_z \sim \sum_{j \geq 0} q_{j,z}$ in the sense of (5.7), which by Lemma 5.18 implies that $(p_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$. Combining this with (5.28) then shows that $(P_{1,z}P_{2,z})_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's. $\square$

5.4. Kernel characterization of holomorphic families of Ψ DO's. We shall now give a characterization of holomorphic families of Ψ_H DO's in terms of holomorphic families with values in $\mathcal{K}^*(U \times \mathbb{R}^{d+1})$. Since the latter is defined in terms of asymptotic expansions of kernels in $\mathcal{K}_*(U \times \mathbb{R}^{d+1})$ (cf. Definition 4.14) a technical difficulty occurs, because for a family $(K_z)_{z \in \Omega} \subset \mathcal{K}_*(U \times \mathbb{R}^{d+1})$ logarithmic singularities may appear as the order of K_z crosses non-negative integer values.

This issue is resolved by making use of holomorphic families of almost homogeneous distributions as follows.

Definition 5.22. A family $(K_z)_{z \in \Omega} \subset \mathcal{K}_{ah}^*(U \times \mathbb{R}^{d+1})$ is holomorphic when:

- (i) The degree $m(z)$ of K_z is a holomorphic function on Ω ;
- (ii) The family $(K_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(U) \otimes \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1}))$;
- (iii) For any $\lambda > 0$ the family $\{K_z(x, \lambda \cdot y) - \lambda^{m(z)} K_z(x, y)\}_{z \in \Omega}$ is in $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$.

We let $\text{Hol}(\Omega, \mathcal{K}_{ah}^*(U \times \mathbb{R}^{d+1}))$ denote the set of holomorphic families of almost homogeneous distributions.

Lemma 5.23. Let $(K_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U) \otimes \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1}))$. Then we have equivalence:

- (i) The family $(K_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, \mathcal{K}_{ah}^*(U \times \mathbb{R}^{d+1}))$ and has degree $m(z)$.
- (ii) We can put $(K_z)_{z \in \Omega}$ into the form,

$$(5.30) \quad K_z(x, y) = \check{p}_{z, \xi \rightarrow y}(x, y) + R_z(x, y), \quad z \in \Omega,$$

for some family $(p_z)_{z \in \Omega} \in \text{Hol}(\Omega, S_{ah}^*(U \times \mathbb{R}^{d+1}))$ of degree $\hat{m}(z) := -(m(z) + d + 2)$ and some family $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$.

Proof. Assume that the family $(K_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, \mathcal{K}_{ah}^*(U \times \mathbb{R}^{d+1}))$ and has degree $m(z)$. Let $\varphi \in C_c^\infty(U \times \mathbb{R}^{d+1})$ be such that $\varphi(y) = 1$ near $y = 0$ and set $p_z = (\varphi(y) K_z(x, y))_{y \rightarrow \xi}$. Thus,

$$(5.31) \quad K_z(x, y) = \check{p}_{z, \xi \rightarrow y}(x, y) + (1 - \varphi(y)) K_z(x, y) = \check{p}_{z, \xi \rightarrow y}(x, y) \mod \text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1})).$$

As $(\varphi(y) K_z(x, y))_{z \in \Omega}$ is in $\text{Hol}(\Omega, C^\infty(U) \otimes \mathcal{E}'_L(\mathbb{R}^{d+1}))$ with $L = \text{supp } \varphi$, we see that $(p_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$. Furthermore, thanks to (4.23) for any $\lambda > 0$ we have:

$$(5.32) \quad p_z(x, \lambda \cdot \xi) - \lambda^{\hat{m}(z)} p_z(x, \xi) = \lambda^{-(d+2)} [(\varphi(\lambda^{-1} \cdot y) - \varphi(y)) K_z(x, \lambda^{-1} \cdot y) + \varphi(y) (K_z(x, \lambda^{-1} \cdot y) - \lambda^{-m} K_z(x, y))]_{y \rightarrow \xi}^\wedge(x, \xi).$$

Since the r.h.s. above is the Fourier transform with respect to y of an element of $\text{Hol}(\Omega, C_c^\infty(U \times \mathbb{R}^{d+1})) \subset \text{Hol}(\Omega, S^{-\infty}(U \times \mathbb{R}^{d+1}))$ we see that $(p_z(x, \lambda \cdot \xi) - \lambda^{\hat{m}(z)} p_z(x, \xi))_{z \in \Omega}$ is in $\text{Hol}(\Omega, S^{-\infty}(U \times \mathbb{R}^{d+1}))$ for any $\lambda > 0$. Combining this with Lemma 5.15 then shows that $(p_z)_{z \in \Omega}$ is a family of almost homogeneous symbols of degree $\hat{m}(z)$. Conversely, let $(p_z)_{z \in \Omega} \in \text{Hol}(\Omega, S_{ah}^*(U \times \mathbb{R}^{d+1}))$ be almost homogeneous of degree $\hat{m}(z)$. Possibly by shrinking Ω we may assume that we have $\sup_{z \in \Omega} \Re \hat{m}(z) \leq k < \infty$. Then the family $(p_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S_{||}^k(U \times \mathbb{R}^{d+1}))$, hence to $\text{Hol}(\Omega, S_{\frac{1}{2}\frac{1}{2}}^k(U \times \mathbb{R}^{d+1}))$.

As mentioned in the course of the proof of Proposition 5.11, the map $q(x, \xi) \rightarrow \check{q}_{\xi \rightarrow y}(x, y)$ is analytic from $S_{\frac{1}{2}\frac{1}{2}}^k(U \times \mathbb{R}^{d+1})$ to $C^\infty(U) \otimes \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1})$, so the family $(\check{p}_{z, \xi \rightarrow y})_{z \in \Omega}$ lies in $\text{Hol}(\Omega, C^\infty(U) \otimes \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1}))$.

Next, using (4.23) we see that, for any $\lambda > 0$, we have

$$(5.33) \quad \check{p}_{z, \xi \rightarrow y}(x, \lambda \cdot y) - \lambda^{m(z)} \check{p}_{z, \xi \rightarrow y}(x, y) = [p_z(x, \lambda^{-1} \cdot \xi) - \lambda^{-\hat{m}(z)} p_z(x, \xi)]_{\xi \rightarrow y}^\wedge(x, y).$$

Since by Lemma 5.15 the r.h.s. of (5.33) is the inverse Fourier transform with respect to ξ of an element of $\text{Hol}(\Omega, S^{-\infty}(U \times \mathbb{R}^{d+1}))$ we see that the family $\{\check{p}_{z, \xi \rightarrow y}(x, \lambda \cdot y) - \lambda^{m(z)} \check{p}_{z, \xi \rightarrow y}(x, y)\}_{z \in \Omega}$ is contained in $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$. It then follows that $(\check{p}_{z, \xi \rightarrow y})_{z \in \Omega}$ is a holomorphic family of almost homogeneous distributions of degree $m(z)$. $\square$

Definition 5.24. A family $(K_z)_{z \in \Omega} \subset \mathcal{K}^*(U \times \mathbb{R}^{d+1})$ is holomorphic when:

- (i) The order m_z of K_z is a holomorphic function of z ;
- (ii) For $j = 0, 1, \dots$ there exists $(K_{j,z}) \in \text{Hol}(\Omega, \mathcal{K}_{\text{ah}}^*(U \times \mathbb{R}^{d+1}))$ of degree $m(z) + j$ such that $K_z \sim \sum_{j \geq 0} K_{j,z}$ in the sense that, for any open $\Omega' \subset \subset \Omega$ and any integer N , as soon as J is large enough we have

$$(5.34) \quad K_z - \sum_{j \leq J} K_{j,z} \in \text{Hol}(\Omega', C^N(U \times \mathbb{R}^{d+1})).$$

Proposition 5.25. For a family $(K_z)_{z \in \Omega} \subset \mathcal{K}^*(U \times \mathbb{R}^{d+1})$ the following are equivalent:

- (i) The family $(K_z)_{z \in \Omega}$ is holomorphic and has order $m(z)$.
- (ii) We can put $(K_z)_{z \in \Omega}$ into the form,

$$(5.35) \quad K_z(x, y) = (p_z)_{\xi \rightarrow y}^\vee(x, y) + R_z(x, y),$$

for some family $(p_z)_{z \in \Omega} \in \text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ of order $\hat{m}(z) := -(m(z) + d + 2)$ and some family $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$.

Proof. Assume that $(K_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$. Let $\varphi \in C^\infty(\mathbb{R}^{d+1})$ be such that $\varphi(y) = 1$ near $y = 0$ and for $z \in \Omega$ let $p_z = (\varphi(y)K_z(x, y))_{y \rightarrow \xi}^\wedge$. Since $(K_z)_{z \in \Omega}$ lies in $\text{Hol}(\Omega, C^\infty(U) \hat{\otimes} \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1}))$ we see that $(p_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$ and we have

$$(5.36) \quad K_z(x, y) = (p_z)_{\xi \rightarrow y}^\vee(x, y) + (1 - \varphi(y))K_z(x, y) = (p_z)_{\xi \rightarrow y}^\vee(x, y) \mod \text{Hol}(\Omega, C^\infty(U \times U)).$$

Let us write $K_z \sim \sum_{j \geq 0} K_{j,z}$ with $(K_{j,z})_{z \in \Omega} \in \text{Hol}(\Omega, \mathcal{K}_{\text{ah}}^*(U \times \mathbb{R}^{d+1}))$ of degree $\hat{m}(z) + j$ and $\sim$ taken in the sense of (5.34). For $j = 0, 1, \dots$ we let $p_{j,z} = (\varphi(y)K_{j,z})_{z \in \Omega}$. Then arguing as in the proof of Lemma 5.23 shows that $(p_{j,z})_{z \in \Omega}$ is a family in $\text{Hol}(\Omega, S_{\text{ah}}^*(U \times \mathbb{R}^{d+1}))$ of degree $\hat{m}(z) - j$.

Next, in the sense of (5.34) we have $(p_z)_{\xi \rightarrow y}^\vee(x, y) \sim \sum_{j \geq 0} (p_{j,z})_{\xi \rightarrow y}^\vee(x, y)$. Under the Fourier transform with respect to y this shows that, for any compact $L \subset U$, any integer N and any open $\Omega' \subset \subset \Omega$, as soon as J is large enough we have estimates,

$$(5.37) \quad |\partial_x^\alpha \partial_\xi^\beta (p_z - \sum_{j \leq J} (\varphi(y)K_{j,z}(x, y))_{y \rightarrow \xi}^\wedge)(x, \xi)| \leq C_{\Omega' N J L \alpha \beta} (1 + |\xi|^2)^{-[N/2]},$$

for $(x, \xi) \in L \times \mathbb{R}^{d+1}$ and $z \in \Omega'$. Hence $p_z \sim \sum_{j \geq 0} p_{j,z}$ in the sense of (5.7). It thus follows that $(p_z)_{z \in \Omega}$ is in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ and has order $\hat{m}(z)$, so using (5.36) we see that the family $(K_z)_{z \in \Omega}$ is of the form (5.35).

Conversely, assume that $(K_z)_{z \in \Omega}$ is of the form $K_z(x, y) = (p_z)_{\xi \rightarrow y}^\vee(x, y) + R_z(x, y)$ for some family $(p_z)_{z \in \Omega}$ in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$ of order $\hat{m}(z) := -(m(z) + d + 2)$ and some family $(R_z)_{z \in \Omega}$ in $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$. By Lemma 5.16 we have $p_z \sim \sum_{j \geq 0} p_{j,z}$ with $(p_{j,z})_{z \in \Omega} \in \text{Hol}(\Omega, S_{\text{ah}}^*(U \times \mathbb{R}^{d+1}))$ of degree $m(z) - j$ and $\sim$ taken in the sense of (5.7). Thus, under the inverse Fourier transform with respect to ξ , we get an asymptotic expansion $(p_z)_{\xi \rightarrow y}^\vee \sim \sum_{j \geq 0} (p_{j,z})_{\xi \rightarrow y}^\vee$ in the sense of (5.34). As Lemma 5.23 tells us that $((p_{j,z})_{\xi \rightarrow y}^\vee)_{z \in \Omega}$ is a holomorphic family of almost homogeneous distributions of degree $\hat{m}(z) + j$, it follows that $((p_z)_{\xi \rightarrow y}^\vee)_{z \in \Omega}$ is a family in $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ of order $\hat{m}(z)$. Since $(K_z)_{z \in \Omega}$ agrees with $((p_z)_{\xi \rightarrow y}^\vee)_{z \in \Omega}$ up to an element of $\text{Hol}(\Omega, C^\infty(U \times U))$, the same is true for $(K_z)_{z \in \Omega}$. $\square$

We are now ready to prove the kernel characterization of holomorphic families of $\Psi_H \text{DO}$'s.

Proposition 5.26. Let $(P_z)_{z \in \Omega} \in \text{Hol}(\Omega, \mathcal{L}(C_c^\infty(U), C^\infty(U)))$. Then the following are equivalent:

- (i) The family $(P_z)_{z \in \Omega}$ is a holomorphic family of $\Psi_H \text{DO}$'s of order $m(z)$.

(ii) There exist $(K_z)_{z \in \Omega} \in \text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ of order $\hat{m}(z) := -(m(z) + d + 2)$ and $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U \times U))$ such that the distribution kernel $k_{P_z}(x, y)$ of P_z is of the form

$$(5.38) \quad k_{P_z}(x, y) = |\psi'_x| P_z(x, -\varepsilon_x(y)) + R_z(x, y),$$

where ψ_x denotes the coordinate change to the privileged coordinates at x .

(iii) There exist $(K_{P_z})_{z \in \Omega} \in \text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ of order $\hat{m}(z) := -(m(z) + d + 2)$ and $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U \times U))$ such that the distribution kernel $k_{P_z}(x, y)$ of P_z is of the form

$$(5.39) \quad k_{P_z}(x, y) = |\varepsilon'_x| P_z(x, -\varepsilon_x(y)) + R_z(x, y).$$

where ε_x denotes the coordinate change to the Heisenberg coordinates at x .

Proof. First, it follows from (4.27) and Proposition 5.25 that (i) and (ii) are equivalent.

Next, for $x \in U$ let ϕ_x denote the transition map from the privileged coordinates at x to the Heisenberg coordinates at x . Recall that the coordinate changes ϕ_x , $x \in U$, give rise to an action on distributions on $U \times \mathbb{R}^{d+1}$ given by

$$(5.40) \quad K(x, y) \longrightarrow \phi_x^* K(x, y), \quad \phi_x^* K(x, y) = K(x, \phi_x^{-1}(y)).$$

Since ϕ_x depends smoothly on x , this action gives rise to continuous linear isomorphisms of $C^N(U \times \mathbb{R}^{d+1})$, $N \geq 0$, and $C^\infty(U \times \mathbb{R}^{d+1})$ onto themselves, hence to analytic isomorphisms. Moreover, since $\phi_x(0) = 0$ this also yields an analytic isomorphism of $C^\infty(U) \hat{\otimes} \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1})$ onto itself. Combining this with the homogeneity property (4.31) we then deduce that (5.40) induces linear isomorphisms of $\text{Hol}(\Omega, \mathcal{K}_{\text{ah}}^*(U \times \mathbb{R}^{d+1}))$ and $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ onto themselves. Together with (4.29) this shows that the statements (ii) and (iii) are equivalent. $\square$

5.5. Holomorphic families of Ψ DO's on a general Heisenberg manifold. Let us now define holomorphic families of Ψ_H DO's on a general Heisenberg manifold.

First, the following holds.

Lemma 5.27. *Let $(K_z)_{z \in \Omega} \in \text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ and assume there exists integer N such that $\inf_{z \in \Omega} \Re \text{ord} K_z \geq 2N$. Then the family $(K_z)_{z \in \Omega}$ is contained in $\text{Hol}(\Omega, C^N(U \times \mathbb{R}^{d+1}))$.*

Proof. Thanks to Proposition 5.26 we may assume that K_z is of the form $K_z(x, y) = \check{p}_{z, \xi \rightarrow y}(x, y)$ with $(p_z)_{z \in \Omega}$ in $\text{Hol}(\Omega, S^*(U \times \mathbb{R}^{d+1}))$. As we have $-(\Re \text{ord} p_z + d + 2) = \Re \text{ord} K_z \geq 2N$ we see that $(p_z)_{z \in \Omega}$ is contained in $\text{Hol}(\Omega, S_{||}^{-(2N+d+2)}(U \times \mathbb{R}^{d+1}))$. Since the map $p \rightarrow \check{p}_{\xi \rightarrow y}$ is continuous from $S_{||}^k(U \times \mathbb{R}^{d+1})$ to $C^\infty(U) \hat{\otimes} C^N(U \times \mathbb{R}^{d+1})$ (see [Po8]), it follows that $(K_z)_{z \in \Omega}$ lies in $\text{Hol}(\Omega, C^N(U \times \mathbb{R}^{d+1}))$. $\square$

Lemma 5.28. *1) Assume $k < -(d + 2)$ and set $N = \lceil -\frac{k+d+2}{2} \rceil$. Then the map $p \rightarrow \check{p}_{\xi \rightarrow y}$ is continuous from $S_{||}^k(U \times \mathbb{R}^{d+1})$ to $C^\infty(U) \hat{\otimes} C^N(U \times \mathbb{R}^{d+1})$.*

2) For $\Re m > 0$ we have $\mathcal{K}^m(U \times \mathbb{R}^{d+1}) \subset C^\infty(U) \hat{\otimes} C^{\lceil \frac{\Re m}{2} \rceil}(\mathbb{R}^{d+1})$.

Proposition 5.29. *Let $\tilde{U}$ be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $\tilde{H} \subset T\tilde{U}$ and a $\tilde{H}$ -frame of $T\tilde{U}$ and let $\phi : (U, H) \rightarrow (\tilde{U}, \tilde{H})$ be a Heisenberg diffeomorphism. Then for any family $(\tilde{P}_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi_{\tilde{H}}^*(\tilde{U}))$ the family $(P_z)_{z \in \Omega} := (\phi^* \tilde{P}_z)_{z \in \Omega}$ is contained in $\text{Hol}(\Omega, \Psi_H^*(U))$.*

Proof. For $x \in U$ and $\tilde{x} \in \tilde{U}$ let ε_x and $\tilde{\varepsilon}_{\tilde{x}}$ denote the coordinate changes to the Heisenberg coordinates at x and $\tilde{x}$ respectively. Then by Proposition 5.26 the distribution kernel $k_{\tilde{P}_z}(\tilde{x}, \tilde{y})$ of $\tilde{P}_z$ is of the form

$$(5.41) \quad k_{\tilde{P}_z}(\tilde{x}, \tilde{y}) = |\tilde{\varepsilon}'_{\tilde{x}}| K_{\tilde{P}_z}(\tilde{x}, -\tilde{\varepsilon}_{\tilde{x}}(\tilde{y})) + \tilde{R}_z(\tilde{x}, \tilde{y}),$$

with $(K_{\tilde{P}_z})_{z \in \Omega}$ in $\text{Hol}(\Omega, \mathcal{K}^*(\tilde{U} \times \tilde{U}))$ and $(\tilde{R}_z)_{z \in \Omega}$ in $\text{Hol}(\Omega, C^\infty(\tilde{U} \times \mathbb{R}^{d+1}))$. Then the proof of Proposition 4.19 in [Po8] shows that the distribution kernel $k_{P_z}(x, y)$ of P_z takes the form,

$$(5.42) \quad k_{P_z}(x, y) = |\varepsilon'_x| K_{P_z}(x, -\varepsilon_x(y)) \\ + (1 - \chi(x, -\varepsilon_x(y))) |\tilde{\varepsilon}'_{\phi(x)}| K_{\tilde{P}_z}(\phi(x), -\tilde{\varepsilon}_{\phi(x)}(\phi(y))) + \tilde{R}_z(\phi(x), \phi(y)),$$

where we have let

$$(5.43) \quad K_{P_z}(x, y) = \chi(x, y) |\partial_y \Phi(x, y)| K_{\tilde{P}_z}(\phi(x), \Phi(x, y)), \quad \Phi(x, y) = -\tilde{\varepsilon}_{\phi(x)} \circ \phi \circ \varepsilon_x^{-1}(-y),$$

and $\chi(x, y) \in C^\infty(U \times \mathbb{R}^{d+1})$ is supported on the open subset $\mathcal{U} = \{(x, y) \in U \times \mathbb{R}^{d+1}; \varepsilon_x^{-1}(-y) \in U\}$, is properly supported with respect to x and satisfies $\chi(x, y) = 1$ near $U \times \{0\}$. In particular, we have

$$(5.44) \quad k_{P_z}(x, y) = |\varepsilon'_x| K_{P_z}(x, -\varepsilon_x(y)) \mod \text{Hol}(\Omega, C^\infty(U \times U)).$$

Let us now prove that $(K_{P_z})_{z \in \Omega}$ is an element of $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$. To this end, possibly by shrinking Ω , we may assume that $\inf_{z \in \Omega} \Re \hat{m}(z) \geq \mu > -\infty$. Moreover, the proof of Proposition 4.19 in [Po8] shows that for any integer N we have

$$(5.45) \quad K_{P_z}(x, y) = \sum_{\langle \alpha \rangle < N} \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} K_{\alpha\beta, z}(x, y) + \sum_{j=1}^3 R_{N, z}^{(j)}(x, y),$$

$$(5.46) \quad K_{\alpha\beta, z}(x, y) = a_{\alpha\beta}(x) y^\beta (\partial_y^\alpha K_{\tilde{P}_z})(\phi(x), \phi'_H(x)y),$$

where the smooth functions $a_{\alpha\beta}(x)$ are as in Proposition 4.19 and the remainder terms $R_{N, z}^{(j)}(x, y)$, $j = 1, 2, 3$, take the forms:

- $R_{N, z}^{(2)}(x, y) = \sum_{\langle \alpha \rangle < N} \sum_{\langle \beta \rangle \div \frac{3}{2}N} r_{M\alpha}(x, y) y^\beta (\partial_y^\alpha K_{\tilde{P}_z})(\phi(x), \phi'_H(x)y)$ for some functions $r_{\alpha\beta}(x, y)$ in $C^\infty(U \times \mathbb{R}^{d+1})$;

- $R_{N, z}^{(3)}(x, y) = \sum_{\langle \alpha \rangle = N} \sum_{\langle \beta \rangle \div \frac{3}{2}N} \int_0^1 r_{\alpha\beta}(t, x, y) (\tilde{y}^\beta \partial_y^\alpha K_{\tilde{P}_z})(\phi(x), t\Phi(x, y) + (1-t)\phi'_H(x)y) dt$, for some functions $r_{\alpha\beta}(t, x, y)$ in $C^\infty([0, 1] \times U \times \mathbb{R}^{d+1})$;

- $R_{N, z}^{(3)}(x, y) = \sum_{\langle \alpha \rangle < N} \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} (1 - \chi(x, y)) K_{\alpha\beta, z}(x, y)$.

Observe that the map $\Phi(x, y) = (\phi(x), \phi'_H(x)y)$ is a smooth diffeomorphism from $U \times \mathbb{R}^{d+1}$ onto $\tilde{U} \times \mathbb{R}^{d+1}$ such that $\Phi(x, 0) = (\phi(x), 0)$ and $\Phi(x, \lambda y) = (\phi(x), \lambda \phi'_H(x)y)$, so along similar lines as that of the proof of Proposition 5.26 we can prove that the map

$$(5.47) \quad \mathcal{D}'(U \times \mathbb{R}^{d+1}) \ni K(x, y) \longrightarrow K(\phi(x), \phi'_H(x)y) \in \mathcal{D}'(\tilde{U} \times \mathbb{R}^{d+1})$$

gives rise to a linear map from $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ to $\text{Hol}(\Omega, \mathcal{K}^*(\tilde{U} \times \mathbb{R}^{d+1}))$ preserving the order. Therefore, the family $(K_{\alpha\beta, z}(x, y))_{z \in \Omega}$ is contained in $\text{Hol}(\Omega, \mathcal{K}^*(\tilde{U} \times \mathbb{R}^{d+1}))$ and has order $\hat{m}(z) + \langle \beta \rangle - \langle \alpha \rangle$. Incidentally, the term $(R_{N, z}^{(3)})_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$.

On the other hand, if $\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle \div \frac{3}{2}N$ then the order $\hat{m}_{\alpha\beta}(z) = \hat{m}(z) + \langle \beta \rangle - \langle \alpha \rangle$ of the family $(\tilde{y}^\beta \partial_y^\alpha K_{\tilde{P}_z})_{z \in \Omega} \in \text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ satisfies $\Re \hat{m}_{\alpha\beta}(z) \geq \Re \hat{m}(z) + \frac{1}{3}\langle \beta \rangle \geq \mu + \frac{N}{2}$. Therefore, it follows from Lemma 5.27 that, for any integer J , as soon as N is large enough $(\tilde{y}^\beta \partial_y^\alpha K_{\tilde{P}_z})_{z \in \Omega}$ is in $\text{Hol}(\Omega, C^J(\tilde{U} \times \mathbb{R}^{d+1}))$, as much so the remainder terms $(R_{N, z}^{(2)})_{z \in \Omega}$ and $(R_{N, z}^{(3)})_{z \in \Omega}$ are in $\text{Hol}(\Omega, C^J(U \times \mathbb{R}^{d+1}))$.

All this shows that in the sense of (5.34) we have $K_{P_z}(x, y) \sim \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle} K_{\alpha\beta, z}(x, y)$, which implies that $(K_{P_z})_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$. Combining this with (5.44) and Proposition 5.26 then shows that $(P_z)_{z \in \Omega}$ is a holomorphic family of $\Psi_H\text{DO}$'s. $\square$

Now, let (M^{d+1}, H) be a Heisenberg manifold and let $\mathcal{E}$ be a smooth vector bundle over M . Then Proposition 5.29 allows us to define holomorphic families with values in $\Psi_H^*(M, \mathcal{E})$ as follows.

Definition 5.30. A family $(P_z)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E})$ is holomorphic when:

- (i) The order $m(z)$ of P_z is a holomorphic function of z ;
- (ii) For φ and ψ in $C_c^\infty(M)$ with disjoint supports $(\varphi P_z \psi)_{z \in \Omega}$ is a holomorphic family of smoothing operators (i.e. is given by a holomorphic family of smooth distribution kernels);
- (iii) For any trivialization $\tau : \mathcal{E}|_U \rightarrow U \times \mathbb{C}^r$ over a local Heisenberg chart $\kappa : U \rightarrow V \subset \mathbb{R}^{d+1}$ the family $(\kappa_* \tau_*(P_z|_U))_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, \Psi_H^*(V, \mathbb{C}^r)) := \text{Hol}(\Omega, \Psi_H^*(V)) \otimes \text{End } \mathbb{C}^r$.

All the preceding properties of holomorphic families of $\Psi_H \text{DO}$'s on an open subset of $\mathbb{R}^{d+1}$ hold *verbatim* for holomorphic families with values in $\Psi_H^*(M, \mathcal{E})$. Moreover, we have:

Proposition 5.31. The principal symbol map $\sigma_* : \Psi_H^*(M, \mathcal{E}) \rightarrow S_*(\mathfrak{g}^*M, \mathcal{E})$ is analytic, in the sense that for any holomorphic family $(P_z)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E})$ the family of symbols $(\sigma_*(P_z))_{z \in \Omega}$ is in $\text{Hol}(\Omega, C^\infty(\mathfrak{g}^*M \setminus 0, \text{End } \mathcal{E}))$.

Proof. Let $(P_z)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E})$ be a holomorphic family of $\Psi_H \text{DO}$'s of order $m(z)$ and let us show that the family of symbols $(\sigma_*(P_z))_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(\mathfrak{g}^*M \setminus 0, \text{End } \mathcal{E}))$. Since this is a purely local issue we may as well assume that $(P_z)_{z \in \Omega}$ is a holomorphic family of scalar $\Psi_H \text{DO}$'s on a local trivializing Heisenberg chart $U \subset \mathbb{R}^{d+1}$.

By Proposition 5.26 we can put the distribution kernel of P_z into the form,

$$(5.48) \quad k_{P_z}(x, y) = |\varepsilon'_x| K_{P_z}(x, -\varepsilon_x(y)) + R_z(x, y),$$

with $(K_{P_z})_{z \in \Omega} \in \text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ of order $\hat{m}(z) = -(m(z) + d + 2)$ and $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, C^\infty(U \times U))$. Let $\varphi \in C^\infty(\mathbb{R}^{d+1})$ be such that $\varphi(y) = 1$ near $y = 0$ and let $p_z = (\varphi(y) K_{P_z}(x, y))_{\xi \rightarrow y}^\wedge$. Then the proof of Proposition 5.25 shows that $(p_z)_{z \in \Omega}$ is a holomorphic family of symbols. Moreover, we have

$$(5.49) \quad K_{P_z}(x, y) = (p_z)_{\xi \rightarrow y}^\vee(x, y) + (1 - \varphi(y)) K_{P_z}(x, y) = (p_z)_{\xi \rightarrow y}^\vee(x, y) \mod \text{Hol}(\Omega, C^\infty(U \times U)).$$

Let $z \in \Omega$ and let $K_{\hat{m}(z)} \in \mathcal{K}_{\hat{m}(z)}(U \times \mathbb{R}^{d+1})$ be the principal kernel of K_{P_z} . Then (5.49) and Proposition 4.15 show that the leading symbol of p_z is the restriction to $U \times (\mathbb{R}^{d+1} \setminus 0)$ of $(K_{\hat{m}(z)})_{y \rightarrow \xi}^\wedge$. Since the latter is equal to $\sigma_{m(z)}(P_z)$, we see that the leading symbol of p_z is just $\sigma_{m(z)}(P_z)$. Since $(p_z)_{z \in \Omega}$ is a holomorphic family of symbols it then follows from Remark 5.8 that the family $(\sigma_{m(z)}(P_z))_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, C^\infty(U \times (\mathbb{R}^{d+1} \setminus 0)))$. The proof is thus achieved. $\square$

5.6. Transposition and holomorphic families of $\Psi_H \text{DO}$'s. Let us now look at the analyticity and anti-analyticity of taking transposes and adjoints of $\Psi_H \text{DO}$'s.

Proposition 5.32. Let $(P_z)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E})$ be a holomorphic family of $\Psi_H \text{DO}$'s. Then the transpose family $(P_z^t)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E}^*)$ is a holomorphic family of $\Psi_H \text{DO}$'s.

Proof. For $z \in \Omega$ let $k_{P_z}(x, y)$ denote the distribution kernel of P_z . The distribution kernel of P_z^t is $k_{P_z^t}(x, y) = k_{P_z}(y, x)^t$, hence is represented outside the diagonal by a holomorphic family of smooth kernels. Therefore, we need only to prove the statement for a holomorphic family of scalar $\Psi_H \text{DO}$'s on a Heisenberg chart $U \subset \mathbb{R}^{d+1}$, as we shall now suppose that the family $(P_z)_{z \in \Omega}$ is. In addition, there is no loss of generality in assuming that the order $m(z)$ of P_z is such that there exists $\mu \in \mathbb{R}$ so that $\Re \hat{m}(z) \geq \mu$ for any $z \in \Omega$.

Next, thanks to Proposition 5.26 the kernel of P_z is of the form

$$(5.50) \quad k_{P_z} = |\varepsilon'_x| K_{P_z}(x, -\varepsilon_x(y)) + R_z(x, y),$$

with $(K_{P_z})_{z \in \Omega}$ in $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$ and $(R_z)_{z \in \Omega}$ in $\text{Hol}(\Omega, C^\infty(U \times U))$. Then the proof of Proposition 4.21 in [Po8] shows that we can write

$$(5.51) \quad k_{P_z}(x, y) = |\varepsilon'_x| K_{P_z^t}(x, -\varepsilon_x(y)) + (1 - \chi(x, -\varepsilon_x(y))) |\varepsilon'_y| K_{P_z}(y, -\varepsilon_y(x)) + R_z(y, x),$$

where we have let

$$(5.52) \quad K_{P_z^t}(x, y) = \chi(x, y) |\varepsilon'_x|^{-1} |\varepsilon'_y| K_{P_z}(\varepsilon_x^{-1}(-y), -\varepsilon_{\varepsilon_x^{-1}(-y)}(x)),$$

and the function $\chi(x, y) \in C^\infty(U \times \mathbb{R}^{d+1})$ is supported in the open subset $\mathcal{U} = \{(x, y) \in U \times \mathbb{R}^{d+1}; \varepsilon_x^{-1}(-y) \in U\}$ and is in such way to be properly supported with respect x and to be equal to 1 near $U \times \{0\}$. In particular, we have

$$(5.53) \quad k_{P_z^t}(x, y) = |\varepsilon'_x| K_{P_z^t}(x, -\varepsilon_x(y)) \quad \text{mod } \text{Hol}(\Omega, C^\infty(U \times U)).$$

Moreover, as shown in [Po8], for any integer N we have

$$(5.54) \quad K_{P_z^t}(x, y) = \sum_{\alpha, \beta, \gamma, \delta}^{(N)} K_{\alpha\beta\gamma\delta, z} + \sum_{j=1}^4 R_{N, z}(x, y),$$

$$(5.55) \quad K_{\alpha\beta\gamma\delta, z} = a_{\alpha\beta\gamma\delta}(x) y^{\beta+\delta} (\partial_x^\gamma \partial_y^\alpha K_{P_z})(x, -y)$$

where the smooth functions $a_{\alpha\beta\gamma\delta}(x)$ are as in Proposition 4.21, the summation goes over all the multi-orders α, β, γ and δ such that $\langle \alpha \rangle < N$, $\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N$ and $|\gamma| \leq |\delta| \leq 2|\gamma| < 2N$ and the remainder terms $R_{N, z}^{(j)}(x, y)$ take the forms:

- $R_{N, z}^{(1)} = \sum_{\langle \alpha \rangle = N} \sum_{\langle \beta \rangle \doteq \frac{3}{2}\langle \alpha \rangle} |\varepsilon'_x|^{-1} |\varepsilon'_y| \int_0^1 r_{N\alpha\beta}(t, x, y) (y^\beta \partial_y^\alpha K_{P_z})(\varepsilon_x^{-1}(-y), \Phi_t(x, y))$, where the functions $r_{N\alpha\beta}(t, x, y)$ are in $C^\infty([0, 1] \times U \times \mathbb{R}^{d+1})$, the equality $k \doteq \frac{3}{2}l$ means that k is equal to $\frac{3}{2}l$ if $\frac{3}{2}l$ is integer and to $\frac{3}{2}l + \frac{1}{2}$ otherwise, and we have let $\Phi_t(x, y) = -y + t(y - \varepsilon_{\varepsilon_x^{-1}(-y)}(x))$;

- $R_{N, z}^{(2)}(x, y) = \sum_{\langle \beta \rangle \doteq \frac{3}{2}N} r_{N\alpha}(x, y) y^\beta (\partial_y^\alpha K_{P_z})(\varepsilon_x^{-1}(-y), -y)$ with $r_{N\alpha}(x, y)$ in $C^\infty(U \times \mathbb{R}^{d+1})$;

- $R_{N, z}^{(3)}(x, y) = \sum_{|\gamma|=N} \sum_{N \leq |\delta| \leq 2N} a_{\alpha\beta\gamma\delta}(x) y^{\beta+\delta} \int_0^1 (1-t)^{N-1} (\partial_x^\gamma \partial_y^\alpha K_{P_z})(\varepsilon_t(x, y), -y)$;

- $R_{N, z}^{(4)}(x, y) = \sum_{\alpha, \beta, \gamma, \delta}^{(N)} (1 - \chi(x, y)) K_{\alpha\beta\gamma\delta, z}$.

Each family $(K_{\alpha\beta\gamma\delta, z})_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$. Moreover, the remainder term $R_{N, z}^{(4)}$ belongs to $\text{Hol}(\Omega, C^\infty(U \times \mathbb{R}^{d+1}))$ and, along similar lines as that of the proof of Proposition 5.29, we can show that for any integer J the other remainder terms $(R_{N, z}^{(j)})_{z \in \Omega}$ are in $\text{Hol}(\Omega, C^J(U \times \mathbb{R}^{d+1}))$ as soon N is large enough. Therefore, we have $K_{P_z^t} \sim \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle} \sum_{|\gamma| \leq |\delta| \leq 2|\gamma|} K_{\alpha\beta\gamma\delta, z}$ in the sense of (5.54), which means that $K_{P_z^t}$ belongs to $\text{Hol}(\Omega, \mathcal{K}^*(U \times \mathbb{R}^{d+1}))$. Combining this with (5.53) and Proposition 5.26 shows that $(P_z^t)_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's. $\square$

Assume now that M is endowed with a density > 0 and $\mathcal{E}$ with a Hermitian metric. Then Proposition 5.32 allows us to carried out the proof of Proposition 4.31 in the holomorphic setting as much so to get:

Proposition 5.33. *Let $(P_z)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E})$ be a holomorphic family of Ψ_H DO's. Then the family $(P_z^*)_{z \in \Omega} \subset \Psi_H^*(M, \mathcal{E}^*)$ is an anti-holomorphic family of Ψ_H DO's, in the sense that $(P_z^*)_{z \in \Omega}$ is a holomorphic family of Ψ_H DO's.*

6. COMPLEX POWERS OF HYPOELLIPTIC DIFFERENTIAL OPERATORS

In this section we will show that the complex powers of a positive hypoelliptic differential operator, *a priori* defined as unbounded operators on $L^2(M, \mathcal{E})$, give rise to a holomorphic family of Ψ_H DO's.

Since we cannot carry out in the Heisenberg setting the standard approach of Seeley [Se] to the complex powers of elliptic operators, we will rely on a new approach based on the pseudodifferential representation of the heat kernel of [BGS], which extended to the Heisenberg setting the results of [Gre].

This section is divided into two subsections. In the first one we recall the pseudodifferential representation of the heat kernel of a hypoelliptic operator of [BGS], and in the second one we deal with the complex powers of a positive hypoelliptic operator.

Throughout this section we let (M^{d+1}, H) be a compact Heisenberg manifold endowed with a positive density and let $\mathcal{E}$ be a Hermitian vector bundle over M of rank r .

6.1. Pseudodifferential representation of the heat kernel. Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of even (Heisenberg) order v which is bounded from below and has an invertible principal symbol. In particular, P is hypoelliptic with loss of $\frac{v}{2}$ -derivatives by Theorem 4.32.

Since P is bounded from below it generates on $L^2(M, \mathcal{E})$ a heat semigroup e^{-tP} , $t \geq 0$. In fact, for $t > 0$ and for any integer $k \geq 1$ the operators $P^k e^{-tP}$ and $e^{-tP} P^k$ coincide and are bounded. Thus e^{-tP} maps continuously $L^2(M, \mathcal{E})$ to $\cap_{k \geq 0} \text{dom } P^k$, which is just $C^\infty(M, \mathcal{E})$ since P is hypoelliptic with loss of $\frac{v}{2}$ -derivatives.

Moreover, as e^{-tP} is selfadjoint it defines by duality a continuous map from $\mathcal{D}'(M, \mathcal{E})$ to $L^2(M, \mathcal{E})$. Since $e^{-tP} = e^{-tP/2} e^{-tP/2}$ it follows that e^{-tP} extends to a continuous map from $\mathcal{D}'(M, \mathcal{E})$ to $C^\infty(M, \mathcal{E})$, that is e^{-tP} is smoothing. In particular, e^{-tP} is given by a smooth kernel $k_t(x, y)$ in $C^\infty(M \times M, \mathcal{E} \boxtimes (\mathcal{E} \otimes |\Lambda|(M)))$, where $|\Lambda|(M)$ denotes the bundle of densities on M .

On the other hand, the heat semi-group allows us to invert the heat equation. Indeed, the operator given by

$$(6.1) \quad Q_0 f(x, t) = \int_0^\infty e^{-sP} f(x, t-s) ds, \quad f \in C_c^\infty(M \times \mathbb{R}, \mathcal{E}),$$

maps continuously into $C^0(\mathbb{R}, L^2(M, \mathcal{E})) \subset \mathcal{D}(M \times \mathbb{R}, \mathcal{E})$ and satisfies

$$(6.2) \quad (P + \partial_t) Q_0 f = Q_0 (P + \partial_t) f = f \quad \forall u \in C_c^\infty(M \times \mathbb{R}, \mathcal{E}).$$

Notice that the operator Q_0 has the *Volterra property* of [Pi], i.e. it has a distribution kernel of the form $K_{Q_0}(x, y, t-s)$ with $K_{Q_0}(x, y, t)$ supported outside the region $\{t < 0\}$. Indeed, at the level of distribution kernels the formula (6.1) implies that we have

$$(6.3) \quad K_{Q_0}(x, y, t) = \begin{cases} k_t(x, y) & \text{if } t > 0, \\ 0 & \text{if } t < 0. \end{cases}$$

The above equalities are the main motivation for using pseudodifferential techniques to study the heat kernel $k_t(x, y)$. The idea is to consider a class of Ψ_H DO's, the Volterra Ψ_H DO's, taking into account:

- (i) The aforementioned Volterra property;
- (ii) The parabolic homogeneity of the heat operator $P + \partial_t$, i.e. the homogeneity with respect to the dilations,

$$(6.4) \quad \lambda.(\xi, \tau) = (\lambda.\xi, \lambda^v \tau), \quad (\xi, \tau) \in \mathbb{R}^{d+2}, \quad \lambda \neq 0.$$

In the sequel for $g \in \mathcal{S}'(\mathbb{R}^{d+2})$ and $\lambda \neq 0$ we let g_λ be the tempered distribution defined by

$$(6.5) \quad \langle g_\lambda(\xi, \tau), f(\xi, \tau) \rangle = |\lambda|^{-(d+2+v)} \langle g(\xi, \tau), f(\lambda^{-1}\xi, \lambda^{-v}\tau) \rangle, \quad f \in \mathcal{S}(\mathbb{R}^{d+2}).$$

Definition 6.1. A distribution $g \in \mathcal{S}'(\mathbb{R}^{d+2})$ is parabolic homogeneous of degree m , $m \in \mathbb{Z}$, when we have $g_\lambda = \lambda^m g$ for any $\lambda \neq 0$.

Let $\mathbb{C}_-$ denote the complex halfplane $\{\Im \tau < 0\}$ with closure $\overline{\mathbb{C}_-} \subset \mathbb{C}$. Then:

Lemma 6.2 ([BGS, Prop. 1.9]). Let $q(\xi, \tau) \in C^\infty((\mathbb{R}^{d+1} \times \mathbb{R}) \setminus 0)$ be parabolic homogeneous of degree m and such that:

(i) q extends to a smooth function on $(\mathbb{R}^{d+1} \times \overline{\mathbb{C}_-}) \setminus 0$ which restricts on $\mathbb{R}^{d+1} \times \mathbb{C}_-$ to an element of $C^\infty(\mathbb{R}^{d+1}) \hat{\otimes} \text{Hol}(\mathbb{C}_-)$.

Then there exists a unique distribution $g \in \mathcal{S}'(\mathbb{R}^{d+2})$ agreeing with q on $\mathbb{R}^{d+1} \setminus 0$ and such that:

- (ii) g is parabolic homogeneous of degree m ;
- (iii) The inverse Fourier transform $\check{g}(x, t)$ vanishes for $t < 0$.

Let U be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $H \subset TU$ and H -frame $X_0, \dots, X_d$ of TU . We define Volterra symbols and Volterra $\Psi_H \text{DO}$'s as follows.

Definition 6.3. $S_{v,m}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$, $m \in \mathbb{Z}$, consists of functions $q(x, \xi, \tau)$ in $C^\infty(U \times (\mathbb{R}^{d+2} \setminus 0))$ such that:

- (i) $q(x, \lambda \xi, \lambda^v \tau) = \lambda^m q(x, \xi, \tau)$ for any $(x, \xi, \tau) \in U \times (\mathbb{R}^{d+2} \setminus 0)$ and any $\lambda \neq 0$;
- (ii) $q(x, \xi, \tau)$ extends to an element of $C^\infty(U \times [(\mathbb{R}^{d+1} \times \overline{\mathbb{C}_-}) \setminus 0])$ in such way that its restricts to an element of $C^\infty(U \times \mathbb{R}^{d+1}) \hat{\otimes} \text{Hol}(\mathbb{C}_-)$ on $U \times \mathbb{R}^{d+1} \times \mathbb{C}_-$.

Definition 6.4. $S_v^m(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$, $m \in \mathbb{Z}$, consists of functions $q(x, \xi, \tau)$ in $C^\infty(U \times \mathbb{R}^{d+1} \times \mathbb{R})$ with an asymptotic expansion $q \sim \sum_{j \geq 0} q_{m-j}$, $q_l \in S_{v,m}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$, in the sense that, for any integer N and any compact $K \subset \bar{U}$, we have

$$(6.6) \quad |\partial_x^\alpha \partial_\xi^\beta \partial_\tau^k (q - \sum_{j < N} q_{m-j})(x, \xi, \tau)| \leq C_{NK\alpha\beta k} (\|\xi\| + |\tau|^{1/v})^{m-N-(\beta)-vk},$$

for $x \in K$ and $|\xi| + |\tau|^{1/v} > 1$.

Definition 6.5. Let $q(x, \xi, \tau) \in S_{v,m}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$ and let $g \in C^\infty(U) \hat{\otimes} \mathcal{S}'(\mathbb{R}^{d+1})$ be the unique homogeneous extension of q provided by Lemma 6.2. Then we let $\check{q}_m(x, y, t)$ denote the inverse Fourier transform of $g(x, \xi, \tau)$ with respect to the variables (ξ, τ) .

Remark 6.6. The above definition makes sense since it follows from the proof of Lemma 6.2 in [BGS] that the extension process of Lemma 6.2 applied to every symbol $q(x, \cdot, \cdot)$, $x \in U$, is smooth with respect to x , so really gives rise to an element of $C^\infty(U) \hat{\otimes} \mathcal{S}'(\mathbb{R}^{d+2})$.

Definition 6.7. $\Psi_{H,v}^m(U \times \mathbb{R}_{(v)})$, $m \in \mathbb{Z}$, consists of continuous operators $Q : C_c^\infty(U_x \times \mathbb{R}_t) \rightarrow C^\infty(U_x \times \mathbb{R}_t)$ such that Q has the Volterra property and can be put into the form

$$(6.7) \quad Q = q(x, -iX, D_t) + R$$

with q in $S_v^m(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$ and R in $\Psi^{-\infty}(U \times \mathbb{R})$.

It is immediate to extend the properties of $\Psi_H \text{DO}$'s on U alluded to in Section 4 to Volterra $\Psi_H \text{DO}$'s on $U \times \mathbb{R}$ except for the asymptotic completeness as in Lemma 4.7, which is crucial for constructing. The problem is that the cut-off arguments of the classical proof the asymptotic completeness of standard ΨDO 's cannot be carried through in Volterra setting because we require analyticity with respect to the time covariable. A proof of the asymptotic completeness of Volterra ΨDO 's is given in [Pi], but simpler proofs which can be carried out *verbatim* for Volterra $\Psi_H \text{DO}$'s can be found in [Po5].

On the other hand, thanks to the Volterra property the kernels of $\Psi_H \text{DO}$'s can be characterized as follows.

Definition 6.8. $\mathcal{K}_{v,m}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$, $m \in \mathbb{Z}$, consists of distributions $K(x, y, t)$ in $C^\infty(U) \otimes \mathcal{S}'_{reg}(\mathbb{R}^{d+2})$ such that:

- (i) The support of $K(x, y, t)$ is contained in $U \times \mathbb{R}^{d+1} \times \mathbb{R}_+$;
- (ii) $K(x, y, t)$ is parabolic homogeneous of degree m with respect to the variables (y, t) .

Definition 6.9. $\mathcal{K}_v^m(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$, $m \in \mathbb{Z}$, is the space of distributions $K(x, y, t)$ in $\mathcal{D}'(U \times \mathbb{R}^{d+2})$ which admit an asymptotic expansion $K \sim \sum_{j \geq 0} K_{m+j}$, $K_{m+j} \in \mathcal{K}_{v,m+j}(U \times \mathbb{R}^{d+2})$, where $\sim$ is taken in the sense of (4.24).

In the sequel, for $x \in U$ we let ψ_x and ε_x respectively denote the changes of variable to the privileged coordinates and the Heisenberg coordinates at x . Then, along the same lines as that of the proofs of Proposition 4.16 and Proposition 4.17, we obtain the following characterization of Volterra Ψ_H DO's.

Proposition 6.10. Let $Q : C_c^\infty(U_x \times \mathbb{R}_t) \rightarrow C^\infty(U_x \times \mathbb{R}_t)$ be a continuous operator with distribution kernel $k_Q(x, t; y, s)$. Then the following are equivalent:

- (i) The operator Q belongs to $\Psi_{H,v}^m(U \times \mathbb{R})$;
- (ii) The kernel of Q can be put into the form,

$$(6.8) \quad k_Q(x, t; y, s) = |\psi'_x| K(x, -\psi_x(y), t - s) + R(x, y, t - s),$$

for some $K(x, y, t) \in \mathcal{K}_v^{\hat{m}}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$, $\hat{m} = -(m + d + 2 + v)$ and $R \in C^\infty(U \times \mathbb{R}^{d+1} \times \mathbb{R})$.

- (iii) The kernel of Q can be put into the form,

$$(6.9) \quad k_Q(x, t; y, s) = |\varepsilon'_x| K_Q(x, -\varepsilon_x(y), t - s) + R(x, y, t - s),$$

for some $K_Q(x, y, t) \in \mathcal{K}_v^{\hat{m}}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$, $\hat{m} = -(m + d + 2 + v)$ and $R \in C^\infty(U \times \mathbb{R}^{d+1} \times \mathbb{R})$.

An interesting consequence of Proposition 6.10 is the following small time asymptotics for the kernel of a Volterra Ψ_H DO.

Proposition 6.11 ([BGS, Thm. 4.5]). Let $Q \in \Psi_{H,v}^m(U \times \mathbb{R}_{(v)})$ have symbol $q \sim \sum_{j \geq 0} q_{m-j}$ and kernel $k_Q(x, y, t - s)$. Then as $t \rightarrow 0^+$ the following asymptotics holds in $C^\infty(U)$,

$$(6.10) \quad k_Q(x, x, t) \sim t^{-\frac{2[\frac{m}{2}] + d + 4}{v}} \sum_{j \geq 0} t^j |\varepsilon'_x| (q_{2[\frac{m}{2}] - 2j})_{(\xi, \tau) \rightarrow (y, t)}^\vee(x, 0, 1).$$

On the other hand, using Proposition 6.10 and arguing along similar lines that of the proof of Proposition 4.19 in [Po8] allows us to prove:

Proposition 6.12. Let U (resp. $\tilde{U}$) be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $H \subset TU$ (resp. $\tilde{H} \subset T\tilde{U}$) and a H -frame of TU (resp. a $\tilde{H}$ -frame of $T\tilde{U}$). Let $\phi : (U, H) \rightarrow (\tilde{U}, \tilde{H})$ be a Heisenberg diffeomorphism and let $\tilde{Q} \in \Psi_{\tilde{H},v}^m(\tilde{U} \times \mathbb{R}_{(v)})$.

1) The operator $Q = (\phi \oplus 1_{\mathbb{R}})^* \tilde{Q} : C_c^\infty(U \times \mathbb{R}^{d+1} \times \mathbb{R}) \rightarrow C^\infty(U \times \mathbb{R}^{d+1} \times \mathbb{R})$ belongs to $\Psi_{H,v}^m(U \times \mathbb{R}_{(v)})$.

2) If the distribution kernel of $\tilde{P}$ is of the form (6.9) with $K_{\tilde{Q}}(\tilde{x}, \tilde{y}, t) \in \mathcal{K}_v^{\hat{m}}(\tilde{U} \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$ then the distribution kernel of P can be written in the form (6.9) with $K_Q(x, y, t) \in \mathcal{K}_v^{\hat{m}}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$ such that

$$(6.11) \quad K_Q(x, y, t) \sim \sum_{\langle \beta \rangle \geq \frac{3}{2} \langle \alpha \rangle} \frac{1}{\alpha! \beta!} a_{\alpha\beta}(x) y^\beta (\partial_y^\beta K_{\tilde{Q}})(\phi(x), \phi'_H(x)y, t),$$

where the functions $a_{\alpha\beta}(x)$ are as in (4.35).

This allows us to define Volterra Ψ_H DO's on the manifold $M \times \mathbb{R}$ and acting on the sections of the bundle $\mathcal{E}$ (or rather on the sections of the pullback of $\mathcal{E}$ by the projection $M \times \mathbb{R} \rightarrow M$, again denoted $\mathcal{E}$).

Definition 6.13. $\Psi_{H,v}^m(M \times \mathbb{R}_{(v)}, \mathcal{E})$, $m \in \mathbb{Z}$, consists of continuous operators $Q : C_c^\infty(M \times \mathbb{R}, \mathcal{E}) \rightarrow C^\infty(M \times \mathbb{R}, \mathcal{E})$ such that:

- (i) Q has the Volterra property;
- (ii) The distribution kernel of Q is smooth off the diagonal of $(M \times \mathbb{R}) \times (M \times \mathbb{R})$;
- (iii) For any trivialization $\tau : \mathcal{E}|_U \rightarrow U \times \mathbb{C}^r$ of $\mathcal{E}$ over a local Heisenberg chart $\kappa : U \rightarrow V \subset \mathbb{R}^{d+1}$ the operator $(\kappa \otimes \text{id})_* \tau_*(Q|_{U \times \mathbb{R}})$ belongs to $\Psi_{H,v}^m(V \times \mathbb{R}_{(v)}, \mathbb{C}^r) := \Psi_{H,v}^m(V \times \mathbb{R}_{(v)}) \otimes \text{End } \mathbb{C}^r$.

Using Proposition 6.12 we can define the global principal symbol of a Volterra Ψ_H DO as follows. Let $\mathfrak{g}^*M$ denote the dual bundle of the Lie algebra bundle $\mathfrak{g}M$ of M and consider the canonical projection $\pi : \mathfrak{g}^*M \times \mathbb{R} \rightarrow M$.

Definition 6.14. $S_{v,m}(\mathfrak{g}^*M \times \mathbb{R}_{(v)}, \mathcal{E})$, $m \in \mathbb{Z}$, is the subspace of $C^\infty((\mathfrak{g}^*M \times \mathbb{R}) \setminus 0, \pi^* \text{End } \mathcal{E})$ consisting of sections $q(x, \xi, \tau)$ such that:

- (i) $q(x, \lambda\xi, \lambda^v\tau) = \lambda^m q(x, \xi, \tau)$ for any $(x, \xi, \tau) \in (\mathfrak{g}^*M \times \mathbb{R}) \setminus 0$ and any $\lambda \in \mathbb{R} \setminus 0$;
- (ii) $q(x, \xi, \tau)$ extends to a section in $C^\infty((\mathfrak{g}^*M \times \bar{\mathbb{C}}_-) \setminus 0, \pi^* \text{End } \mathcal{E})$ which restricts to an element of $C^\infty(\mathfrak{g}^*M, \pi^* \text{End } \mathcal{E}) \hat{\otimes} \text{Hol}(\mathbb{C}_-)$ on $\mathfrak{g}^*M \times \mathbb{C}_-$.

Using (6.11) and arguing as in the proof of Proposition 4.24 in [Po8] we get:

Proposition 6.15. For any $Q \in \Psi_{H,v}^m(M \times \mathbb{R}_{(v)}, \mathcal{E})$ there is a unique symbol $\sigma_m(Q) \in S_{v,m}(\mathfrak{g}^*M \times \mathbb{R}_{(v)}, \mathcal{E})$ such that if in a local trivializing Heisenberg chart $U \subset \mathbb{R}^{d+1}$ we let $K_{Q,\hat{m}}(x, y, t) \in \mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$ be the leading kernel for the kernel $K_Q(x, y, t)$ in the form (6.9) for Q , then we have

$$(6.12) \quad \sigma_m(Q)(x, \xi, \tau) = [K_{Q,\hat{m}}]_{(y,t) \rightarrow (\xi,\tau)}^\wedge(x, \xi, \tau), \quad (x, \xi) \in U \times \mathbb{R}^{d+1} \setminus 0.$$

Equivalently, on any trivializing Heisenberg coordinates centered at $a \in M$ the symbol $\sigma_m(Q)(a, \cdot, \cdot)$ coincides with the (local) principal symbol of Q at $x = 0$.

Definition 6.16. For $Q \in \Psi_{H,v}^m(M \times \mathbb{R}_{(v)}, \mathcal{E})$ the symbol $\sigma_m(Q) \in S_{v,m}(\mathfrak{g}^*M \times \mathbb{R}, \mathcal{E})$ provided by Proposition 6.15 is called the (global) principal symbol of Q .

Granted this we can define the model operator of a Volterra Ψ_H DO as follows.

Definition 6.17. Let $Q \in \Psi_{H,v}^m(M \times \mathbb{R}_{(v)}, \mathcal{E})$ have principal symbol $\sigma_m(Q)$. Then the model operator of Q at $a \in M$ is the left-convolution operator by $\sigma_m(Q)_{(\xi,\tau) \rightarrow (y,t)}^\vee(a, \cdot, \cdot)$, i.e. the continuous endomorphism $Q^a : \mathcal{S}(G_a M \times \mathbb{R}, \mathcal{E}_a) \rightarrow \mathcal{S}(G_a M \times \mathbb{R}, \mathcal{E}_a)$ given by

$$(6.13) \quad Q^a u(y, t) = \langle \sigma_m(Q)_{(\xi,\tau) \rightarrow (y,t)}^\vee(x, z, t), u(y \cdot z^{-1}, t - s) \rangle, \quad u \in \mathcal{S}(G_a M, \mathcal{E}_a).$$

Remark 6.18. The model operator Q^a can be defined as an endomorphism of $\mathcal{S}(G_a M, \mathcal{E}_a)$, not just as an endomorphism of $\mathcal{S}_0(G_a M, \mathcal{E}_a)$ as in Definition 4.27, because $\sigma_m(Q)_{(\xi,\tau) \rightarrow (y,t)}^\vee(a, \cdot, \cdot)$ makes sense as an element of $\mathcal{S}'(G_a M, \mathcal{E}_a)$.

Proposition 6.19. The group laws on the fibers of $GM \times \mathbb{R}$ give rise to a convolution product,

$$(6.14) \quad * : S_{v,m_1}(\mathfrak{g}^*M \times \mathbb{R}_{(v)}, \mathcal{E}) \times S_{v,m_2}(\mathfrak{g}^*M \times \mathbb{R}_{(v)}, \mathcal{E}) \longrightarrow S_{v,m_1+m_2}(\mathfrak{g}^*M \times \mathbb{R}_{(v)}, \mathcal{E}),$$

such that for any symbols $q_{m_j} \in S_{v,m_j}(\mathfrak{g}^*M \times \mathbb{R}_{(v)}, \mathcal{E})$, $j = 1, 2$, we have

$$(6.15) \quad q_{m_1} * q_{m_2}(x, \xi, \tau) = [q_{m_1}(x, \cdot, \cdot) *^x q_{m_2}(x, \cdot, \cdot)](\xi, \tau), \quad (x, \xi, \tau) \in (\mathfrak{g}^*M \times \mathbb{R}) \setminus 0,$$

where $*^x$ denote the convolution product for symbols on $G_x M \times \mathbb{R}$.

In a local trivializing Heisenberg chart the symbolic calculus for Volterra Ψ_H DO's reduces the existence of a Volterra Ψ_H DO parametrix to the invertibility of the local and global principal symbols. Therefore, we obtain:

Proposition 6.20. *Let $Q \in \Psi_{H,v}^m(M \times \mathbb{R}_{(v)}, \mathcal{E})$, $m \in \mathbb{Z}$. Then we have equivalence:*

(i) *The principal symbol of Q is invertible with respect to the product (6.14) of Volterra-Heisenberg symbols;*

(ii) *The operator Q admits a parametrix in $\Psi_{H,v}^{-m}(M \times \mathbb{R}_{(v)}, \mathcal{E})$.*

In the case of the heat operator $P + \partial_t$, comparing a parametrix with the inverse (6.1) and using (6.3) allows us to obtain the pseudodifferential representation of the heat kernel of P below.

Proposition 6.21 ([BGS, pp. 362–363]). *Suppose that the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol. Then:*

1) *The heat operator $P + \partial_t$ has an inverse $(P + \partial_t)^{-1}$ in $\Psi_{H,v}^{-v}(M \times \mathbb{R}_{(v)}, \mathcal{E})$.*

2) *Let $K_{(P+\partial_t)^{-1}}(x, y, t - s)$ denote the kernel of $(P + \partial_t)^{-1}$. Then the heat kernel $k_t(x, y)$ of P satisfies*

$$(6.16) \quad k_t(x, y) = K_{(P+\partial_t)^{-1}}(x, y, t) \quad \text{for } t > 0.$$

Combining this with Proposition 6.11 then gives the heat kernel asymptotics for P in the form below.

Proposition 6.22 ([BGS, Thm. 5.6]). *If the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol, then as $t \rightarrow 0^+$ the following asymptotics holds in $C^\infty(M, (\text{End } \mathcal{E}) \otimes |\Lambda|(M))$,*

$$(6.17) \quad k_t(x, x) \sim t^{-\frac{d+2}{v}} \sum t^{\frac{2j}{v}} a_j(P)(x), \quad a_j(P)(x) = |\varepsilon'_x|(q_{-v-2j})_{(\xi, \tau) \rightarrow (y, t)}^\vee(x, 0, 1),$$

where the equality on the right shows how to compute $a_j(P)(x)$ in a local trivializing Heisenberg chart by means of the symbol $q_{-v-2j}(x, \xi, \tau)$ of degree $-v - 2j$ of any parametrix of $P + \partial_t$ in $\Psi_{H,v}^{-v}(M \times \mathbb{R}_{(v)}, \mathcal{E})$.

We will give in Section 8 criterions for the invertibility principal symbol of $P + \partial_t$ to be invertible. Nevertheless, in the case of a sublaplacian we have:

Proposition 6.23 ([BGS, Thm. 5.22]). *Let $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint sublaplacian which is bounded from below and assume that the condition (4.57) is satisfied at every point of M . Then the principal symbol of $\Delta + \partial_t$ is an invertible Volterra-Heisenberg symbol, hence Proposition 6.21 and Proposition 6.22 hold for Δ .*

Remark 6.24. As shown by Proposition 4.38 the condition (4.57) at a point of M is equivalent to the Rockland condition for Δ when the Levi form is degenerate, but is a stronger condition when the Levi form is nondegenerate. It will be shown in Section 8 that when the Levi form is everywhere nondegenerate the Rockland condition for Δ is enough to insure us the invertibility of the principal symbol of $\Delta + \partial_t$ even when the Levi form is nondegenerate at a point. In addition, it will be also shown in [Po10] that any selfadjoint sublaplacian with an invertible principal symbol is bounded from below, so the assumption on the boundedness from below of Δ is in fact superfluous.

Example 6.25. The above result is true for the following sublaplacians:

- (a) A selfadjoint sum of squares $\Delta = X_1 X_1^* + \dots + X_m X_m^*$ where $X_1, \dots, X_d$ span H ;
- (b) The Kohn Laplacian on a CR manifold and acting on (p, q) -forms under condition $Y(q)$;
- (c) The horizontal sublaplacian on a Heisenberg manifold acting on horizontal forms of degree k under condition $X(k)$;
- (d) The conformal powers of the horizontal sublaplacian acting on functions on a strictly pseudoconvex CR manifold.

6.2. Complex powers. Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of even (Heisenberg) order v such that P has an invertible principal symbol and is positive, i.e., we have $\langle Pu, u \rangle \geq 0$ for any $u \in C^\infty(M, \mathcal{E})$. Let $\Pi_0(P)$ be the orthogonal projection onto $\ker P$. Then the operator $P_0 := (1 - \Pi_0(P))P + \Pi_0(P)$ is selfadjoint with spectrum contained in $[c, \infty)$ for some $c > 0$. Thus by standard functional calculus, for any $s \in \mathbb{C}$, the power P_0^s is a well defined unbounded operator on $L^2(M, \mathcal{E})$. We then define the power P^s , $s \in \mathbb{C}$, by letting

$$(6.18) \quad P^s = (1 - \Pi_0(P))P_0^s = P_0^s - \Pi_0(P),$$

so that P^s coincides with P_0^s on $(\ker P)^\perp$ and is zero on $\ker P$. In particular, we have $P^0 = 1 - \Pi_0(P)$ and P^{-1} is the partial inverse of P .

The key result of this section is the following.

Theorem 6.26. *Suppose that the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol. Then:*

- (i) *For any $s \in \mathbb{C}$ the operator P^s defined by (6.18) is a Ψ_H DO of order vs ;*
- (ii) *The family $(P^s)_{s \in \mathbb{C}}$ forms a holomorphic 1-parameter group of Ψ_H DO's.*

Proof. Let us first assume that $\mathcal{E}$ is the trivial line bundle over M , so that P is a scalar operator. For $\Re s > 0$ the function $x \rightarrow x^{-s}$ is bounded on $[0, \infty)$, so the operators P_0^{-s} and P^{-s} are bounded. Moreover, by the Mellin formula we have

$$(6.19) \quad P^{-s} = (1 - \Pi_0(P))P_0^s = \frac{1}{\Gamma(s)} \int_0^\infty t^s (1 - \Pi_0(P)) e^{-tP} \frac{dt}{t}.$$

This leads us to define

$$(6.20) \quad A_s = \int_0^1 t^{s-1} e^{-tP} dt, \quad \Re s > 0.$$

Then we have

$$(6.21) \quad \begin{aligned} \Gamma(s)P^{-s} - A_s &= \int_0^1 t^{s-1} \Pi_0(P) e^{-tP} dt + \int_1^\infty t^{s-1} (1 - \Pi_0(P)) e^{-tP} dt, \\ &= \frac{1}{2} \Pi_0(P) + e^{-P/2} \left(\int_0^\infty (1+t)^{s-1} e^{-tP} dt \right) e^{-P/2}. \end{aligned}$$

Since $\Pi_0(P)$ and $e^{-P/2}$ are smoothing operators and $(\int_0^\infty (1+t)^{s-1} e^{-tP} dt)_{\Re s > 0}$ is a holomorphic family of bounded operators on $L^2(M)$, we get

$$(6.22) \quad (\Gamma(s)P^{-s} - A_s)_{\Re s > 0} \in \text{Hol}(\Re s > 0, \Psi^{-\infty}(M)).$$

Let us now show that $(A_s)_{\Re s > 0}$ defined by (6.20) is a holomorphic family of Ψ_H DO's such that $\text{ord} A_s = -vs$. To this end observe that, in terms of distribution kernels, the formula (6.20) means that A_s has distribution kernel

$$(6.23) \quad k_{A_s}(x, y) = \int_0^1 t^{s-1} k_t(x, y) dt.$$

where $k_t(x, y)$ denotes the heat kernel of P .

On the other hand, since P is bounded from below and the principal symbol of $P + \partial_t$ is invertible, Theorem 6.21 tells us that $P + \partial_t$ has an inverse $Q_0 := (P + \partial_t)^{-1}$ in $\Psi_{H,v}^{-v}(M \times \mathbb{R}_{(v)}, \mathcal{E})$ and that the distribution kernel $K_{Q_0}(x, y, t - s)$ of Q_0 is related to the heat kernel of P by

$$(6.24) \quad K_{Q_0}(x, y, t) = k_t(x, y) \quad \text{for } t > 0.$$

Therefore, for $\Re s > 0$ we have

$$(6.25) \quad k_{A_s}(x, y) = \int_0^1 t^{s-1} K_{Q_0}(x, y, t) dt.$$

Let φ and ψ be smooth functions on M with disjoint supports. Then using (6.25) we see that $\varphi A_s \psi$ has distribution kernel

$$(6.26) \quad k_{\varphi A_s \psi}(x, y) = \int_0^1 t^{s-1} \varphi(x) K_{Q_0}(x, y, t) \psi(y) dt.$$

Since the distribution kernel of a Volterra- Ψ_H DO is smooth off the diagonal of $(M \times \mathbb{R}) \times (M \times \mathbb{R})$ the distribution $K_{Q_0}(x, y, t)$ is smooth on the region $\{x \neq y\} \times \mathbb{R}$, so (6.26) defines a holomorphic family of smooth kernels. Thus,

$$(6.27) \quad (\varphi A_s \psi)_{\Re s > 0} \in \text{Hol}(\Re s > 0, \Psi^{-\infty}(M)).$$

Next, the following holds.

Lemma 6.27. *Let $V \subset \mathbb{R}^{d+1}$ be a Heisenberg chart, let $Q \in \Psi_{H,v}^{-v}(V \times \mathbb{R}_{(v)})$ have distribution kernel $K_Q(x, y, t-s)$ and for $\Re s > 0$ let $B_s : C_c(V) \rightarrow C(V)$ be given by the distribution kernel,*

$$(6.28) \quad k_{B_s}(x, y) = \int_0^1 t^{s-1} K_Q(x, y, t) dt, \quad \Re s > 0.$$

Then $(B_s)_{\Re s > 0}$ is a holomorphic family of Ψ_H DO's such that $\text{ord} B_s = -vs$

Proof of the lemma. Let ε_x denote the change to the Heisenberg coordinates at x . By Proposition 6.10 on $V \times V \times \mathbb{R}$ the distribution $K_Q(x, y, t)$ is of the form

$$(6.29) \quad K_Q(x, y, t) = |\epsilon'_x| K(x, -\varepsilon_x(y), t) + R(x, y, t),$$

for some $K \in \mathcal{K}_v^{-(d+2)}(V \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$ and some $R \in C^\infty(U \times U \times \mathbb{R})$. Let us write $K \sim \sum_{j \geq 0} K_{j-(d+2)}$ with $K_l \in \mathcal{K}_{v,l}(V \times \mathbb{R}^{d+1} \times \mathbb{R}_{(v)})$. Thus, for any integer N , as soon as J large enough we have

$$(6.30) \quad K(x, y, t) = \sum_{j \leq J} K_{j-(d+2)}(x, y, t) + R_{NJ}(x, y, t), \quad R_{NJ} \in C^N(U \times \mathbb{R}^{d+2}).$$

In particular, on $V \times V$ we have

$$(6.31) \quad k_{B_s}(x, y) = |\epsilon'_x| K_s(x, \varepsilon_x(y)) + R_s(x, y), \quad K_s(x, y) = \sum_{j \leq J} K_{j,s}(x, y) + R_{NJ,s}(x, y),$$

where we have let

$$(6.32) \quad K_s(x, y) = \int_0^1 t^{s-1} K(x, y, t) dt, \quad K_{j,s}(x, y) = \int_0^1 t^{s-1} K_{j-(d+2)}(x, y, t) dt, \quad j \geq 0,$$

and $(R_s)_{\Re s > 0}$ and $(R_{NJ,s})_{\Re s > 0}$ are in $\text{Hol}(\Re s > 0, C^\infty(V \times V))$ and $\text{Hol}(\Re s > 0, C^N(V \times V))$ respectively.

Notice that $K_{j-(d+2)}(x, y, t)$ is in $C^\infty(V) \hat{\otimes} \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1} \times \mathbb{R})$ and is parabolic homogeneous of degree $j - (d+2) \geq -(d+2)$. Thus the family $(K_{j,s})_{\Re s > 0}$ belongs to $\text{Hol}(\Re s > 0, C^\infty(U) \hat{\otimes} \mathcal{D}'_{\text{reg}}(\mathbb{R}^{d+1}))$. Moreover, for any $\lambda > 0$, the difference $K_{j,s}(x, \lambda y) - \lambda^{vs+j-(d+2)} K_{j,s}(x, y)$ is equal to

$$(6.33) \quad \int_1^{\lambda^2} t^{s-1} K(x, y, t) dt \in \text{Hol}(\Re s > 0, C^\infty(V \times \mathbb{R}^{d+2})).$$

Hence $(K_{j,s})_{\Re s > 0}$ is a holomorphic family of almost homogeneous distributions of degree $vs - (d+2) + j$. Combining this with (6.31) then shows that $(K_s)_{\Re s > 0}$ belongs to $\text{Hol}(\Re s > 0, \mathcal{K}^*(V \times \mathbb{R}^{d+1}))$ and has order $vs - (d+2)$. Therefore, using (6.31) and Proposition 5.26 we see that $(B_s)_{\Re s > 0}$ is a holomorphic family of Ψ_H DO's such that $\text{ord} B_s = -(\text{ord} K_s + d+2) = -vs$. $\square$

It follows from Lemma 6.27 that for any local Heisenberg chart $\kappa : U \rightarrow V$ the family $(\kappa_* A_{s|U})_{\Re s > 0}$ is a holomorphic family of Ψ_H DO's on V of order $-vs$. Combining this with (6.27) and (6.22) then shows that $(A_s)_{\Re s > 0}$ and $(P^s)_{\Re s < 0}$ are holomorphic families of Ψ_H DO's of orders $-vs$ and vs respectively.

Now, let $s \in \mathbb{C}$ and let k be a positive integer such that $k > \Re s$. Then we have

$$(6.34) \quad P^s u = P^{s-k} P^k u \quad \text{for any } u \in C^\infty(M, \mathcal{E}).$$

As P^k is a differential operator and P^{s-k} is a Ψ_H DO of order $m(s-k)$ this proves that P^s is a Ψ_H DO of order ms . In fact, as by Proposition 5.19 the product of Ψ_H DO's is analytic this actually shows that $(P^s)_{s \in \mathbb{C}}$ is a holomorphic family of Ψ_H DO's such that $\text{ord} P^s = vs$ for every $s \in \mathbb{C}$.

Finally, when $\mathcal{E}$ is a general vector bundle we can similarly prove that the complex powers P^s , $s \in \mathbb{C}$, forms a holomorphic family of Ψ_H DO's such that $\text{ord} P^s = vs$ for any $s \in \mathbb{C}$. The proof is thus complete. $\square$

Example 6.28. Theorem 6.26 holds for the following sublaplacians:

- (a) A selfadjoint sums of squares $\Delta = \nabla_{X_1}^* \nabla_{X_1} + \dots + \nabla_{X_m}^* \nabla_{X_m}$, where $X_1, \dots, X_m$ span H and ∇ is a connection on $\mathcal{E}$, under the condition that the Levi form is nonvanishing.
- (b) The Kohn Laplacian $\square_b$ on a compact CR manifold acting on (p, q) -forms when the condition $Y(q)$ holds everywhere.
- (c) The horizontal sublaplacian Δ_b on a compact Heisenberg manifold (M^{d+1}, H) acting on sections of $\Lambda_{\mathbb{C}}^k H^*$ when the condition $X(k)$ holds everywhere.

In Section 8 we will actually make use of Theorem 6.26 to show that when the Levi form has constant rank the Rockland condition is enough to insure us that the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol (see Theorem 8.5). Therefore, we obtain:

Theorem 6.29. *Assume that the Levi form of (M, H) has constant rank and let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a positive differential operator of even (Heisenberg) order v such that P satisfies the Rockland condition at every point of M . Then:*

- (i) *For any $s \in \mathbb{C}$ the operator P^s defined by (6.18) is a Ψ_H DO of order vs ;*
- (ii) *The family $(P^s)_{s \in \mathbb{C}}$ forms a holomorphic 1-parameter group of Ψ_H DO's.*

Example 6.30. Theorem 6.29 is valid for the contact Laplacian Δ_R on a compact orientable contact manifold (M^{2n+1}, θ) . In this case Δ_R^s has order $2s$ on degree $k = 0, \dots, 2n$ with $k \neq n$ and has order $4s$ on degree n .

Remark 6.31. The above example allows us to fill a technical gap in the proof in [JK] of the Baum-Connes conjecture for $SU(n, 1)$ (see [Po9]).

7. WEIGHTED SOBOLEV SPACES

Let (M^{d+1}, H) be a compact Heisenberg manifold endowed with a smooth positive density and assume that the Levi form of (M, H) is non-vanishing. We shall now construct weighted Sobolev spaces $W_H^s(M)$, $s \in \mathbb{R}$, which extend to any real parameter the weighted Sobolev spaces $S_k^2(M)$, $k \in \mathbb{N}$, of Folland-Stein [FS1] (see also [RSt]). As a consequence these Sobolev spaces will provide us with sharp regularity estimates for hypoelliptic Ψ_H DO's.

Let $X_1, \dots, X_m$ be real vector fields spanning H and consider the positive sum of squares,

$$(7.1) \quad \Delta_X = X_1^* X_1 + \dots + X_m^* X_m.$$

As mentioned in Example 6.28 (a), since the Levi form of (M, H) is non-vanishing Theorem 6.26 is valid for $1 + \Delta_X$. Thus, the complex powers $(1 + \Delta_X)^s$, $s \in \mathbb{C}$, gives rise to a holomorphic 1-parameter group of Ψ_H DO's such that $\text{ord}(1 + \Delta_X)^s = 2s$ for any $s \in \mathbb{C}$.

Definition 7.1. $W_H^s(M)$, $s \in \mathbb{R}$, consists of all distributions $u \in \mathcal{D}'(M)$ such that $(1 + \Delta_X)^{\frac{s}{2}}u$ is in $L^2(M)$. It is endowed with the Hilbert norm given by

$$(7.2) \quad \|u\|_{W_H^s} = \|(1 + \Delta_X)^{\frac{s}{2}}u\|_{L^2}, \quad u \in W_H^s(M).$$

Proposition 7.2. 1) Neither $W_H^s(M)$, nor its topology, depend on the choice of the vector fields $X_1, \dots, X_m$.

2) We have the following continuous embeddings:

$$(7.3) \quad \begin{array}{llll} L_s^2(M) & \hookrightarrow & W_H^s(M) & \hookrightarrow & L_{s/2}^2(M) & \text{if } s \geq 0, \\ L_{s/2}^2(M) & \hookrightarrow & W_H^s(M) & \hookrightarrow & L_s^2(M) & \text{if } s < 0. \end{array}$$

Proof. 1) Let $Y_1, \dots, Y_p$ be other vector fields spanning H . The operator $(1 + \Delta_Y)^s(1 + \Delta_X)^{-s}$ is a Ψ_H DO of order 0, so is bounded on $L^2(M)$ by Proposition 4.8. Therefore, we get the estimates,

$$(7.4) \quad \|(1 + \Delta_Y)^s u\|_{L^2} = \|(1 + \Delta_Y)^s(1 + \Delta_X)^{-s}(1 + \Delta_X)^s u\|_{L^2} \leq C_{XYs} \|(1 + \Delta_X)^s u\|_{L^2},$$

which hold for any $u \in C^\infty(M)$. Interchanging the roles of the X_j 's and of the Y_k 's also gives the estimates

$$(7.5) \quad \|(1 + \Delta_X)^s u\|_{L^2} \leq C_{YXs} \|(1 + \Delta_Y)^s u\|_{L^2}, \quad u \in C^\infty(M).$$

Therefore, whether we use the X_j 's or the Y_k 's to define the $W_H^s(M)$ changes neither the space, nor its topology.

2) Let $s \in [0, \infty)$. Since $(1 + \Delta_X)^{\frac{s}{2}}$ is a Ψ_H DO of order s , Proposition 4.8 tells us that it is bounded from $L_s^2(M)$ to $L^2(M)$. Thus,

$$(7.6) \quad \|u\|_{W_H^s} = \|(1 + \Delta_X)^{\frac{s}{2}}u\|_{L^2} \leq C_s \|u\|_{L_s^2}, \quad u \in W_H^s(M),$$

which shows that $L_s^2(M)$ embeds continuously into $W_H^s(M)$.

On the other hand, as $(1 + \Delta_X)^{-\frac{s}{2}}$ has order $-s$ Proposition 4.8 also tells us that $(1 + \Delta_X)^{-\frac{s}{2}}$ is bounded from $L^2(M)$ to $L_{s/2}^2(M)$. Therefore, for we get the estimates

$$(7.7) \quad \|u\|_{L_{s/2}^2} = \|(1 + \Delta_X)^{-\frac{s}{2}}(1 + \Delta_X)^{\frac{s}{2}}u\|_{L_{s/2}^2} \leq C_s \|(1 + \Delta_X)^{\frac{s}{2}}u\|_{L^2} = C_s \|u\|_{W_H^s},$$

which hold for any $u \in W_H^s(M)$ and show that $W_H^s(M)$ embeds continuously into $L_{s/2}^2(M)$.

Finally, when $s < 0$ we can similarly show that we have continuous embeddings $L_{s/2}^2(M) \hookrightarrow W_H^s(M)$ and $W_H^s(M) \hookrightarrow L_s^2(M)$. $\square$

As an immediate consequence of Proposition 7.2 we obtain:

Proposition 7.3. The following equalities between topological spaces hold:

$$(7.8) \quad C^\infty(M) = \cap_{s \in \mathbb{R}} W_H^s(M) \quad \text{and} \quad \mathcal{D}'(M) = \cup_{s \in \mathbb{R}} W_H^s(M).$$

Let us now compare the Weighted Sobolev spaces $W_H^s(M)$ to the weighted Sobolev spaces $S_k^2(M)$, $k = 1, 2, \dots$, of Folland-Stein [FS1].

In we sequel we let $\mathbb{N}_m = \{1, \dots, m\}$ and for any $I = (i_1, \dots, i_k)$ in $\mathbb{N}_m^k$ we set

$$(7.9) \quad X_I = X_{i_1} \dots X_{i_k}.$$

Definition 7.4 ([FS1]). The Hilbert space $S_k^2(M)$, $k \in \mathbb{N}$, consists of functions $u \in L^2(M)$ such that $(X_I)u \in L^2(M)$ for any $I \in \cup_{j=1}^k \mathbb{N}_m^j$. It is endowed with the Hilbertian norm given by

$$(7.10) \quad \|u\|_{S_k^2}^2 = \|u\|_{L^2}^2 + \sum_{1 \leq j \leq k} \sum_{I \in \mathbb{N}_m^j} \|X_I u\|_{L^2}^2, \quad u \in S_k^2(M).$$

Proposition 7.5. For $k = 1, 2, \dots$ the weighted Sobolev spaces $W_H^k(M)$ and $S_k^2(M)$ agree as spaces and bear the same topology.

Proof. First, if $I \in \cup_{j=1}^k \mathbb{N}_m^j$ then by Proposition 7.8 the differential operator X_I is bounded from $W_H^k(M)$ to $L^2(M)$, so we get the estimate,

$$(7.11) \quad \|u\|_{S^k}^2 = \|u\|_{L^2}^2 + \sum_{1 \leq j \leq k} \sum_{I \in \mathbb{N}_m^j} \|X_I u\|_{L^2}^2 \leq C_k^2 \|u\|_{W_H^k}^2 \quad u \in C^\infty(M).$$

On the other hand, for any $l \in \mathbb{N}$ the differential operators X_j and X_j^* , $j = 1, \dots, m$, are bounded linear maps from $S_{l+1}^2(M)$ to $S_l^2(M)$. Thus, for any integer p , the operator $(1 + \Delta_X)^p$ is bounded from $S_{l+2p}^2(M)$ to $S_l^2(M)$. It follows that, when k is even, we have

$$(7.12) \quad \|u\|_{W_H^k} = \|(1 + \Delta_X)^{\frac{k}{2}} u\|_{L^2} \leq C_k \|u\|_{S^k}, \quad u \in C^\infty(M).$$

Moreover, for any $u \in C^\infty(M)$ we have

$$(7.13) \quad \|(1 + \Delta_X)^{\frac{1}{2}} u\|_{L^2}^2 = \langle (1 + \sum_{1 \leq j \leq m} X_j^* X_j) u, u \rangle_{L^2} = \|u\|_{L^2}^2 + \sum_{1 \leq j \leq m} \|X_j^2 u\|_{L^2}^2 = \|u\|_{S_1^2}^2.$$

Thus $(1 + \Delta_X)^{\frac{1}{2}}$ is bounded from $S_1^2(M)$ to $L^2(M)$. Therefore, if k is odd, say $k = 2p + 1$, then the operator $(1 + \Delta_X)^{\frac{k}{2}} = (1 + \Delta_X)^p (1 + \Delta_X)^{\frac{1}{2}}$ is bounded from $S_k^2(M)$ to $L^2(M)$. Hence (7.13) is valid in the odd case as well. Together with (7.11) this implies that $W_H^k(M)$ and $S_k^2(M)$ agree as spaces and bear the same topology. $\square$

Now, let $\mathcal{E}$ be a Hermitian vector bundle over M . We can also define weighted Sobolev spaces of sections of $\mathcal{E}$ as follows. Let $\nabla : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, T^*M \times \mathcal{E})$ be a connection on $\mathcal{E}$ and define

$$(7.14) \quad \Delta_{\nabla, X} = \nabla_{X_1}^* \nabla_{X_1} + \dots + \nabla_{X_m}^* \nabla_{X_m}.$$

As in the scalar case, the complex powers $(1 + \Delta_{\nabla, X})^s$, $s \in \mathbb{C}$, form an analytic 1-parameter group of Ψ_H DO's such that $\text{ord}(1 + \Delta_{\nabla, X})^s = 2s$ for any $s \in \mathbb{C}$.

Definition 7.6. $W_H^s(M, \mathcal{E})$, $s \in \mathbb{R}$, consists of all distributional sections $u \in \mathcal{D}'(M, \mathcal{E})$ such that $(1 + \Delta_{\nabla, X})^{\frac{s}{2}} u \in L^2(M, \mathcal{E})$. It is endowed with the Hilbertian norm given by

$$(7.15) \quad \|u\|_{W_H^s} = \|(1 + \Delta_{\nabla, X})^{\frac{s}{2}} u\|_{L^2}, \quad u \in W_H^s(M, \mathcal{E}).$$

Along similar lines as that of the proof of Proposition 7.2 we can prove:

Proposition 7.7. 1) As a topological space $W_H^s(M, \mathcal{E})$, $s \in \mathbb{R}$, is independent of the choice of the connection ∇ and of the vector fields $X_1, \dots, X_m$.

2) We have the following continuous embeddings:

$$(7.16) \quad \begin{array}{llll} L_s^2(M, \mathcal{E}) & \hookrightarrow & W_H^s(M, \mathcal{E}) & \hookrightarrow & L_{s/2}^2(M, \mathcal{E}) & \text{if } s \geq 0, \\ L_{s/2}^2(M, \mathcal{E}) & \hookrightarrow & W_H^s(M, \mathcal{E}) & \hookrightarrow & L_s^2(M, \mathcal{E}) & \text{if } s < 0. \end{array}$$

As a consequence we obtain the equalities of topological spaces,

$$(7.17) \quad C^\infty(M, \mathcal{E}) = \cap_{s \in \mathbb{R}} W_H^s(M, \mathcal{E}) \quad \text{and} \quad \mathcal{D}'(M, \mathcal{E}) = \cup_{s \in \mathbb{R}} W_H^s(M, \mathcal{E}).$$

Notice that we can also define Folland-Stein spaces $S_k^2(M, \mathcal{E})$, $k = 1, 2, \dots$, as in the scalar case, by using the differential operators $\nabla_{X_I} = \nabla_{X_{i_1}} \dots \nabla_{X_{i_k}}$, $I \in \cup_{j=1}^k \mathbb{N}_m^j$. Then, by arguing as in the proof of Proposition 7.5, we can show that the spaces $W_H^k(M, \mathcal{E})$ and $S_k^2(M, \mathcal{E})$ agree and bear the same topology.

Now, the Sobolev spaces $W_H^s(M, \mathcal{E})$ yield sharp regularity results for Ψ_H DO's.

Proposition 7.8. Let $P \in \Psi_H^m(M, \mathcal{E})$, set $k = \Re m$ and let $s \in \mathbb{R}$.

1) The operator P extends to a continuous linear mapping from $W_H^{s+k}(M, \mathcal{E})$ to $W_H^s(M, \mathcal{E})$.

2) Assume that the principal symbol of P is invertible. Then for any $u \in \mathcal{D}'(M, \mathcal{E})$ we have

$$(7.18) \quad Pu \in W_H^s(M, \mathcal{E}) \implies u \in W_H^{s+k}(M, \mathcal{E}).$$

In fact, for any $s' \in \mathbb{R}$ we have the estimate,

$$(7.19) \quad \|u\|_{W_H^{s+k}} \leq C_{ss'}(\|Pu\|_{W_H^s} + \|u\|_{W_H^{s'}}), \quad u \in W_H^{s+k}(M, \mathcal{E}).$$

Proof. 1) As $P_s = (1 + \Delta_{\nabla, X})^{\frac{s}{2}} P (1 + \Delta_{\nabla, X})^{-\frac{(s+k)}{2}}$ is a Ψ_H DO with purely imaginary order, Proposition 4.8 tells us that it gives rise to a bounded operator on $L^2(M, \mathcal{E})$. Therefore, we have

$$(7.20) \quad \|Pu\|_{W_H^s} = \|P_s(1 + \Delta_{\nabla, X})^{s+k}u\|_{L^2} \leq C_s\|u\|_{W_H^{s+k}}, \quad u \in C^\infty(M, \mathcal{E}),$$

It then follows that P extends to a continuous linear mapping from $W_H^{s+k}(M, \mathcal{E})$ to $W_H^s(M, \mathcal{E})$.

2) Since the principal symbol of P is invertible by Proposition 4.32 there exist Q in $\Psi_H^{-m}(M, \mathcal{E})$ and R in $\Psi^{-\infty}(M, \mathcal{E})$ such that $QP = 1 - R$. Therefore, for any $u \in \mathcal{D}'(M, \mathcal{E})$ we have $u = QPu + Ru$. Thanks to the first part we know that Q maps $W_H^s(M, \mathcal{E})$ to $W_H^{s+k}(M, \mathcal{E})$. Since R is smoothing, and so Ru always is smooth, it follows that if Pu is in $W_H^s(M, \mathcal{E})$ then u must be in $W_H^{s+k}(M, \mathcal{E})$.

In fact, as Q is actually bounded from W_H^s to W_H^{s+k} and (7.17) implies that R is bounded from any space $W_H^{s'}(M, \mathcal{E})$ to $W_H^{s+k}(M, \mathcal{E})$ we have estimates,

$$(7.21) \quad \|u\|_{W_H^{s+k}} \leq \|QPu\|_{W_H^{s+k}} + \|Ru\|_{W_H^{s+k}} \leq C_{ss'}(\|Pu\|_{W_H^s} + \|u\|_{W_H^{s'}}), \quad u \in W_H^{s+k}(M, \mathcal{E}).$$

The proof is thus complete. $\square$

Remark 7.9. Combining Proposition 7.8 and the embeddings (7.3) and (7.16) we recover the Sobolev regularity of Ψ_H DO's as in Proposition 4.8 as well as the hypoelliptic estimates (4.49).

On the other hand, the spaces $W_H^s(M, \mathcal{E})$ can be localized as follows.

Definition 7.10. We say that $u \in \mathcal{D}'(M, \mathcal{E})$ is W_H^s near a point $x_0 \in M$ whenever there exists $\varphi \in C^\infty(M)$ such that $\varphi(x_0) \neq 0$ and φu is in $W_H^s(M, \mathcal{E})$.

This definition depends only on the germ of u at x_0 because we have:

Lemma 7.11. Let $u \in \mathcal{D}'(M, \mathcal{E})$ be W_H^s near x_0 . Then for any $\varphi \in C^\infty(M)$ with a small enough support about x_0 the distribution φu lies in $W_H^s(M, \mathcal{E})$.

Proof. Let $\varphi \in C^\infty(M)$ be such that $\varphi(x_0) \neq 0$ and φu is in $W_H^s(M, \mathcal{E})$ and let $\psi \in C^\infty(M)$ be so that $\psi(x_0) \neq 0$ and $\text{supp } \psi \cap \varphi^{-1}(0) \neq \emptyset$. Then $\chi := \frac{\psi}{\varphi}$ is in $C^\infty(M)$ and we have

$$(7.22) \quad (1 + \Delta_{\nabla, X})^{\frac{s}{2}} \psi u = (1 + \Delta_{\nabla, X})^{\frac{s}{2}} \chi \varphi u = (1 + \Delta_{\nabla, X})^{\frac{s}{2}} \chi (1 + \Delta_{\nabla, X})^{-\frac{s}{2}} (1 + \Delta_{\nabla, X})^{\frac{s}{2}} \varphi u.$$

Since $(1 + \Delta_{\nabla, X})^{\frac{s}{2}} \varphi u$ is in $L^2(M, \mathcal{E})$ and $(1 + \Delta_{\nabla, X})^{\frac{s}{2}} \chi (1 + \Delta_{\nabla, X})^{-\frac{s}{2}}$ is a zero'th order Ψ_H DO, so maps $L^2(M, \mathcal{E})$ to itself, it follows that ψu lies in $W_H^s(M, \mathcal{E})$. Hence the lemma. $\square$

We can now get a localized version of (7.18).

Proposition 7.12. Let $P \in \Psi_H^m(M, \mathcal{E})$ have an invertible principal symbol, set $k = \Re m$ and let $s \in \mathbb{R}$. Then for any $u \in \mathcal{D}'(M, \mathcal{E})$ we have

$$(7.23) \quad Pu \text{ is } W_H^s \text{ near } x_0 \implies u \text{ is } W_H^s \text{ near } x_0.$$

Proof. Assume that Pu is W_H^s near x_0 and let $\varphi \in C^\infty(M)$ be such that $\varphi(x_0) \neq 0$ and φu is in $W_H^s(M, \mathcal{E})$. Thanks to Lemma 7.11 we may assume that $\varphi = 1$ near x_0 . Let $\psi \in C^\infty(M)$ be such that $\psi(x_0) \neq 0$ and $\varphi = 1$ near $\text{supp } \psi$. Since the principal symbol of P is invertible there exist Q in $\Psi_H^{-m}(M, \mathcal{E})$ and R in $\Psi^{-\infty}(M, \mathcal{E})$ such that $QP = 1 - R$. Thus for any $u \in \mathcal{D}'(M, \mathcal{E})$ we have

$$(7.24) \quad \psi u = \psi QPu + \psi Ru = \psi Q\varphi Pu + \psi Q(1 - \varphi)Pu + \psi Ru.$$

In the above equality Ru is a smooth function since R is a smoothing operator. Similarly, as ψ and $1 - \varphi$ have disjoint supports, the operator $\psi Q(1 - \varphi)P$ is a smoothing operator and so $\psi Q(1 - \varphi)Pu$ is a smooth function. In addition, since φPu is in $W_H^s(M, \mathcal{E})$ and ψQ is a Ψ_H DO of order $-m$, and so maps $W_H^s(M, \mathcal{E})$ to $W_H^{s+k}(M, \mathcal{E})$, it follows that ψu is in W_H^{s+k} . Hence u is W_H^{s+k} near x_0 . $\square$

Next, the first part of Proposition 7.8 admits generalization to holomorphic families of Ψ_H DO's.

Proposition 7.13. *Let $\Omega \subset \mathbb{C}$ be open and let $(P_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi_H^*(M, \mathcal{E}))$. Assume there exists a real m such that $\Re \text{ord} P_z \leq m < \infty$. Then for any $s \in \mathbb{R}$ the family $(P_z)_{z \in \Omega}$ defines a holomorphic family with values in $\mathcal{L}(W_H^{s+m}(M, \mathcal{E}), W_H^s(M, \mathcal{E}))$.*

Proof. First, let $V \subset \mathbb{R}^{d+1}$ be a Heisenberg chart with H -frame $Y_0, Y_1, \dots, Y_d$ and let $(Q_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi_H^*(V))$ be such that $\Re \text{ord} Q_z \leq 0$. Then we can write

$$(7.25) \quad Q_z = p_z(x, -iY) + R_z,$$

for some families $(p_z)_{z \in \Omega} \in \text{Hol}(\Omega, S^*(V \times \mathbb{R}^{d+1}))$ and $(R_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi^\infty(V))$. Since $\Re \text{ord} p_z = \Re \text{ord} Q_z \leq 0$ we see that $(p_z)_{z \in \Omega}$ belongs to $\text{Hol}(\Omega, S_{\parallel}^0(U \times \mathbb{R}^{d+1}))$.

Next, for $j = 1, \dots, d$ let σ_j denote the standard symbol of $-iY_j$ and set $\sigma = (\sigma_0, \dots, \sigma_d)$. Then it follows from the proof of Proposition 5.11 that the family $(p_z(x, \sigma(x, \xi)))_{z \in \Omega}$ lies in $\text{Hol}(\Omega, S_{\frac{1}{2}, \frac{1}{2}}(V \times \mathbb{R}^{d+1}))$. Since by [Hw] the quantization map $q \rightarrow q(x, D_x)$ is continuous from $S_{\frac{1}{2}, \frac{1}{2}}(U \times \mathbb{R}^{d+1})$ to $\mathcal{L}(L_{\text{loc}}^2(U), L^2(U))$, we deduce that $(p_z(x, -iX))_{z \in \Omega}$ and $(Q_z)_{z \in \Omega}$ are holomorphic families with values in $\mathcal{L}(L_{\text{loc}}^2(U), L^2(U))$.

It follows from the above result that any family $(Q_z)_{z \in \Omega} \in \text{Hol}(\Omega, \Psi_H^*(M, \mathcal{E}))$ such that $\Re \text{ord} Q_z \leq 0$ gives rise to a holomorphic family with values in $\mathcal{L}(L^2(M, \mathcal{E}))$. This applies in particular to the family

$$(7.26) \quad Q_z^{(s)} = (1 + \Delta_{\nabla, X})^{\frac{s}{2}} Q_z (1 + \Delta_{\nabla, X})^{-\frac{m+s}{2}}, \quad z \in \Omega.$$

As $Q_z = (1 + \Delta_{\nabla, X})^{-\frac{s}{2}} Q_z^{(s)} (1 + \Delta_{\nabla, X})^{\frac{m+s}{2}}$ it follows that $(Q_z)_{z \in \Omega}$ gives rise to a holomorphic family with values in $\mathcal{L}(W_H^{s+m}(M, \mathcal{E}), W_H^s(M, \mathcal{E}))$. $\square$

Combining Proposition 7.13 with Theorem 6.29 we immediately get:

Proposition 7.14. *Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a positive differential operator of even (Heisenberg) order w with an invertible principal symbol. Then, for any reals m and s , the family of the complex powers of P satisfies*

$$(7.27) \quad (P^z)_{\Re z < m} \in \text{Hol}(\Re z < m, \mathcal{L}(W_H^{s+mw}(M, \mathcal{E}), W_H^s(M, \mathcal{E}))).$$

8. ROCKLAND CONDITION AND THE HEAT EQUATION

Let (M^{d+1}, H) be a compact Heisenberg manifold equipped with a smooth positive density and let $\mathcal{E}$ be a Hermitian vector bundle over M . In this section we shall show that for a selfadjoint differential operator $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ which is bounded from below, the invertibility of the principal symbol of P is enough to insure us the invertibility of the principal symbol of $P + \partial_t$. This is needed for the pseudodifferential representation of the heat kernel and the invertibility of the heat operator $P + \partial_t$ in [BGS], as well as in Theorem 6.26 for the complex powers of P to give rise to a holomorphic family of Ψ_H DO's.

Let us first look at symbols that are positive in the sense below.

Definition 8.1. *A symbol $p \in S_m(\mathfrak{g}^*M, \mathcal{E})$ is said to be positive when it can be put into the form $p = \bar{q} * q$ for some symbol $q \in S_{\frac{m}{2}}(\mathfrak{g}^*M, \mathcal{E})$.*

Lemma 8.2. *Let $P \in \Psi_H^m(M, \mathcal{E})$ have principal symbol $\sigma_m(P) \in S^m(\mathfrak{g}^*M, \mathcal{E})$. Then we have equivalence:*

- (i) *The symbol $\sigma_m(P)$ is positive;*
- (ii) *There exist $Q \in \Psi_H^{\frac{m}{2}}(M, \mathcal{E})$ and $R \in \Psi_H^{m-1}(M, \mathcal{E})$ so that $P = Q^*Q + R$.*

Proof. Assume that $\sigma_m(P)$ is a positive symbol, so that there exists $q_{\frac{m}{2}} \in S_{\frac{m}{2}}(\mathfrak{g}^*M, \mathcal{E})$ such that $p = \overline{q_{\frac{m}{2}}} * q_{\frac{m}{2}}$. By Proposition 4.26 the principal symbol map $\sigma_{\frac{m}{2}} : \Psi_H^{\frac{m}{2}}(M, \mathcal{E}) \rightarrow S^{\frac{m}{2}}(\mathfrak{g}^*M, \mathcal{E})$ is surjective, so there exists $Q \in \Psi_H^{\frac{m}{2}}(M, \mathcal{E})$ such that $\sigma_{\frac{m}{2}}(Q) = q_{\frac{m}{2}}$. Then by Proposition 4.29 and Proposition 4.31 the operator Q^*Q has principal symbol $\overline{q_{\frac{m}{2}}} * q_{\frac{m}{2}} = \sigma_m(P)$, hence coincides with P modulo $\Psi_H^{m-1}(M, \mathcal{E})$.

Conversely, if P is of the form $P = Q^*Q + R$ with $Q \in \Psi_H^{\frac{m}{2}}(M, \mathcal{E})$ and $R \in \Psi_H^{m-1}(M, \mathcal{E})$ then P and Q^*Q have same principal symbol. Then $\sigma_m(P) = \overline{\sigma_{\frac{m}{2}}(Q)} * \sigma_{\frac{m}{2}}(Q)$, that is, $\sigma_m(P)$ is a positive symbol. $\square$

From now on we assume that the Levi form of (M, H) has constant rank and we let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a differential operator of even Heisenberg order m .

Lemma 8.3. *Assume that P satisfies the Rockland condition at every point and has a positive principal symbol. Then the principal symbol of the heat operator $P + \partial_t$ is an invertible Volterra-Heisenberg symbol.*

Proof. Let us first assume that $\mathcal{E}$ is the trivial line bundle. Since proving the invertibility of the principal symbol of $P + \partial_t$ is a local problem with respect to the space covariable, we may as well work in a local Heisenberg chart.

Moreover, as the Levi form has constant rank it follows from [Po6] that GM is a fiber bundle of Lie group with fiber $G = \mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$, where $2n$ is the rank of the Levi form. Therefore, by considering a trivialization of this fiber bundle by means of a suitable local H -frame (see [Po6]) we may further assume that GU is the trivial bundle $U \times G$. In particular, the families of model operators $(P^x)_{x \in U}$ can be seen as a smooth family of self-adjoint convolution operators on $S_0(G)$.

Next, as the global principal symbol $\sigma_m(P)$ of P is positive there exists $\tilde{p}_{\frac{m}{2}} \in S_{\frac{m}{2}}(U \times G)$ such that $\sigma_m(P) = \overline{\tilde{p}_{\frac{m}{2}}} * \tilde{p}_{\frac{m}{2}}$, so if we let $\tilde{P}^x$ be the left-convolution operator with symbol $\tilde{p}_{\frac{m}{2}}(x, \cdot)$ then by Proposition 4.29 and Proposition 4.31 we have $P^x = (\tilde{P}^x)^* \tilde{P}^x$. Thus, by Proposition 4.34 for every non-trivial irreducible representation π of G we have $\overline{\pi P^x} = (\overline{\pi \tilde{P}^x})^* \pi \tilde{P}^x$, which shows that $\overline{\pi P^x}$ is positive.

Now, as P^x is positive and satisfies the Rockland condition it then follows from [FS2] that:

- The operator $P^x + \partial_t$ satisfies the Rockland condition on $G \times \mathbb{R}$ (see [FS2, Lem. 4.21]);
- There exists $K^x(y, t) \in C^\infty((G \times \mathbb{R}) \setminus \{0\})$ homogeneous of degree $-(d+2)$ such that $K^x(y, t) = 0$ for $t < 0$ and $K^x(y, t)$ is a fundamental solution of $P^x + \partial_t$, that is,

$$(8.1) \quad (P^x + \partial_t)K^x(y, t) = \delta(y, t),$$

where $\delta(y, t)$ denotes the Dirac distribution at the origin on $G \times \mathbb{R}$ (see [FS2, Lem. 4.24] in the case $m > \frac{d+2}{2}$ and [FS2, pp. 136–137] for the general case).

Let Q^x be the left-convolution operator on $G \times \mathbb{R}$ with $K^x(y, t)$. As K^x belongs to $\mathcal{K}_{v, -(d+2)}(G \times \mathbb{R}_{(m)})$ this is a left-invariant Volterra Ψ_H DO with symbol $q_{-m}^x(\xi, \tau) = (K^x)^\wedge(\xi, \tau)$. Moreover, thanks to (8.1) for any $f \in \mathcal{S}(G \times \mathbb{R})$ we have

$$(8.2) \quad (P^x + \partial_t)Q^x f = (P^x + \partial_t)(K * f) = [(P^x + \partial_t)K] * f = \delta * f = f.$$

Hence $(P^x + \partial_t)Q^x = 1$, which gives $(\sigma_m(P)(x, \cdot) + i\tau) * q_{-m}^x = 1$.

Next, the distribution $K^x(y, -t)$ belongs to $\mathcal{K}_{v, -(d+2)}(G \times \mathbb{R}_{(m)})$ and is a fundamental solution of $P^x - \partial_t = (P^x + \partial_t)^t$, so the same arguments as above show that the left-convolution operator $\tilde{Q}^x$ with $K^x(y, -t)$ is a left-invariant Volterra Ψ_H DO such that $(P^x + \partial_t)^t \tilde{Q}^x = 1$. Taking transposes we obtain $(\tilde{Q}^x)^t (P^x + \partial_t)$, so if we let $\tilde{q}_{-m}^x \in S_{v, -m}(G \times \mathbb{R}_{(m)})$ be the symbol of $(\tilde{Q}^x)^t$ then we get $\tilde{q}_{-m}^x * (\sigma_m(P)(x, \cdot) + i\tau) = 1$. Henceforth $\sigma_m(P)(x, \cdot) + i\tau$ is a two-sided invertible Volterra-Heisenberg symbol on $G \times \mathbb{R}$ with inverse $q_{-m}^x = \tilde{q}_{-m}^x$.

On the other hand, since for every $x \in U$ the operator $P^x + \partial_t$ satisfies the Rockland condition on $G \times \mathbb{R}$, it follows from [CGGP, Thm. 5(d)] that there exists $\tilde{K}(x, y, t) \in C^\infty(U \times [(G \times \mathbb{R}) \setminus 0])$ homogeneous of degree $-(d+2)$ with respect to (y, t) such that, for every $x \in U$, the left-convolution operator by $\tilde{K}(x, \cdot, \cdot)$ is a right-inverse for $P^x + \partial_t$. Necessarily, this operator agrees with the two-sided inverse Q^x , so K^x agrees with $\tilde{K}(x, \cdot, \cdot)$.

This shows that K^x depends smoothly only on x , that is, $K(x, y, t) = K^x(x, t)$ is an element of $\mathcal{K}_{v, -(d+2)}(U \times G \times \mathbb{R})$. It then follows that $q_{-m}(x, \xi, \tau) = q_{-m}^x(xi, \tau) = K_{(y,t) \rightarrow (\xi, \tau)}^\wedge(x, \xi, \tau)$ belongs to class $S_{v, -m}(U \times G \times \mathbb{R}_{(m)})$ and satisfies

$$(8.3) \quad q_{-m} * (\sigma_m(P) + i\tau) = (\sigma_m(P) + i\tau) * q_{-m} = 1.$$

Hence $\sigma_m(P)(x, \xi) + i\tau$ is an invertible Volterra-Heisenberg symbol with inverse $q_{-m}(x, \xi, \tau)$.

Finally, suppose that $\mathcal{E}$ is not the trivial line bundle. Then, as the aforementioned results of [FS2] and [CGGP] remain valid for systems, by working in a local trivializing Heisenberg chart and by arguing as above we can show that the principal symbol $P + \partial_t$ is an invertible Volterra-Heisenberg symbol. $\square$

Granted Proposition 8.3 we can give the following criterion for a selfadjoint hypoelliptic differential operator to have a positive principal symbol.

Proposition 8.4. *Suppose that P satisfies the Rockland condition at every point. Then the following are equivalent:*

- (i) *The operator P is bounded from below.*
- (ii) *The principal symbol $\sigma_m(P)$ of P is positive.*

Proof. Assume that P is bounded from below. Possibly by replacing P by $P + c$ with c large enough we may assume that P is positive. Observe that P^2 satisfies the Rockland condition at every point. Indeed, at every point $x \in M$ the model operator of P^2 is $(P^2)^x = (P^x)^2$, so for any nontrivial irreducible representation π of $G_x M$ on $C^\infty(\pi)$ we have $\overline{\pi(P^2)^x} = \overline{\pi(P^x)^2} = \overline{\pi P^x \pi P^x}$. Since $\overline{\pi P^x}$ is injective on $C^\infty(\pi)$ we see that so is $\overline{\pi(P^2)^x}$, that is, P^2 satisfies the Rockland condition at x .

Granted this, we can apply Lemma 8.3 to deduce that the principal symbol of $P^2 + \partial_t$ is an invertible Volterra-Heisenberg symbol. As P^2 is positive we then may use Theorem 6.26 to see that $(P^2)^{\frac{1}{4}} = P^{\frac{1}{2}}$ is a Ψ_H DO of order $\frac{m}{2}$. Since $P = (P^{\frac{1}{2}})^2$ it then follows from Lemma 8.2 that the principal symbol of P is a positive symbol.

Conversely, suppose that P has a positive principal symbol. Then thanks to Lemma 8.2 the operator P can be written as $P = Q^*Q + R$ with $Q \in \Psi_H^{\frac{m}{2}}(M, \mathcal{E})$ and $R \in \Psi_H^{m-1}(M, \mathcal{E})$. Let $P_1 = Q^*Q$. Since P and P_1 have same principal symbol, the operator P_1 also satisfies the Rockland condition at every point.

Next, since P_1 is positive for $\lambda < -1$ we have

$$(8.4) \quad (P_1 - \lambda)^{-1} = \int_0^\infty e^{-tP_1} e^{t\lambda} dt,$$

where the integral converges in $\mathcal{L}(L^2(M, \mathcal{E}))$. Let $\alpha = \frac{m-1}{m}$. Then we have

$$(8.5) \quad R(P_1 - \lambda)^{-1} = R P_1^{-\alpha} R_{(\lambda)}, \quad R_{(\lambda)} = \int_0^\infty P_1^\alpha e^{-tP_1} e^{t\lambda} dt.$$

Since $P_1 = Q^*Q$ has a positive principal symbol and satisfies the Rockland condition, Lemma 8.3 tells us that the principal symbol of $P_1 + \partial_t$ is an invertible Volterra-Heisenberg symbol. This allows us to apply Theorem 6.29 to deduce that $P_1^{-\alpha}$ is a Ψ_H DO of order $-m\alpha = -(m-1)$. Therefore $RP_1^{-\alpha}$ is a Ψ_H DO of order 0, hence is bounded on $L^2(M, \mathcal{E})$.

On the other hand, we have

$$(8.6) \quad \|R_{(\lambda)}\|_{L^2} \leq \int_0^\infty t^{-\alpha} \|(tP_1)^\alpha e^{-tP_1}\| e^{t\lambda} dt.$$

As $\alpha \in (0, 1)$ the function $x \rightarrow x^\alpha e^{-x}$ maps $[0, \infty)$ to $[0, 1]$. Therefore, we have $\|(tP_1)^\alpha e^{-tP_1}\| \leq 1$, from which we get

$$(8.7) \quad \|R_{(\lambda)}\|_{L^2} \leq \int_0^\infty t^{-\alpha} e^{t\lambda} dt = |\lambda|^{\alpha-1} \int_0^\infty u^{-\alpha} e^{-u} du.$$

Now, by combining (8.5)–(8.7) we obtain

$$(8.8) \quad \|R(P_1 - \lambda)^{-1}\|_{L^2} \leq \|RP_1^{-\alpha}\|_{L^2} \|R_{(\lambda)}\|_{L^2} \leq C_\alpha |\lambda|^{\alpha-1},$$

where the constant C_α does not depend on λ . Since $\alpha < 1$ it follows that for λ negatively large enough we have $\|R(P_1 - \lambda)^{-1}\|_{L^2} \leq \frac{1}{2}$, so that $1 + R(P_1 - \lambda)^{-1}$ is invertible. Since we have

$$(8.9) \quad P - \lambda = P_1 + R - \lambda = (P_1 + R - \lambda)(P_1 - \lambda)^{-1},$$

it follows that $P - \lambda$ has a right inverse for λ negatively large enough. Since we can similarly show that for λ negatively large enough $P - \lambda$ is left-invertible, we deduce that as soon as λ negatively large enough $P - \lambda$ admits a bounded two-sided inverse. This means that the spectrum of P is contained in some interval $[c, \infty)$, that is, P is bounded from below. $\square$

Now, by combining Lemma 8.3 and Proposition 8.4 with Proposition 6.21 and Proposition 6.22 we get the main result of this section:

Theorem 8.5. *Assume that the Levi form of (M, H) has constant rank and let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of even Heisenberg order m such that P is bounded from below and satisfies the Rockland condition at every point. Then:*

- 1) *The principal symbol $P + \partial_t$ is an invertible Volterra-Heisenberg symbol;*
- 2) *The heat operator $P + \partial_t$ has an inverse in $\Psi_{H,v}^{-m}(M \times \mathbb{R}_{(m)}, \mathcal{E})$;*
- 3) *The heat kernel $k_t(x, y)$ of P has an asymptotics in $C^\infty(M, (\text{End } \mathcal{E}) \otimes |\Lambda|(M))$ of the form*

$$(8.10) \quad k_t(x, x) \sim_{t \rightarrow 0^+} t^{-\frac{d+2}{m}} \sum t^{\frac{2j}{m}} a_j(P)(x), \quad a_j(P)(x) = |\varepsilon'_x| \check{q}_{-m-2j}(x, 0, 1),$$

where the equality on the right-hand side shows how to compute $a_j(P)(x)$ in a local trivializing Heisenberg chart by means of the (local) symbol $q_{-m-2j}(x, \xi, \tau)$ of degree $-m-2j$ of any Volterra- Ψ_H DO parametrix for $P + \partial_t$ in this chart.

Example 8.6. Theorem 8.5 holds for the following examples:

- (a) The conformal powers $\square_\theta^{(k)}$ of the horizontal sublaplacian acting on functions on a compact strictly pseudoconvex CR manifold;
- (b) The contact Laplacian on a compact orientable contact manifold.

9. SPECTRAL ASYMPTOTICS FOR HYPOELLIPTIC OPERATORS

In this section we will make use of Theorem 8.5 to derive spectral asymptotics for hypoelliptic operators on Heisenberg manifolds.

Let (M^{d+1}, H) be a compact Heisenberg manifold equipped with a smooth positive density and let $\mathcal{E}$ be a Hermitian vector bundle over M . Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of Heisenberg order m such that P is bounded from below and the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol. Recall that by Proposition 6.23 and Theorem 8.5 the latter condition is satisfied in the following cases:

- The operator P is a sublaplacian and satisfies the condition (4.57) at every point.
- The Levi form of (M, H) has constant rank and P satisfies the Rockland condition at every point.

Let $\lambda_0(P) \leq \lambda_1(P) \leq \dots$ denote the eigenvalues of P counted with multiplicity and let $N(P; \lambda)$ be the counting function of P , that is,

$$(9.1) \quad N(P; \lambda) = \#\{k \in \mathbb{N}; \lambda_k(P) \leq \lambda\}, \quad \lambda \in \mathbb{R}.$$

Theorem 9.1. *Under the above assumptions the following hold.1) As $t \rightarrow 0^+$ we have*

$$(9.2) \quad \text{Tr } e^{-tP} \sim t^{-\frac{d+2}{m}} \sum t^{\frac{2j}{m}} A_j(P), \quad A_j(P) = \int_M \text{tr}_{\mathcal{E}} a_j(P)(x),$$

where the density $a_j(P)(x)$ is the coefficient of $t^{\frac{j-d+2}{m}}$ in the heat kernel asymptotics (8.10) for P .

2) We have $A_0(P) > 0$.

3) As $\lambda \rightarrow \infty$ we have

$$(9.3) \quad N(P; \lambda) \sim \nu_0(P) \lambda^{\frac{d+2}{m}}, \quad \nu_0(P) = \Gamma(1 + \frac{d+2}{m})^{-1} A_0(P).$$

4) As $k \rightarrow \infty$ we have

$$(9.4) \quad \lambda_k(P) \sim \left(\frac{k}{\nu_0(P)} \right)^{\frac{m}{d+2}}.$$

Proof. First, as $\text{Tr } e^{-tP} = \int_M \text{tr}_{\mathcal{E}} k_t(x, x)$ the asymptotics (9.2) immediately follows from (8.10).

Second, we have $A_0(P) = \lim_{t \rightarrow 0^+} t^{\frac{d+2}{m}} \text{Tr } e^{-tP} \geq 0$, so if we can show that $A_0(P) \neq 0$ then we get $A_0(P) > 0$ and the asymptotics (9.3) will follow from (9.2) by Karamata's Tauberian theorem (see [Ha, Thm. 108]). This will also yield (9.4), since the latter is equivalent to (9.3) (e.g. [Sh, Sect. 13]). Therefore, the bulk of the proof is to show that $A_0(P) \neq 0$.

Now, notice that there is at least one integer $< \frac{m}{d+2}$ such that $A_j(P) \neq 0$. Otherwise, by (9.2) there would exist a constant $C > 0$ such that $\text{Tr } e^{-tP} \leq C$ for $0 \leq t \leq 1$. Thus,

$$(9.5) \quad k e^{-t\lambda_k(P)} \leq \sum_{j < k} e^{-t\lambda_j(P)} \leq \text{Tr } e^{-tP} \leq C, \quad 0 < t < 1.$$

Therefore, letting $t \rightarrow 0^+$ would give $k \leq C$ for every $k \in \mathbb{N}$, which is not possible.

On the other hand, when $m \geq d+2$ the only integer $< \frac{m}{d+2}$ is $j = 0$, so in this case we must have $A_0(P) \neq 0$.

Next, assume $m < d+2$ and $A_0(P) = 0$. Let $\mu = \frac{d+2}{m} - j_0$ where j_0 is the smallest integer j such that $A_j(P) \neq 0$. Since $1 \leq j_0 < \frac{d+2}{m}$ we have $0 < \mu \leq \frac{d+2}{m} - 1$. Moreover, the asymptotics (9.2) becomes

$$(9.6) \quad \text{Tr } e^{-tP} = A_{j_0}(P) t^{-\mu} + O(t^{1-\mu}) \quad \text{as } t \rightarrow 0^+.$$

This implies that we have $A_{j_0}(P) \geq 0$. Since we have $A_{j_0}(P) \neq 0$ by definition of j_0 , we get $A_{j_0}(P) > 0$. Therefore, as alluded to above, it follows from Karamata's Tauberian theorem that as $k \rightarrow \infty$ we have

$$(9.7) \quad \lambda_k(P) \sim \left(\frac{k}{\beta}\right)^{\frac{1}{\mu}}, \quad \beta = \Gamma(1 + \mu)^{-1} A_{j_0}(P).$$

It follows that $\lambda_k(P^{-\frac{d+2}{2m}}) = O(k^{-\frac{1}{2}-\delta})$, with $\delta = \frac{1}{2}(\frac{1}{\mu}\frac{d+2}{m} - 1) > 0$. In particular, we have $\sum_{k \geq 0} \lambda_k(P^{-\frac{d+2}{2m}})^2 < \infty$, that is, $P^{-\frac{d+2}{2m}}$ is a Hilbert-Schmidt operator on $L^2(M, \mathcal{E})$.

In addition, observe that $P^{-\frac{d+2}{2m}}$ is a Ψ_H DO of order $-\frac{(d+2)}{2}$ and that any $Q \in \Psi_H^{-\frac{(d+2)}{2}}(M, \mathcal{E})$ can be written as

$$(9.8) \quad Q = \Pi_0(P)Q + (1 - \Pi_0(P))P^{-\frac{d+2}{2m}}P^{\frac{d+2}{2m}}Q,$$

where $\Pi_0(P)$ denotes the orthogonal projection onto $\ker P$. Recall that the space of Hilbert-Schmidt operators is a two-sided ideal of $\mathcal{L}(L^2(M, \mathcal{E}))$. Observe that in (9.8) the projection $\Pi_0(P)$ is a smoothing operator, so is a Hilbert-Schmidt operator and $P^{\frac{d+2}{2m}}Q$ is a bounded operator on $L^2(M, \mathcal{E})$ because this is a zero'th order Ψ_H DO. It thus follows that any $Q \in \Psi_H^{-\frac{(d+2)}{2}}(M, \mathcal{E})$ is a Hilbert-Schmidt operator on $L^2(M, \mathcal{E})$.

Now, we get a contradiction as follows. Let $\kappa : U \rightarrow V$ be a Heisenberg chart over which there is a trivialization $\tau : \mathcal{E}_U \rightarrow U \times \mathbb{C}^r$ of $\mathcal{E}$ and such that the open $V \subset \mathbb{R}^{d+1}$ is bounded. Let $\varphi \in C_c^\infty(\mathbb{R}^{d+1})$ have non-empty support L , let $\chi \in C_c^\infty(V \times V)$ be such that $\chi(x, y) = 1$ near $L \times L$, and let $Q : C_c^\infty(V) \rightarrow C^\infty(V)$ be given by the kernel,

$$(9.9) \quad k_Q(x, y) = |\varepsilon'_x|\varphi(x)\|\varepsilon_x(y)\|^{-\frac{d+2}{2}}\chi(x, y).$$

The kernel $k_Q(x, y)$ has a compact support contained in $V \times V$ and, as $\varphi(x)(1 - \chi(x, y))$ vanishes near the diagonal of $V \times V$, we have

$$(9.10) \quad k_Q(x, y) = |\varepsilon'_x|\varphi(x)\|\varepsilon_x(y)\|^{-\frac{d+2}{2}} \bmod C^\infty(V \times V).$$

Since $\varphi(x)\|y\|^{-\frac{d+2}{2}}$ belongs to $\mathcal{K}^{-\frac{d+2}{2}}(V \times \mathbb{R}^{d+1})$, it follows from Proposition 4.17 that Q is a Ψ_H DO of order $-\frac{d+2}{2}$.

Let $Q_0 = \tau^*\kappa^*(Q \otimes 1)$. Then Q_0 belongs to $\Psi_H^{-\frac{d+2}{2}}(M, \mathcal{E})$, hence is a Hilbert-Schmidt operator on $L^2(M, \mathcal{E})$. This implies that Q is a Hilbert-Schmidt operator on $L^2(V)$, so by [GK, p. 109] its kernel lies in $L^2(V \times V)$. This cannot be true, however, because we have

$$(9.11) \quad \begin{aligned} \int_{V \times V} |k_Q(x, y)|^2 dx dy &\geq \int_{L \times L} |\varepsilon'_x|^2 |\varphi(x)|^2 \|\varepsilon_x(y)\|^{-(d+2)} dx dy \\ &= \int_L |\varepsilon'_x| |\varphi(x)|^2 \left(\int_{\varepsilon_x(L)} \|y\|^{-(d+2)} dy \right) dx = \infty. \end{aligned}$$

We have thus obtained a contradiction, so we must have $A_0(P) \neq 0$. The proof is now complete. $\square$

Example 9.2. Theorem 9.1 is valid for the following operators:

(a) Real selfadjoint sublaplacian $\Delta = \nabla_{X_1}^* \nabla_{X_1} + \dots + \nabla_{X_m}^* \nabla_{X_m}^2 + \mu(x)$, where ∇ is connection on $\mathcal{E}$, the vector fields $X_1, \dots, X_m$ span H and $\mu(x)$ is a selfadjoint section of $\text{End } \mathcal{E}$, provided that the Levi form of (M, H) is non-vanishing;

(b) The Kohn Laplacian on a compact CR manifold and acting on (p, q) -forms when the condition $Y(q)$ holds everywhere;

(c) The horizontal sublaplacian on a compact Heisenberg manifold acting on horizontal forms of degree k when the condition $X(k)$ holds everywhere;

- (d) The contact Laplacian on a compact orientable contact manifold.
- (e) The conformal powers of the horizontal sublaplacian acting on functions on a compact strictly pseudoconvex CR manifold.

Several authors have obtained Weyl asymptotics closely related to (1.12) for bicharacteristic hypoelliptic operators ([II], [Me1], [MS]), including sublaplacians on Heisenberg manifolds, and for more general hypoelliptic operators ([Mo1], [Mo2]) using different approaches involving other pseudodifferential calculi.

While these authors deal in a setting more general than the Heisenberg setting, as far as the Heisenberg setting is concerned, our approach using the Volterra-Heisenberg calculus presents the following advantages:

- (i) The pseudodifferential analysis is somewhat simpler, since the Volterra-Heisenberg calculus yields for free a heat kernel asymptotics, once the principal symbol of the heat operator is shown to be invertible, for which it is enough to use the Rockland condition in many cases;
- (ii) As the Volterra-Heisenberg calculus take fully into account the underlying Heisenberg geometry of the manifold and is invariant by change of Heisenberg coordinates, we can explicitly compute the coefficient $\nu_0(P)$ in (9.3) for operators admitting a normal form. This is illustrated below by Proposition 9.4, which will be used in the next two section to give geometric expressions for the Weyl asymptotics (9.3) for geometric operators on CR and contact manifolds.

Next, prior to dealing with operators admitting a normal form we need the following.

Lemma 9.3. *Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of Heisenberg order m such that P is bounded from below and the principal symbol of $P + \partial_t$ is an invertible Volterra-Heisenberg symbol. Then, for any $a \in M$, the following hold.*

1) *The model operator $P^a + \partial_t$ admits a unique fundamental solution $K^a(x, t)$ which, in addition, belongs to the class $\mathcal{K}_{v, -(d+2)}(G_a M \times \mathbb{R}_{(m)})$.*

2) *In Heisenberg coordinates centered at a we have*

$$(9.12) \quad a_0(P)(0) = K^a(0, 1).$$

Proof. Since it is enough to prove the result in a trivializing Heisenberg chart near a , we may as well assume that $\mathcal{E}$ is the trivial line bundle, since the proof in the general case follows along similar lines.

Let $\sigma_{-m}(Q) \in S_{v, -m}(\mathfrak{g}^*U \times \mathbb{R}_{(m)})$ be the inverse of $\sigma_m(P) + i\tau$ and let Q^a be the left-invariant Volterra- Ψ_H DO on $G_a U \times \mathbb{R}$ with symbol $\sigma_{-m}(Q)(a, \cdot, \cdot)$. In particular, the operator Q^a is the inverse of $P^a + \partial_t$ on $\mathcal{S}(G_a U \times \mathbb{R})$ and it agrees with the left-convolution by $K^a = [\sigma_{-m}(a, \cdot, t)]_{(\xi, \tau) \rightarrow (y, t)}^\vee$. Therefore, the left-convolution operator with $[(P^a + \partial_t)K^a]$ agrees with $(P^a + \partial_t)Q^a = 1$. Thus,

$$(9.13) \quad (P^a + \partial_t)K^a(y, t) = \delta(y, t),$$

that is, $K^a(y, t)$ is a fundamental solution for $P^a + \partial_t$.

Let $K \in \mathcal{S}'(G_a U \times \mathbb{R})$ be another fundamental solution for $P^a + \partial_t$. As it follows from the proof of Lemma 8.3, the left-convolution operator Q by K is a right-inverse for $P^a + \partial_t$, so agrees with Q^a . Hence $K = K^a$, which shows that K^a is the unique fundamental solution of $P^a + \partial_t$.

In addition, K^a is in the class $\mathcal{K}_{v, -(d+2)}(G_a M \times \mathbb{R}_{(m)})$ because this the inverse Fourier transform of a symbol in $S_{v, -m}(\mathfrak{g}^*U \times \mathbb{R}_{(m)})$.

Finally, let $q_{-m} \in S_{v, -m}(U \times \mathbb{R}^{d+1} \times \mathbb{R}_{(m)})$ be the local principal symbol for a parametrix Q for $P + \partial_t$ on $U \times \mathbb{R}$. As Q has global principal symbol $\sigma_{-m}(Q)$ it follows from Proposition 6.15 that $q_{-m}(0, \cdot, \cdot) = \sigma_{-m}(Q)(a, \cdot, \cdot)$. Moreover, since we are in the Heisenberg coordinates centered at a , the map ε_0 to the Heisenberg coordinates centered at 0 is just the identity. Combining this with (9.3) then gives

$$(9.14) \quad a_0(P)(0) = |\varepsilon'_0| \check{q}_m(0, 0, 1) = [\sigma_{-m}(Q)]_{(\xi, \tau) \rightarrow (y, t)}(a, 0, 1) = K^a(0, 1).$$

The proof is thus achieved. $\square$

Now, assume that the Levi form of (M, H) has constant rank $2n$. Therefore, by Proposition 2.7 the tangent Lie group bundle GM is a fiber bundle with typical fiber $G = \mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$.

Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint differential operator of Heisenberg order m such that P is bounded from below and satisfies the Rockland condition at every point.

We further assume that the density $d\rho(x)$ of M and the model operator of P at a point admit normal forms. By this it is meant that there exist $\rho_0 > 0$ and a differential operator $P_0 : C^\infty(G, \mathbb{C}^r) \rightarrow C^\infty(G, \mathbb{C}^r)$ such that near every point x_0 , there exist trivializing Heisenberg coordinates, herewith called normal trivializing Heisenberg coordinates centered at x_0 , in which the following hold:

- (i) We have $d\rho(x)|_{x=0} = [\rho_0 dx]|_{x=0}$;
- (ii) At $x = 0$ the tangent group is G and the model operator of P is P_0 .

In the sequel, we let $\text{vol}_\rho M$ denote the volume of M with respect to ρ , that is,

$$(9.15) \quad \text{vol}_\rho M = \int_M d\rho(x).$$

As it turns out the assumptions (i) and (ii) allows us to relate the Weyl asymptotics (9.3) for P to $\text{vol}_\rho M$ as follows.

Proposition 9.4. *Under the assumptions (i) and (ii) above, as $\lambda \rightarrow \infty$ we have*

$$(9.16) \quad N(P; \lambda) \sim \rho_0^{-1} \nu_0(P_0) (\text{vol}_\rho M) \lambda^{\frac{d+2}{m}}, \quad \nu_0(P_0) = \Gamma(1 + \frac{d+2}{m})^{-1} \text{tr}_{\mathbb{C}^r} K_0(0, 1),$$

where $K_0(x, t)$ denotes the fundamental solution of $P_0 + \partial_t$.

Proof. By Proposition 9.1 as $\lambda \rightarrow \infty$ we have

$$(9.17) \quad N(P; \lambda) \sim \nu_0(P) \lambda^{\frac{d+2}{m}}, \quad \nu_0(P) = \Gamma(1 + \frac{d+2}{m})^{-1} \int_M \text{tr}_{\mathcal{E}} a_0(P)(x).$$

On the other hand, by Lemma 9.3 in normal trivializing Heisenberg coordinates centered at point $a \in M$ we have

$$(9.18) \quad [\text{tr}_{\mathbb{C}^r} a_0(P)(x) dx]|_{x=0} = [\text{tr}_{\mathbb{C}^r} K_0(0, 1) dx]|_{x=0} = \text{tr}_{\mathbb{C}^r} K_0(0, 1) \rho_0^{-1} [d\rho(x)]|_{x=0}.$$

Hence $\text{tr}_{\mathcal{E}} a_0(P)(x) = \rho_0^{-1} \text{tr}_{\mathbb{C}^r} K_0(0, 1) d\rho(x)$. Thus,

$$(9.19) \quad \nu_0(P) = \Gamma(1 + \frac{d+2}{m})^{-1} \int_M \rho_0^{-1} \text{tr}_{\mathbb{C}^r} K_0(0, 1) d\rho(x) = \rho_0^{-1} \nu_0(P_0) \text{vol}_\rho M,$$

with $\nu_0(P_0) = \Gamma(1 + \frac{d+2}{m})^{-1} \text{tr}_{\mathbb{C}^r} K_0(0, 1)$. Combining this (9.17) then proves the lemma. $\square$

10. WEYL ASYMPTOTICS AND CR GEOMETRY

The aim of this section is to express in geometric terms the Weyl asymptotics (9.3) for the Kohn Laplacian and the horizontal sublaplacian on a nondegenerate CR manifold with a Levi metric.

Let (M^{2n+1}, θ) be a κ -strictly pseudoconvex pseudohermitian manifold with $0 \leq \kappa \leq \frac{n}{2}$. Let $T_{1,0} \subset T_{\mathbb{C}}M$ be the CR structure of M and define $T_{0,1} = \overline{T_{1,0}}$ and $H = \Re(T_{1,0} \otimes T_{0,1})$. The Levi form L_θ on $T_{1,0}$ is given by

$$(10.1) \quad L_\theta(Z, W) = -id\theta(Z, \bar{W}) = i\theta([Z, \bar{W}]), \quad Z, W \in T_{1,0}.$$

Let X_0 be the Reeb field associated to θ , so that $\iota_{X_0}\theta = 1$ and $\iota_{X_0}d\theta = 0$. Then we have

$$(10.2) \quad [Z, \bar{W}] = -iL_\theta(Z, W)X_0 \mod T_{1,0} \oplus T_{0,1} \quad \forall Z, W \in T_{1,0}M.$$

We endow M with a Levi metric as follows (see also [FS1]). Let $\tilde{h}$ be a (positive definite) Hermitian metric on $T_{1,0}$. Then there exists a Hermitian-valued section A of $\text{End}_{\mathbb{C}} T_{1,0}$ such that

$$(10.3) \quad L_{\theta}(Z, W) = \tilde{h}(AZ, W), \quad Z, W \in T_{1,0}.$$

Let $A = U|A|$ be the polar decomposition of A . Since by assumption the Hermitian form L_{θ} has signature $(n - \kappa, \kappa, 0)$, hence is nondegenerate, the section A is invertible and the section U is an orthogonal matrix with only the eigenvalues 1 and -1 with multiplicities $n - \kappa$ and κ respectively. Therefore, we have the splitting,

$$(10.4) \quad T_{1,0} = T_{1,0}^+ \oplus T_{1,0}^-, \quad T_{1,0}^{\pm} = \ker(U \mp 1),$$

where the restriction of L_{θ} to $T_{1,0}^+$ (resp. $T_{1,0}^-$) is positive definite (resp. negative definite). We then get a Levi metric on $T_{1,0}$ by letting

$$(10.5) \quad h(Z, W) = \tilde{h}(|A|Z, W), \quad Z, W \in T_{1,0}.$$

In particular, on the direct summands $T_{1,0}^{\pm}$ of the splitting (10.4) we have

$$(10.6) \quad L_{\theta}(Z, W) = h(UZ, W) = \pm h(Z, W), \quad Z, W \in T_{1,0}^{\pm}.$$

We now extend h into a Hermitian metric h on $T_{\mathbb{C}}M$ by making the following requirements:

$$(10.7) \quad h(X_0, X_0) = 1, \quad h(Z, W) = \overline{h(\bar{Z}, \bar{W})} \quad \forall Z, W \in T_{0,1},$$

$$(10.8) \quad \text{The splitting } T_{\mathbb{C}}M = \mathbb{C}X_0 \oplus T_{1,0} \oplus T_{0,1} \text{ is orthogonal with respect to } h.$$

This allows us to express the Levi form $\mathcal{L}_{\mathbb{C}} : (H \otimes \mathbb{C}) \times (H \otimes \mathbb{C}) \rightarrow T_{\mathbb{C}}M/(H \otimes \mathbb{C})$ as follows. Since $\theta(X_0) = 1$ we have

$$(10.9) \quad \mathcal{L}_{\mathbb{C}}(X, Y) = \theta([X, Y])X_0 = -d\theta(X, Y)X_0, \quad X, Y \in H \otimes \mathbb{C}.$$

Therefore, it follows from (10.6) that we have

$$(10.10) \quad \mathcal{L}(Z, \bar{W}) = -iL_{\theta}(Z, W) = h(Z, iUW), \quad Z, W \in T_{1,0}.$$

Since $\mathcal{L}$ is antisymmetric and the integrability condition $[T_{1,0}, T_{1,0}] \subset T_{1,0}$ implies that $\mathcal{L}_{\mathbb{C}}$ vanishes on $T_{1,0} \times T_{1,0}$ and on $T_{0,1} \times T_{0,1}$, we get

$$(10.11) \quad \mathcal{L}_{\mathbb{C}}(X, Y) = h(X, LY), \quad X, Y \in H \otimes \mathbb{C},$$

where L is the antilinear antisymmetric section of $\text{End}_{\mathbb{R}}(H \otimes \mathbb{C})$ such that

$$(10.12) \quad L(Z + \bar{W}) = iUW - i(\bar{U}\bar{Z}), \quad Z, W \in T_{1,0}.$$

In particular, since $U^* = U$ and $U^2 = 1$ we have $|L| = 1$. Let $Z_1, \dots, Z_n$ be a local orthonormal frame for $T_{1,0}$ (with respect to h) and such that $Z_1, \dots, Z_{n-\kappa}$ span $T_{1,0}^+$ and $Z_{n-\kappa+1}, \dots, Z_n$ span $T_{1,0}^-$. Then $\{X_0, Z_j, \bar{Z}_j\}$ is an orthonormal frame. In the sequel we will call such a frame an *admissible orthonormal frame of $T_{\mathbb{C}}M$* . Then from (10.6) we get:

$$(10.13) \quad L_{\theta}(Z_j, \bar{Z}_k) = \epsilon_j h(Z_j, Z_k) = \epsilon_j \delta_{jk},$$

where $\epsilon_j = 1$ for $j = 1, \dots, n - \kappa$ and $\epsilon_j = -1$ for $j = n - \kappa + 1, \dots, n$.

Let $\{\theta, \theta^j, \bar{\theta}^j\}$ be the dual coframe of $T_{\mathbb{C}}^*$ associated to $\{X_0, Z_j, \bar{Z}_j\}$. Then the volume form of h is locally given by

$$(10.14) \quad \sqrt{h(x)}dx = i^n \theta \wedge \theta^1 \wedge \theta^{\bar{1}} \wedge \dots \wedge \theta^n \wedge \theta^{\bar{n}}.$$

On the other hand, because of (10.13) we have $d\theta = i \sum_{j=1}^n \epsilon_j \theta^j \wedge \bar{\theta}^j \mod \theta \wedge T^*M$, so the form $\theta \wedge d\theta^n$ is equal to

$$(10.15) \quad n! i^n \epsilon_1 \dots \epsilon_n \theta \wedge \theta^1 \wedge \theta^{\bar{1}} \wedge \dots \wedge \theta^n \wedge \theta^{\bar{n}} = n! i^n (-1)^{\kappa} \theta \wedge \theta^1 \wedge \theta^{\bar{1}} \wedge \dots \wedge \theta^n \wedge \theta^{\bar{n}}.$$

Therefore, we get the following global formula for the volume form,

$$(10.16) \quad \sqrt{h(x)}dx = \frac{(-1)^\kappa}{n!} \theta \wedge d\theta^n.$$

In particular, we see that the volume form depends only on θ and not on the choice of the Levi metric.

Definition 10.1. *The pseudohermitian volume of (M, θ) is*

$$(10.17) \quad \text{vol}_\theta M = \frac{(-1)^\kappa}{n!} \int_M \theta \wedge d\theta^n.$$

We shall now relate the asymptotics (9.3) for the Kohn Laplacian and the Tanaka sublaplacian to the volume $\text{vol}_\theta M$. To this end consider the Heisenberg group $\mathbb{H}^{2n+1} = \mathbb{R} \times \mathbb{R}^{2n}$ equipped with the standard left-invariant basis of $\mathfrak{h}^{2n+1}$ given by

$$(10.18) \quad X_0 = \frac{\partial}{\partial x_0}, \quad X_j = \frac{\partial}{\partial x_j} + x_{n+j} \frac{\partial}{\partial x_0}, \quad X_{n+j} = \frac{\partial}{\partial x_{n+j}} - x_j \frac{\partial}{\partial x_0}, \quad 1 \leq j \leq n.$$

For $\mu \in \mathbb{C}$ let $\mathcal{L}_\mu$ denote the Folland-Stein sublaplacian,

$$(10.19) \quad \mathcal{L}_\mu = -\frac{1}{2}(X_1^2 + \dots + X_{2n}^2) - i\mu X_0.$$

The Rockland condition for $\mathcal{L}_\mu$ reduces to $\mu \neq \pm n, \pm(n+1), \dots$ (see [FS1], [BG], [Tay]), so in this case the operator $\mathcal{L}_\mu$ is hypoelliptic. Moreover, when $\mu \neq \pm n, \pm(n+1), \dots$ the heat operator $\mathcal{L}_\mu + \partial_t$ is also hypoelliptic and admits a unique fundamental solution since its symbol is an invertible Volterra-Heisenberg symbol (see [BGS]).

In the sequel it will be convenient to use the variable $x' = (x_1, \dots, x_{2n})$.

Lemma 10.2. *For $|\Re \mu| < n$ the fundamental solution of $\mathcal{L}_\mu + \partial_t$ is given by*

$$(10.20) \quad k_\mu(x_0, x', t) = \chi(t)(2\pi t)^{-(n+1)} \int_{-\infty}^{\infty} e^{it^{-1}x_0\xi_0 - \mu\xi_0} \left(\frac{\xi_0}{\sinh \xi_0}\right)^n \exp\left[-\frac{1}{2t} \frac{\xi_0}{\tanh \xi_0} |x'|^2\right] d\xi_0.$$

where $\chi(t)$ denotes the characteristic function of $(0, \infty)$.

Proof. The fundamental solution $k_\mu(x_0, x', t)$ is solution to the equation,

$$(10.21) \quad (\mathcal{L}_\mu + \partial_t)k(x_0, x', t) = \delta(x_0) \otimes \delta(x') \otimes \delta(t).$$

Let us make a Fourier transform with respect to x_0 . Thus, letting $\hat{k}(\xi_0, x', t) = \hat{k}_{x_0 \rightarrow \xi_0}(\xi_0, x', t)$, the equation (10.21) becomes

$$(10.22) \quad (\hat{\mathcal{L}}_0 + \mu\xi_0 + \partial_t)\hat{k}_\mu = \delta(x') \otimes \delta(t),$$

$$(10.23) \quad \hat{\mathcal{L}}_0 = -\frac{1}{2} \sum_{j=1}^n \left(\frac{\partial}{\partial x_j} - ix_{n+j}\xi_0 \right)^2 - \frac{1}{2} \sum_{j=1}^n \left(\frac{\partial}{\partial x_{n+j}} + ix_j\xi_0 \right)^2.$$

Notice that if we fix ξ_0 we have $\hat{\mathcal{L}}_0 = \frac{1}{2}H_{A(\xi_0)}$, where $H_{A(\xi_0)} = -\sum_{j=1}^{2n} (\partial_j - \sum_{k=1}^{2n} A(\xi_0)_{jk}x_k)^2$ is the harmonic oscillator associated to the real antisymmetric matrix,

$$(10.24) \quad A(\xi_0) = \xi_0 J, \quad J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}.$$

Therefore, the fundamental solution of $\hat{\mathcal{L}}_0 + \mu\xi_0 + \partial_t$ is given by a version of the Melher formula (see [GJ], [Po4, p. 225]), that is,

$$(10.25) \quad \hat{k}_0(\xi_0, x', t) = \chi(t)(2\pi t)^{-n} \det^{\frac{1}{2}}\left(\frac{tA(\xi_0)}{\sinh(tA(\xi_0))}\right) \exp\left[-\frac{1}{2t}\left\langle \frac{tA(\xi_0)}{\tanh(tA(\xi_0))}x', x'\right\rangle\right],$$

$$= \chi(t)(2\pi t)^{-n} \left(\frac{t\xi_0}{\sinh t\xi_0}\right)^n \exp\left[-\frac{1}{2t} \frac{t\xi_0}{\tanh t\xi_0} |x'|^2\right].$$

A solution of (10.22) is now given by $\hat{k}_\mu(\xi_0, x', t) = e^{-\mu\xi_0 t} \hat{k}_0(\xi_0, x', t)$. Moreover, as we have

$$(10.26) \quad |h_0(\xi_0, x', t)| \leq \pi^{-n} |\xi_0|^n e^{-tn|\xi_0|},$$

we further see that for $|\Re\mu| < n$ the function $e^{-t\mu\xi_0} \hat{k}_0$ is integrable with respect to ξ_0 . Since $k_\mu(x_0, x', t)$ is the inverse Fourier transform with respect to ξ_0 of $\hat{k}_\mu(\xi_0, x', t)$ it follows that we have

$$(10.27) \quad k_\mu(x_0, x', t) = \chi(t)(2\pi t)^{-(n+1)} \int_{-\infty}^{\infty} e^{it^{-1}x_0\xi_0 - \mu\xi_0} \left(\frac{\xi_0}{\sinh \xi_0}\right)^n \exp\left[-\frac{1}{2t} \frac{\xi_0}{\tanh \xi_0} |x'|^2\right] d\xi_0.$$

The lemma is thus proved. $\square$

Next, for $|\Re\mu| < n$ we let

$$(10.28) \quad \nu(\mu) = \frac{1}{(n+1)!} k_\mu(0, 0, 1) = \frac{(2\pi)^{-(n+1)}}{(n+1)!} \int_{-\infty}^{\infty} e^{-\mu\xi_0} \left(\frac{\xi_0}{\sinh \xi_0}\right)^n d\xi_0.$$

Lemma 10.3. *Let $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a selfadjoint sublaplacian which is bounded from below and assume there exists $\mu \in (-n, n)$ such that near any point of M there is an admissible orthonormal frame $Z_1, \dots, Z_n$ of $T_{1,0}$ with respect to which Δ takes the form,*

$$(10.29) \quad \Delta = - \sum_{j=1}^n (\bar{Z}_j Z_j + Z_j \bar{Z}_j) - i\mu X_0 + \text{lower order terms}.$$

Then as $\lambda \rightarrow \infty$ we have

$$(10.30) \quad N(\Delta; \lambda) \sim \nu(\mu) \operatorname{rk} \mathcal{E} (\operatorname{vol}_\theta M) \lambda^{n+1}.$$

Proof. Let $Z_1, \dots, Z_n$ be a local admissible orthonormal frame of $T_{1,0}$. Then from (10.1) and (10.6) we obtain

$$(10.31) \quad [Z_j, \bar{Z}_k] = -L_\theta(Z_j, Z_k)X_0 = -i\epsilon_j \delta_{jk} X_0 \mod T_{1,0} \oplus T_{0,1}.$$

In addition, we let $X_1, \dots, X_{2n}$ be the vector fields in H such that

$$(10.32) \quad Z_j = \begin{cases} \frac{1}{2}(X_j - iX_{n+j}) & \text{for } j = 1, \dots, n - \kappa, \\ \frac{1}{2}(X_{n+j} - iX_j) & \text{for } j = n - \kappa + 1, \dots, n. \end{cases}$$

Then $X_1, \dots, X_0$ is a local frame of $H = \Re(T_{1,0} \oplus T_{0,1})$ and from (10.31) we get

$$(10.33) \quad [X_j, X_{n+k}] = -2\delta_{jk} X_0 \mod H,$$

$$(10.34) \quad [X_0, X_j] = [X_j, X_k] = [X_{n+j}, X_{n+k}] = 0 \mod H.$$

Moreover, in terms of the vector fields $X_1, \dots, X_{2n}$ the formula (10.29) becomes

$$(10.35) \quad \Delta = -\frac{1}{2}(X_1^2 + \dots + X_{2n}^2) + i\mu X_0 + \text{lower order terms}.$$

Combining this with (10.12) shows that the condition (4.57) for Δ , is given in terms of the eigenvalues of $|L| = 1$ and becomes

$$(10.36) \quad \mu \notin \{\pm(n+k); k = 0, 1, 2, \dots\}.$$

Since by assumption we have $\mu \in (-n, n)$, the condition (4.57) is fulfilled at every point, so by Proposition 6.23 the principal symbol of $\Delta + \partial_t$ is an invertible Volterra-Heisenberg symbol. This allows us to apply Theorem 9.1 to deduce that as $\lambda \rightarrow \infty$ we have

$$(10.37) \quad N(\Delta; \lambda) \sim \nu_0(\Delta) \lambda^{n+1},$$

where $\nu_0(P)$ is given by (9.3).

Let us now work in Heisenberg coordinates centered at a point $a \in M$ related to a local H -frame $X_0, X_1, \dots, X_{2n}$ as above. Because of (10.34) the model vector fields of $X_0, \dots, X_{2n}$ at a are

$$(10.38) \quad X_0^a = \frac{\partial}{\partial x_0}, \quad X_j^a = \frac{\partial}{\partial x_j} + x_{n+j} \frac{\partial}{\partial x_0}, \quad X_{n+j}^a = \frac{\partial}{\partial x_{n+j}} - x_j \frac{\partial}{\partial x_0}, \quad 1 \leq j \leq n.$$

These are the vector fields (10.18), so in the Heisenberg coordinates $G_a M$ agrees with the Heisenberg group $\mathbb{H}^{2n+1}$. In addition, using (10.35) we see that the model operator of Δ is

$$(10.39) \quad \Delta^a = -\frac{1}{2}((X_1^a)^2 + \dots + (X_{2n}^a)^2) - i\mu X_0^a = \mathcal{L}_\mu.$$

Since $\mu \in (-n, n)$ it follows that Δ satisfies the Rockland at every point. On the other hand, since we are in Heisenberg coordinates we have $X_j(0) = \frac{\partial}{\partial x_j}$. In particular, we have $\theta(0) = dx_0$. Moreover, as $d\theta(X, Y) = -\theta([X, Y])$ we deduce from (10.34) that

$$(10.40) \quad d\theta(X_j, X_{n+k}) = 2\delta_{jk} \quad \text{and} \quad d\theta(X_j, X_k) = 0,$$

for $j = 0, 1, \dots, n$ and $k = 1, \dots, n$. It then follows that $d\theta(0) = 2 \sum_{j=1}^n dx_j \wedge dx_{n+j}$. Therefore, the volume form is given by

$$(10.41) \quad \frac{(-1)^\kappa}{n!2^n} \theta \wedge d\theta^n(0) = (-1)^\kappa dx_0 \wedge dx_1 \wedge dx_{n+1} \wedge \dots \wedge dx_n \wedge dx_{2n}.$$

Note also that because of (10.32) the orientation of M is that given by $i^n X_0 \wedge Z_1 \wedge \bar{Z}_1 \wedge \dots \wedge Z_n \wedge \bar{Z}_n = (-1)^\kappa X_0 \wedge X_1 \wedge X_n \wedge \dots \wedge X_{n+1} \wedge X_{2n}$. Therefore, at $x = 0$ the volume form of M is in the same orientation class as $(-1)^\kappa dx_0 \wedge dx_1 \wedge dx_{n+1} \wedge \dots \wedge dx_n \wedge dx_{2n}$. In terms of density, together with (10.41), this means that we have

$$(10.42) \quad \frac{(-1)^\kappa}{n!2^n} \theta \wedge d\theta^n(0) = dx|_{x=0}.$$

All this shows that in the above Heisenberg coordinates the volume form and the model operator of Δ have normal forms in the sense of Proposition 9.4 with $\rho_0 = 1$ and $P_0 = \mathcal{L}_\mu \otimes 1_{\mathbb{C}^r}$. Therefore, we may apply (9.16) to get

$$(10.43) \quad \nu_0(\Delta) = \nu_0(\mathcal{L}_\mu \otimes 1_{\mathbb{C}^r}) \int_M \frac{(-1)^\kappa}{n!2^n} \theta \wedge d\theta^n = \nu_0(\mathcal{L}_\mu) \text{rk } \mathcal{E} \text{vol}_\theta M, .$$

where $\nu_0(\mathcal{L}_\mu) = \frac{1}{(n+1)!} k_\mu(0, 0, 1) = \nu(\mu)$. The lemma is thus proved. $\square$

We are now ready to relate the asymptotics (9.3) for the Kohn Laplacian to the pseudohermitian volume of M .

Theorem 10.4. *Let $\square_b : C^\infty(M, \Lambda^{*,*}) \rightarrow C^\infty(M, \Lambda^{*,*})$ be the Kohn Laplacian on M associated to a Levi metric on M . Then for $p, q = 1, \dots, n$, with $q \neq \kappa$ and $q \neq n - \kappa$, as $\lambda \rightarrow \infty$ we have*

$$(10.44) \quad N(\square_b|_{\Lambda^{p,q}}; \lambda) \sim \alpha_{n\kappa pq} (\text{vol}_\theta M) \lambda^{n+1},$$

$$(10.45) \quad \alpha_{n\kappa pq} = \frac{1}{2^{n+1}} \binom{n}{p} \sum_{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa)} \binom{n-\kappa}{k} \binom{\kappa}{q-k} \nu(n-2(\kappa-q+2k)).$$

Proof. As M is κ -strictly pseudoconvex the $Y(q)$ -condition reduces to $q \notin \{\kappa, n - \kappa\}$, so in this case $\square_b^{p,q}$ has an invertible principal symbol. Since $\square_b$ is a positive it follows from Theorem 9.1 that as $\lambda \rightarrow \infty$ we have

$$(10.46) \quad N(\square_b^{p,q}; \lambda) \sim \nu_0(\square_b^{p,q}) \lambda^{n+1}.$$

It then remains to show that $\nu_0(\square_b) = \alpha_{n\kappa pq} \text{vol}_\theta M$ with $\alpha_{n\kappa pq}$ given by (10.45).

Let $\{X_0, Z_j, \bar{Z}_j\}$ be a local admissible orthonormal frame of $T_{\mathbb{C}}M$ and let $\{\theta, \theta^j, \theta^{\bar{j}}\}$ be the dual coframe of $T_{\mathbb{C}}^*M$. For ordered subsets $J = \{j_1, \dots, j_p\}$ and $K = \{k_1, \dots, k_q\}$ of $\{1, \dots, n\}$ with $j_1 < \dots < j_p$ and $k_1 < \dots < k_q$ we let

$$(10.47) \quad \theta^{J, \bar{K}} = \theta^{j_1} \wedge \dots \wedge \theta^{j_p} \wedge \theta^{\bar{k}_1} \wedge \dots \wedge \theta^{\bar{k}_q}.$$

Then $\{\theta^{J, \bar{K}}\}$ form an orthonormal frame of $\Lambda^{*,*}$ and, as shown in [BG, Sect. 20], with respect to this frame on (p, q) -forms $\square_b$ takes the form,

$$(10.48) \quad \square_b|_{\Lambda^{p,q}} = \text{diag}\{\square_{J\bar{K}}\} + \text{lower order terms},$$

$$(10.49) \quad \square_{J\bar{K}} = -\frac{1}{2} \sum_{1 \leq j \leq n} (Z_j \bar{Z}_j + \bar{Z}_j Z_j) + \frac{1}{2} \sum_{j \in K} [Z_j, \bar{Z}_j] - \frac{1}{2} \sum_{j \notin K} [Z_j, \bar{Z}_j].$$

Moreover, using (10.31) we see that the leading part of $\square_{J\bar{K}}$ is equal to

$$(10.50) \quad -\frac{1}{2} \sum_{1 \leq j \leq n} (Z_j \bar{Z}_j + \bar{Z}_j Z_j) - \frac{i}{2} \mu_K X_0, \quad \mu_K = \sum_{j \in K} \epsilon_j - \sum_{j \notin K} \epsilon_j.$$

Notice that since $\epsilon_j = 1$ for $j = 1, \dots, n - \kappa$ and $\epsilon_j = -1$ for $j = n - \kappa + 1, \dots, n$ we have $\mu_K \in [-n, n]$ and $\mu_K = \pm n$ if, and only if, we have $K = \{1, \dots, n - \kappa\}$ or $K = \{n - \kappa + 1, \dots, n\}$. Thus if $|K| = q$ with $q \notin \{\kappa, n - \kappa\}$ then $\square_{J\bar{K}}$ is two times a sublaplacian of the form (10.29) with $\mu = \mu_K$ in $(-n, n)$.

On the other hand, complex conjugation is an isometry, so from the orthogonal splitting (10.4) we get the orthogonal splitting $T_{0,1} = T_{0,1}^+ \oplus T_{0,1}^-$ with $T_{0,1}^\pm = \overline{T_{1,0}^\pm}$. By duality these splittings give rise to the orthogonal decompositions,

$$(10.51) \quad \Lambda^{1,0} = \Lambda_+^{1,0} \oplus \Lambda_-^{1,0}, \quad \Lambda^{0,1} = \Lambda_+^{0,1} \oplus \Lambda_-^{0,1},$$

$$(10.52) \quad \Lambda^{p,q} = \bigoplus_{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa)} \Lambda^{p,q,k}, \quad \Lambda^{p,q,k} = \Lambda^{p,0} \wedge (\Lambda_+^{0,1})^k \wedge (\Lambda_-^{0,1})^{q-k}.$$

Since (10.48) and (10.49) show that $\square_b$ is scalar modulo lower order terms, on $\Lambda^{p,q}$ we can write

$$(10.53) \quad \square_b = \sum_{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa)} \square_{p;q,k} + \text{lower order terms}, \quad \square_{pqk} = \Pi_{pqk} \square_b \Pi_{pqk},$$

where $\Pi_{p;q,k}$ denotes the orthogonal projection onto $\Lambda^{p;q,k}$. In particular, as $\nu_0(\square_b^{p,q})$ depends only on the principal symbol of $\square_b^{p,q}$ we get

$$(10.54) \quad \nu_0(\square_b|_{\Lambda^{p,q}}) = \sum_{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa)} \nu_0(\square_{p;q,k}).$$

We are thus reduced to express each coefficient $\nu_0(\square_{p;q,k})$ in terms of $\text{vol}_\theta M$.

Next, if $\theta^{J\bar{K}}$ is a section of $\Lambda^{p,q,k}$ then we have

$$(10.55) \quad \#K \cap \{1, \dots, n - \kappa\} = k, \quad \#K \cap \{n - \kappa + 1, \dots, n\} = q - k,$$

$$(10.56) \quad \#K^c \cap \{1, \dots, n - \kappa\} = n - \kappa - k, \quad \#K^c \cap \{n - \kappa + 1, \dots, n\} = \kappa - q + k,$$

from which we get $\mu_K = k - (q - k) - [(n - \kappa - k) - (\kappa - q + k)] = n + 2q - 2\kappa - 4k$. Combining this with (10.48)–(10.50) then gives

$$(10.57) \quad \square_{p;qk} = -\frac{1}{2} \sum_{1 \leq j \leq n} (Z_j \bar{Z}_j + \bar{Z}_j Z_j) - \frac{i}{2} (n + 2q - 2\kappa - 4k) X_0 + \text{lower order terms}.$$

We are now in position to apply Lemma 10.3 to $2\square_{p;qk}$ to get

$$(10.58) \quad \begin{aligned} \nu_0(\square_{p;qk}) &= 2^{-(n+1)} \nu_0(2\square_{p;qk}) = 2^{-(n+1)} (\text{rk } \Lambda^{p;q,k}) \nu(n + 2q - 2\kappa - 4k) \cdot \text{vol}_\theta M \\ &= \frac{1}{2^{n+1}} \binom{n}{p} \binom{n-\kappa}{k} \binom{\kappa}{q-k} \nu(n + 2q - 2\kappa - 4k) \cdot \text{vol}_\theta M. \end{aligned}$$

Combining this with (10.46) and (10.54) then shows that as $\lambda \rightarrow \infty$ we have

$$(10.59) \quad N(\square_{b|\Lambda^{p,q}}; \lambda) \sim \alpha_{n\kappa pq} (\text{vol}_\theta M) \lambda^{n+1},$$

with $\alpha_{n\kappa pq}$ given by (10.45). The proof is thus achieved. $\square$

We now turn to the horizontal sublaplacian on (p, q) -forms.

Theorem 10.5. *Let $\Delta_b : C^\infty(M, \Lambda^{*,*}) \rightarrow C^\infty(M, \Lambda^{*,*})$ be the horizontal sublaplacian associated to the Levi metric on M . Then for $p, q = 1, \dots, n$, with $(p, q) \neq (\kappa, n - \kappa)$ and $(p, q) \neq (n - \kappa, \kappa)$, as $\lambda \rightarrow \infty$ we have*

$$(10.60) \quad N(\Delta_{b|\Lambda^{p,q}}; \lambda) \sim \beta_{n\kappa pq} (\text{vol}_\theta M) \lambda^{n+1},$$

$$(10.61) \quad \beta_{n\kappa pq} = \sum_{\substack{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa) \\ \max(0, p-\kappa) \leq l \leq \min(p, n-\kappa)}} \binom{n-\kappa}{l} \binom{\kappa}{p-l} \binom{n-\kappa}{k} \binom{\kappa}{q-k} \nu(2(q-p) + 4(l-k)).$$

Proof. Thanks to (3.11) we know that we have

$$(10.62) \quad \Delta_b = \square_b + \bar{\square}_b,$$

where $\bar{\square}_b$ denotes the conjugate operator of $\square_b$, that is $\bar{\square}_b \alpha = \overline{\square_b \bar{\alpha}}$, or equivalently the Laplacian of the ∂_b -complex.

As in (10.52) we have orthogonal splittings,

$$(10.63) \quad \Lambda^{p,q} = \bigoplus_{\max(0, p-\kappa) \leq l \leq \min(p, n-\kappa)} \Lambda^{p,l;q} = \bigoplus_{\substack{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa) \\ \max(0, p-\kappa) \leq l \leq \min(p, n-\kappa)}} \Lambda^{p,l;q,k},$$

$$(10.64) \quad \Lambda^{p,l;q} = (\Lambda_+^{1,0})^l \wedge (\Lambda_-^{1,0})^{p-l} \wedge \Lambda^{0,q}, \quad \Lambda^{p,l;q,k} = \Lambda^{p,l;0} \wedge \Lambda^{p,l;0}.$$

Since $\square_b$ is a scalar operator modulo lower order terms, the same is true for $\bar{\square}_b$ and Δ_b . Therefore, on $\Lambda^{p,q}$ we can write

$$(10.65) \quad \bar{\square}_b = \sum_{\max(0, p-\kappa) \leq l \leq \min(p, n-\kappa)} \bar{\square}_{p,l;q} + \text{lower order terms}, \quad \bar{\square}_{p,l;q} = \Pi_{p,l;q} \bar{\square}_b \Pi_{p,l;q},$$

$$(10.66) \quad \Delta_b = \sum_{\substack{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa) \\ \max(0, p-\kappa) \leq l \leq \min(p, n-\kappa)}} \Delta_{p,l;q,k} + \text{lower order terms}, \quad \Delta_{p,l;q,k} = \Pi_{p,l;q,k} \Delta_b \Pi_{p,l;q,k},$$

where $\Pi_{p,l;q}$ and $\Pi_{p,l;q,k}$ denote the orthogonal projections onto $\Lambda^{p,l;q}$ and $\Lambda^{p,l;q,k}$ respectively. In particular, as in (10.67) we have

$$(10.67) \quad \nu_0(\Delta_{b|\Lambda^{p,q}}) = \sum_{\substack{\max(0, q-\kappa) \leq k \leq \min(q, n-\kappa) \\ \max(0, p-\kappa) \leq l \leq \min(p, n-\kappa)}} \nu_0(\Delta_{p,l;q,k}).$$

Next, let $\{X_0, Z_j, \bar{Z}_j\}$ be a local admissible orthonormal frame for $T_{\mathbb{C}}M$. Since $\bar{\square}_{p,l;q} = \overline{\square_{q,p;l}}$, using (10.57) we see that on $\Lambda^{p,l;q}$ we have

$$(10.68) \quad \bar{\square}_{p,l;q} = -\frac{1}{2} \sum_{1 \leq j \leq n} (Z_j \bar{Z}_j + \bar{Z}_j Z_j) + \frac{i}{2} (n + 2p - 2\kappa - 4l) X_0 + \text{lower order terms}.$$

Therefore, on $\Lambda^{p,l;q,k}$ we can write

$$(10.69) \quad \Delta_{p,l;q,k} = \square_{p;q,k} + \bar{\square}_{p,l;q} \\ = \sum_{1 \leq j \leq n} (Z_j \bar{Z}_j + \bar{Z}_j Z_j) - i(2(q-p) + 4(l-k)) X_0 + \text{lower order terms}.$$

Thanks to (10.69) we can apply Lemma 10.3 to get

$$(10.70) \quad \nu_0(\Delta_{p,l;q,k}) = \text{rk } \Lambda^{p,l;q,k} \nu(2(q-p) + 4(l-k)) \text{vol}_{\theta} M \\ = \binom{n-\kappa}{l} \binom{\kappa}{p-l} \binom{n-\kappa}{k} \binom{\kappa}{q-k} \nu(2(q-p) + 4(l-k)) \text{vol}_{\theta} M.$$

Combining this with (10.67) then shows that as $\lambda \rightarrow \infty$ we have

$$(10.71) \quad N(\Delta_b; \lambda) \sim \beta_{npq} (\text{vol}_{\theta} M) \lambda^{n+1},$$

with β_{npq} given by (10.61). The theorem is thus proved. $\square$

Finally, we deal with the conformal powers of the horizontal sublaplacian.

Theorem 10.6. *Assume that M is strictly pseudoconvex (i.e. $\kappa = 0$) and for $k = 1, \dots, n+1$ let $\square_{\theta}^{(k)} : C^{\infty}(M) \rightarrow C^{\infty}(M)$ be a k 'th conformal power of Δ_b acting on functions. Then as $\lambda \rightarrow \infty$ we have*

$$(10.72) \quad N(\square_{\theta}^{(k)}; \lambda) \sim \nu(0) (\text{vol}_{\theta} M) \lambda^{\frac{n+1}{k}}.$$

Proof. Since $\nu_0(\square_{\theta}^{(k)})$ depends only on the principal symbol of $\square_{\theta}^{(k)}$, which is the same as that of Δ_b^k , we have $\nu_0(\square_{\theta}^{(k)}) = \nu_0(\Delta_b^k)$. Therefore, using Theorem 10.5 we see that as $\lambda \rightarrow \infty$ we have

$$(10.73) \quad N(\square_{\theta}^{(k)}; \lambda) \sim N(\Delta_b^k; \lambda) = N(\Delta_b; \lambda)^{\frac{1}{k}} \sim \beta_{n000} (\text{vol}_{\theta} M) \lambda^{\frac{n+1}{k}}.$$

Since $\beta_{n000} = 2^n \nu(0)$ the result follows. $\square$

11. WEYL ASYMPTOTICS AND CONTACT GEOMETRY

In this section we express in more geometric terms the Weyl asymptotics for the horizontal sublaplacian and the contact Laplacian on a compact orientable contact manifold (M^{2n+1}, θ) .

We let $H = \ker \theta$ and let X_0 be the Reeb fields associated to θ , so that $\iota_{X_0} d\theta = 0$ and $\iota_{X_0} \theta = 1$. Since M is orientable H admits a calibrated almost complex structure $J \in C^{\infty}(M, \text{End } H)$, $J^2 = -1$, so that for any nonzero section X of H we have $d\theta(X, JX) = -d\theta(JX, X) > 0$. We then endow M with the orientation defined by θ and the almost complex structure J , so that we have $\theta \wedge d\theta^n > 0$, and with the Riemannian metric,

$$(11.1) \quad g_{\theta} = d\theta(\cdot, J\cdot) + \theta^2.$$

A local orthonormal frame $X_1, \dots, X_{2n}$ of H will be called admissible when we have $X_{n+j} = JX_j$ for $j = 1, \dots, n$. If $\theta^1, \dots, \theta^{2n}$ denotes the dual frame then we have $d\theta = \sum_{j=1}^n \theta^j \wedge \theta^{n+j}$, so the volume form of g_{θ} is equal to

$$(11.2) \quad \theta^1 \wedge \theta^{n+1} \wedge \dots \wedge \theta^n \wedge \theta^{2n} \wedge \theta = \frac{1}{n!} d\theta^n \wedge \theta.$$

In particular, the volume form is independent of the choice of the almost complex structure and depends only on the contact form.

Definition 11.1. The contact volume of (M^{2n+1}, θ) is given by

$$(11.3) \quad \text{vol}_\theta M = \frac{1}{n!} \int_M d\theta^n \wedge \theta.$$

Lemma 11.2. Let $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be selfadjoint sublaplacian such that Δ is bounded from below. We further assume that there exists $\mu \in (-n, n)$ so that near any point of M there is an admissible orthonormal frame $X_1, \dots, X_{2n}$ of H with respect to which Δ takes the form,

$$(11.4) \quad \Delta = -(X_1^2 + \dots + X_{2n}^2) - i\mu X_0 + \text{lower order terms}.$$

Then as $\lambda \rightarrow \infty$ we have

$$(11.5) \quad N(\Delta; \lambda) \sim 2^{-n} \nu(\mu) \text{rk } \mathcal{E}(\text{vol}_\theta M) \lambda^{n+1}.$$

Proof. Let $X_1, \dots, X_{2n}$ be a admissible local orthonormal frame and for $j = 1, \dots, 2n$ let $\tilde{X}_j = \sqrt{2}X_j$. Then $X_0, \tilde{X}_1, \dots, \tilde{X}_{2n}$ is a local H -frame of TM with respect to which Δ takes the form,

$$(11.6) \quad \Delta = -\frac{1}{2}(\tilde{X}_1^2 + \dots + \tilde{X}_{2n}^2) - i\mu(x)X_0 + \text{lower order terms}.$$

Moreover, for $j, k = 1, \dots, 2n$ we have

$$(11.7) \quad \theta([\tilde{X}_j, J\tilde{X}_k]) = -2d\theta(X_j, JX_k) = -2g_\theta(X_j, X_k) = -2\delta_{jk}.$$

Therefore, for $j, k = 1, \dots, n$ we get:

$$(11.8) \quad [X_j, X_{n+k}] = -2\delta_{jk}X_0 \mod H,$$

$$(11.9) \quad [X_0, X_j] = [X_j, X_k] = [X_{n+j}, X_{n+k}] = 0 \mod H.$$

The equalities (11.6)–(11.9) are the same as (10.33)–(10.35) in the case $\kappa = 0$. Therefore, along the same lines as that of the proof Lemma 10.3 we get

$$(11.10) \quad \nu_0(\Delta) = \frac{\nu(\mu)}{n!2^n} \int_M d\theta^n \wedge \theta = 2^{-n} \nu(\mu) \text{vol}_\theta M.$$

Combining this with Theorem 9.1 then proves the asymptotics (11.5). $\square$

Theorem 11.3. Let $\Delta_b : C^\infty(M, \Lambda_{\mathbb{C}}^* H^*) \rightarrow C^\infty(M, \Lambda_{\mathbb{C}}^* H^*)$ be the horizontal sublaplacian on M associated to the metric g_θ above and assume $k \neq n$. Then as $\lambda \rightarrow \infty$ we have

$$(11.11) \quad N(\Delta_{b|_{\Lambda_{\mathbb{C}}^k H^*}}; \lambda) \sim \gamma_{nk}(\text{vol}_\theta M) \lambda^{n+1}, \quad \gamma_{nk} = 2^{-n} \sum_{p+q=k} \binom{n}{p} \binom{n}{q} \nu(p-q).$$

Proof. As explained in Section 3 the almost complex structure of H gives rise to an orthogonal decomposition $\Lambda_{\mathbb{C}}^k H^* = \bigoplus_{p+q=k} \Lambda^{p,q}$. If $X_1, \dots, X_{2n}$ is a local admissible orthonormal frame of H then, as shown by Rumin [Ru, Prop. 2], on $\Lambda^{p,q}$ the operator Δ_b takes the form,

$$(11.12) \quad \Delta_b = -(X_1^2 + \dots + X_{2n}^2) + i(p-q)X_0 + \text{lower order terms}.$$

where the lower order part is not scalar. Therefore, modulo lower order terms, Δ_b preserves the bidegree. We thus may write

$$(11.13) \quad \Delta_{b|_{\Lambda_{\mathbb{C}}^k H^*}} = \sum_{p+q=k} \Delta_{p,q} + \text{lower order terms}, \quad \Delta_{p,q} = \Pi_{p,q} \Delta_b \Pi_{p,q},$$

where $\Pi_{p,q}$ denotes the orthogonal projection of $\Lambda_{\mathbb{C}}^* H^*$ onto $\Lambda^{p,q}$. In particular, we have

$$(11.14) \quad \nu_0(\Delta_{b|_{\Lambda_{\mathbb{C}}^k H^*}}) = \sum_{p+q=k} \nu_0(\Delta_{p,q}).$$

Moreover, since $\Delta_{p,q}$ takes the form (11.12) with respect to any admissible orthonormal frame of H , we may apply Lemma 11.2 to get

$$(11.15) \quad \nu_0(\Delta_{p,q}) = 2^{-n} \sum_{p+q=k} \binom{n}{p} \binom{n}{q} \nu(p-q).$$

Combining this with (11.14) then gives the asymptotics (11.11). $\square$

Finally, in the case of the contact Laplacian we can prove:

Theorem 11.4. *Let $\Delta_R : C^\infty(M, \Lambda^* \oplus \Lambda_*^n) \rightarrow C^\infty(M, \Lambda^* \oplus \Lambda_*^n)$ be the contact Laplacian on M .*

1) *For $k = 0, \dots, 2n$ with $k \neq n$ there exists a universal constant $\nu_{nk} > 0$ depending only on n and k such that as $\lambda \rightarrow \infty$ we have*

$$(11.16) \quad N(\Delta_{R|_{\Lambda^k}}) \sim \nu_{nk} (\text{vol}_\theta M) \lambda^{n+1}.$$

2) *For $j = 1, 2$ there exists a universal constant $\nu_n^{(j)} > 0$ depending only on n and j such that as $\lambda \rightarrow \infty$ we have*

$$(11.17) \quad N(\Delta_{R|_{\Lambda_*^n}}) \sim \nu_n^{(j)} (\text{vol}_\theta M) \lambda^{\frac{n+1}{2}}.$$

Proof. Let $a \in M$ and consider a chart around a together with an admissible orthonormal frame $X_1, \dots, X_{2n}$ of H . Since $X_0, X_1, \dots, X_{2n}$ form a H -frame this chart is a Heisenberg chart. Moreover, as shown in the proofs of Lemma 10.3 and Lemma 11.2 the following hold:

(i) We have X_j^a , where $X_0^0, \dots, X_{2n}^0$ are the left-invariant vector fields (10.18) on $\mathbb{H}^{2n+1}$. In particular, we have $G_a M = \mathbb{H}^{2n+1}$ and $H_a = H_0^0$, where H_0^0 denotes the left-invariant Heisenberg hyperplane bundle of $\mathbb{H}^{2n+1}$.

(ii) We have $\theta(0) = dx_0 = \theta^0(0)$ and $d\theta(0) = 2 \sum_{j=1}^n dx_j \wedge dx_{n+j} = d\theta^0(0)$, where $\theta^0 = dx_0 + \sum_{j=1}^n (x_j dx_{n+j} - x_{n+j} dx_j)$ is the standard left-invariant contact form of $\mathbb{H}^{2n+1}$.

(iii) The density on M given by the contact volume form $\frac{1}{n!} \theta \wedge d\theta^n$ agrees at $x = 0$ with the density dx on $\mathbb{R}^{d+1}$.

On the other hand, for $k = 0, 1, \dots, 2n$ the fiber at a of the bundle Λ_*^k depends only on H_a and on the values of θ and $d\theta$ at a . Therefore, it follows from the statements (i) and (ii) that in the Heisenberg coordinates centered at a the fibers at $x = 0$ of the bundles $\Lambda^* \oplus \Lambda_*^n$ of M and $\mathbb{H}^{2n+1}$ agree.

Next, let Δ_R^0 denote the contact Laplacian on $\mathcal{H}^{2n+1}$. Then we have:

Lemma 11.5. *In the Heisenberg coordinates centered at a the model operator Δ_R^a agrees with Δ_R^0 .*

Proof of the lemma. First, note that in view of the formulas (3.23)–(3.27) for Δ_R and of Proposition 4.29 and Proposition 4.30, we only have to show that in the Heisenberg coordinates centered at a the model operators d_R^a and D_R^a agree with the operators d_R^0 and D_R^0 on $\mathbb{H}^{2n+1}$.

Let $\theta^1, \dots, \theta^{2n}$ be the coframe of H^* dual to $X_1, \dots, X_{2n}$. This coframe gives rise to a trivialization of $\Lambda_{\mathbb{C}}^* H^*$ in the chart with respect to which we have $d_b = \sum_{j=1}^{2n} \varepsilon(\theta^j) X_j$. Furthermore, since $X_j(0) = \frac{d}{dx_j}$ we have $\theta_j(0) = dx_j$, so the model operator of d_b is $d_b^a = \sum_{j=1}^{2n} \varepsilon(dx_j) X_j^0 = d_b^0$, where d_b^0 is the d_b -operator on $\mathbb{H}^{2n+1}$. In particular, as for $k = n+1, \dots, 2n$ we have $d_R = d_b$ on Λ^k , we get $d_R^a = d_b^a = d_b^0 = d_R^0$.

On the other hand, as shown in [Ru] for $k = 0, \dots, n-1$ we have $d_R^* = d_b^*$ on Λ^k . Thus $(d_R^a)^* = (d_R^*)^a = (d_b^*)^a = (d_b^a)^* = (d_b^0)^* = (d_R^0)^*$, which by taking adjoints gives $d_R^a = d_R^0$.

Finally, it is proved in [Ru] that $D_R = X_0 + d_b \varepsilon(d\theta)^* d_b$ on Λ_1^n . Therefore, we get $D_R^a = X_0^a + d_b^a (\varepsilon(d\theta)^a)^* d_b^a = X_0^0 + d_b^0 \varepsilon(d\theta^0)^* d_b^0 = D_R^0$. The proof that $\Delta_R^a = \Delta_R^0$ is thus complete. $\square$

Thanks to the statements (i) and (iii) and the claim above we may apply Proposition 9.4. Letting $K_0(x, t)$ be the fundamental solution of $\Delta_R^0 + \partial_t$ we then obtain:

- For $k = 0, \dots, 2n$, with $k \neq n$, as $\lambda \rightarrow \infty$ we have

$$(11.18) \quad N(\Delta_{R|_{\Lambda^k}}) \sim \nu_{nk}(\text{vol}_\theta M) \lambda^{n+1}, \quad \nu_{nk} = \frac{2^{-n}}{(n+1)!} \text{tr}_{\Lambda^{k,0}} K_{0|_{\Lambda^{k,0}}}(0, 1).$$

- For $j = 1, 2$ as $\lambda \rightarrow \infty$, we have

$$(11.19) \quad N(\Delta_{R|_{\Lambda_j^n}}) \sim \nu_{n,j}(\text{vol}_\theta M) \lambda^{\frac{n+1}{2}}, \quad \nu_n^{(j)} = 2^{-n} \Gamma(1 + \frac{n+1}{2})^{-1} \text{tr}_{\Lambda_j^{n,0}} K_{0|_{\Lambda_j^{n,0}}}(0, 1).$$

In particular, the constant ν_{nk} (resp. $\nu_n^{(j)}$) depends only on n and k (resp. n and j), hence is a universal constant. Furthermore, Theorem 9.1 implies that ν_{nk} and $\nu_n^{(j)}$ are positive numbers. $\square$

REFERENCES

- [AS] Atiyah, M.F.; Singer, I.M.: *The index of elliptic operators. I*. Ann. of Math. (2) **87** (1968), 484–530.
- [Be] Bellaïche, A.: *The tangent space in sub-Riemannian geometry*. Sub-Riemannian geometry, 1–78, Progr. Math., 144, Birkhäuser, Basel, 1996.
- [Bea] Beals, R.: *Opérateurs invariants hypoelliptiques sur un groupe de Lie nilpotent*. Séminaire Goulaouic-Schwartz (1976–77), Exposé No. 19. Ecole Polytech., Palaiseau, 1977.
- [BG] Beals, R.; Greiner, P.C.: *Calculus on Heisenberg manifolds*. Annals of Mathematics Studies, vol. 119. Princeton University Press, Princeton, NJ, 1988.
- [BGS] Beals, R.; Greiner, P.C.; Stanton, N.K.: *The heat equation on a CR manifold*. J. Differential Geom. **20** (1984), no. 2, 343–387.
- [BS1] Beals, R.; Stanton, N.K.: *The heat equation for the $\bar{\partial}$ -Neumann problem. I*. Comm. Partial Differential Equations **12** (1987), no. 4, 351–413.
- [BS2] Beals, R.; Stanton, N.K.: *The heat equation for the $\bar{\partial}$ -Neumann problem. II*. Canad. J. Math. **40** (1988), no. 2, 502–512.
- [Bo] Boutet de Monvel, L.: *Hypoelliptic operators with double characteristics and related pseudo-differential operators*. Comm. Pure Appl. Math. **27** (1974), 585–639.
- [BGH] Boutet de Monvel, L.; Grigis, A.; Helffer, B.: *Parametrixes d’opérateurs pseudo-différentiels à caractéristiques multiples*. Journées EDP (Rennes, 1975) pp. 93–121. Astérisque, No. 34–35, Soc. Math. France, Paris, 1976.
- [CGGP] Christ, M.; Geller, D.; Głowacki, P.; Polin, L.: *Pseudodifferential operators on groups with dilations*. Duke Math. J. **68** (1992) 31–65.
- [Co] Connes, A.: *Noncommutative geometry*. Academic Press, Inc., San Diego, CA, 1994.
- [CM] Connes, A.; Moscovici, H.: *The local index formula in noncommutative geometry*. Geom. Funct. Anal. **5** (1995), no. 2, 174–243.
- [Dy1] Dynin, A.: *Pseudodifferential operators on the Heisenberg group*. Dokl. Akad. Nauk SSSR **225** (1975) 1245–1248.
- [Dy2] Dynin, A.: *An algebra of pseudodifferential operators on the Heisenberg groups*. Symbolic calculus. Dokl. Akad. Nauk SSSR **227** (1976), 792–795.
- [EMM] Epstein, C.L.; Melrose, R.B.; Mendoza, G.: *The Heisenberg algebra, index theory and homology*. To appear. Partially available at <http://www-math.mit.edu/~rbm>.
- [ET] Eliashberg, Y.; Thurston, W.: *Confoliations*. University Lecture Series, 13, AMS, Providence, RI, 1998.
- [FS1] Folland, G.; Stein, E.: *Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group*. Comm. Pure Appl. Math. **27** (1974) 429–522.
- [FS2] Folland, G.; Stein, E.: *Hardy spaces on homogeneous groups*. Mathematical Notes, 28. Princeton University Press, Princeton, 1982.
- [GJ] Glimm, J.; Jaffe, A.: *Quantum physics. A functional integral point of view*. Springer-Verlag, 1987.
- [Gl] Głowacki, P.: *Stable semigroups of measures as commutative approximate identities on nongraded homogeneous groups*. Invent. Math. **83** (1986), no. 3, 557–582.
- [GK] Gohberg, I.C.; Kreĭn, M.G.: *Introduction to the theory of linear nonselfadjoint operators*. Trans. of Math. Monographs 18, AMS, Providence, 1969.
- [GG] Gover, A.R.; Graham, C.R.: *CR invariant powers of the sub-Laplacian*. E-print [arXiv:math.DG/0301092](https://arxiv.org/abs/math.DG/0301092), 31 pages.

- [Gr] Greenleaf, A.: *The first eigenvalue of a sub-Laplacian on a pseudo-Hermitian manifold*. Comm. Partial Differential Equations **10** (1985), no. 2, 191–217.
- [Gre] Greiner, P.: *An asymptotic expansion for the heat equation*. Arch. Rational Mech. Anal. **41** (1971) 163–218.
- [Gri] Grigis, A.: *Hypoellipticité et paramétrix pour des opérateurs pseudodifférentiels à caractéristiques doubles*. Journées EDP (Rennes, 1975) pp. 183–205. Astérisque, No. 34-35, Soc. Math. France, Paris, 1976.
- [Gro] Gromov, M.: *Carnot-Carathéodory spaces seen from within. Sub-Riemannian geometry*, 79–323, Progr. Math., 144, Birkhäuser, Basel, 1996.
- [Gru] Grubb, G.: *Functional calculus of pseudodifferential boundary problems*. Second edition. Progress in Mathematics, 65. Birkhäuser, Boston, MA, 1996.
- [Gu1] V.W. Guillemin *A new proof of Weyl's formula on the asymptotic distribution of eigenvalues*. Adv. in Math. **55** (1985), no. 2, 131–160.
- [Gu2] Guillemin, V.W.: *Gauged Lagrangian distributions*. Adv. Math. **102** (1993), no. 2, 184–201.
- [Ha] Hardy, G.H.: *Divergent Series*. Oxford, at the Clarendon Press, 1949.
- [HN1] Helffer, B.; Nourrigat, F.: *Hypoellipticité pour des groupes nilpotents de rang de nilpotence 3*. Comm. Partial Differential Equations **3** (1978), no. 8, 643–743.
- [HN2] Helffer, B.; Nourrigat, J.: *Caractérisation des opérateurs hypoelliptiques homogènes invariants à gauche sur un groupe de Lie nilpotent gradué*. Comm. Partial Differential Equations **4** (1979), no. 8, 899–958.
- [HN3] Helffer, B.; Nourrigat, J.: *Hypoellipticité maximale pour des opérateurs polynômes de champs de vecteurs*. Prog. Math., No. 58, Birkhäuser, Boston, 1986.
- [HS] Hilsaum, M.; Skandalis, G.: *Morphismes K -orientés d'espaces de feuilles et fonctorialité en théorie de Kasparov (d'après une conjecture d'A. Connes)*. Ann. Sci. École Norm. Sup. (4) **20** (1987), no. 3, 325–390.
- [Hö1] L. Hörmander *Pseudo-differential operators and hypoelliptic equations. Singular integrals* (Proc. Sympos. Pure Math., Vol. X, Chicago, Ill., 1966), pp. 138–183. Amer. Math. Soc., Providence, R.I., 1967.
- [Hö2] L. Hörmander *Hypoelliptic second order differential equations*. Acta Math. **119** (1967), 147–171.
- [Hö3] Hörmander, L.: *A class of hypoelliptic pseudodifferential operators with double characteristics*. Math. Ann. **217** (1975), no. 2, 165–188.
- [Hw] I.L. Hwang *The L^2 -boundedness of pseudodifferential operators*. Trans. Amer. Math. Soc. **302** (1987), no. 1, 55–76.
- [II] Iwasaki, C.; Iwasaki, N.: *Parametrix for a degenerate parabolic equation and its application to the asymptotic behavior of spectral functions for stationary problems*. Publ. Res. Inst. Math. Sci. **17** (1981), no. 2, 577–655.
- [JL1] Jerison, D.; Lee, J.M.: *The Yamabe problem on CR manifolds*. J. Differential Geom. **25** (1987), no. 2, 167–197.
- [JL2] Jerison, D.; Lee, J.M.: *Extremals for the Sobolev inequality on the Heisenberg group and the CR Yamabe problem*. J. Amer. Math. Soc. **1** (1988), no. 1, 1–13.
- [JL3] Jerison, D.; Lee, J.M.: *Intrinsic CR normal coordinates and the CR Yamabe problem*. J. Differential Geom. **29** (1989), no. 2, 303–343.
- [JK] Julg, P.; Kasparov, G.: *Operator K -theory for the group $SU(n, 1)$* . J. Reine Angew. Math. **463** (1995), 99–152.
- [Ko] Koenig, K.: *On maximal Sobolev and Hölder estimates for the tangential Cauchy-Riemann operator and boundary Laplacian*. Amer. J. Math. **124** (2002), no. 1, 129–197.
- [Koh1] J. J. Kohn *Boundaries of complex manifolds*. 1965 Proc. Conf. Complex Analysis (Minneapolis, 1964) pp. 81–94 Springer, Berlin
- [Koh2] Kohn, J. J.: *Superlogarithmic estimates on pseudoconvex domains and CR manifolds*. Ann. of Math. (2) **156** (2002), no. 1, 213–248.
- [KR] Kohn, J.J.; Rossi, H.: *On the extension of holomorphic functions from the boundary of a complex manifold*. Ann. of Math. **81** (1965) 451–472.
- [KV] M. Kontsevich, S. Vishik *Geometry of determinants of elliptic operators. Functional analysis on the eve of the 21st century*, Vol. 1 (New Brunswick, NJ, 1993), 173–197, Progr. Math., 131, Birkhäuser Boston, Boston, MA, 1995.
- [Kr1] Krainer, T.: *Parabolic pseudodifferential operators and long-time asymptotics of solutions*. PhD dissertation, University of Potsdam, 2000.
- [Kr2] Krainer, T.: *Volterra families of pseudodifferential operators and The calculus of Volterra Mellin pseudo-differential operators with operator-valued symbols. Parabolicity, Volterra calculus, and conical singularities*, pp. 1–45 and 47–91. Oper. Theory Adv. Appl., 138, Birkhäuser, Basel, 2002.
- [KS] Krainer, T.; Schulze, B.W.: *On the inverse of parabolic systems of partial differential equations of general form in an infinite space-time cylinder. In Parabolicity, Volterra calculus, and conical singularities*, pp. 93–278. Oper. Theory Adv. Appl., 138, Birkhäuser, Basel, 2002.
- [Le] Lee, J.M.: *The Fefferman metric and pseudo-Hermitian invariants*. Trans. Amer. Math. Soc. **296** (1986), no. 1, 411–429.

- [MMS] Mathai, V.; Melrose, R.B.; Singer, I.M: *Fractional analytic index*. E-print, ArXiv, Feb. 04.
- [MS] Menikoff, A.; Sjöstrand, J.: *On the eigenvalues of a class of hypoelliptic operators. II*. Global analysis (Proc. Biennial Sem. Canad. Math. Congr., Univ. Calgary, Calgary, Alta., 1978), pp. 201–247, Lecture Notes in Math., 755, Springer, Berlin, 1979.
- [Me1] Melrose, R.B.: *Hypoelliptic operators with characteristic variety of codimension two and the wave equation*. Séminaire Goulaouic-Schwartz, 1979–1980 (French), Exp. No. 11, 13 pp., Ecole Polytech., Palaiseau, 1980.
- [Me2] Melrose, R.B.: *The Atiyah-Patodi-Singer index theorem*. A.K. Peters, Boston, 1993.
- [Me3] Melrose, R.B.: *Star products and local line bundles*. Preprint, 2004.
- [Mo1] Mohamed, A.: *Etude spectrale d'opérateurs hypoelliptiques à caractéristiques multiples. I*. Ann. Inst. Fourier (Grenoble) **32** (1982), no. 3, ix, 39–90.
- [Mo2] Mohamed, A.: *Etude spectrale d'opérateurs hypoelliptiques à caractéristiques multiples. II*. Comm. Partial Differential Equations **8** (1983), no. 3, 247–316.
- [Ni] Nicoara, A.: *Regularity results for the tangential Cauchy-Riemann operator and the Szegő projection on weakly pseudoconvex CR manifolds*. PhD dissertation, Princeton University, 2002.
- [Pi] Piriou, A.: *Une classe d'opérateurs pseudo-différentiels du type de Volterra*. Ann. Inst. Fourier **20** (1970) 77–94.
- [Po1] Ponge, R.: *Calcul hypoelliptique sur les variétés de Heisenberg, résidu non commutatif et géométrie pseudo-hermitienne*. PhD Thesis, Univ. Paris-Sud (Orsay), Dec. 2000.
- [Po2] Ponge, R.: *Calcul fonctionnel sous-elliptique et résidu non commutatif sur les variétés de Heisenberg*. C. R. Acad. Sci. Paris, Série I, **332** (2001) 611–614.
- [Po3] Ponge, R.: *Géométrie spectrale et formules d'indices locales pour les variétés CR et contact*. C. R. Acad. Sci. Paris, Série I, **332** (2001) 735–738.
- [Po4] Ponge, R.: *A new short proof of the local index formula and some of its applications*. Comm. Math. Phys. **241** (2003) 215–234. Erratum Comm. Math. Phys. **248** (2004) 639.
- [Po5] Ponge, R.: *On the asymptotic completeness of the Volterra calculus*. Appendix by H. Mikaleyan and R. Ponge. E-print, arXiv, Oct. 03, 14 pages. To appear in J. Anal. Math..
- [Po6] Ponge, R.: *The tangent groupoid of a Heisenberg manifold*. E-print, arXiv, Apr. 04, 17 pages.
- [Po7] Ponge, R.: *A new proof of the regularity of the eta invariant for Dirac operators*. E-print, arXiv, Nov. 04, 10 pages.
- [Po8] Ponge, R.: *Intrinsic notion of principal symbol for the Heisenberg calculus*. E-print, arXiv, April 05.
- [Po9] Ponge, R.: *Remarks on Julg-Kasparov's proof of the Baum-Connes conjecture for the group $SU(n, 1)$* . E-print, arXiv, April 05.
- [Po10] Ponge, R.: *Functional calculus and spectral asymptotics for hypoelliptic operators on Heisenberg manifolds. II*. E-print, arXiv, May 05.
- [Po11] Ponge, R.: *Noncommutative geometry, Heisenberg calculus and CR geometry*. E-print, arXiv, May 05.
- [Po12] Ponge, R.: *New approaches to the construction of complex powers of elliptic ΨDO 's*. E-print, arXiv, May 05.
- [Ro1] Rockland, C.: *Hypoellipticity on the Heisenberg group-representation-theoretic criteria*. Trans. Amer. Math. Soc. **240** (1978) 1–52.
- [Ro2] Rockland, C.: *Intrinsic nilpotent approximation*. Acta Appl. Math. **8** (1987), no. 3, 213–270.
- [RSt] Rothschild, L.; Stein, E.: *Hypoelliptic differential operators and nilpotent groups*. Acta Math. **137** (1976) 247–320.
- [Ru] Rumin, M.: *Formes différentielles sur les variétés de contact*. J. Differential Geom. **39** (1994), no.2, 281–330.
- [Se] R.T. Seeley *Complex powers of an elliptic operator. Singular integrals* (Proc. Sympos. Pure Math., Vol. X, Chicago, Ill., 1966), pp. 288–307. Amer. Math. Soc., Providence, R.I., 1967.
- [Sh] M. Shubin *Pseudodifferential operators and spectral theory*. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin-New York, 1987.
- [Ta] Tanaka, N.: *A differential geometric study on strongly pseudo-convex manifolds*. Lectures in Mathematics, Department of Mathematics, Kyoto University, No. 9. Kinokuniya Book-Store Co., Ltd., Tokyo, 1975.
- [Tay] Taylor, M.E.: *Noncommutative microlocal analysis. I*. Mem. Amer. Math. Soc. **52** (1984), no. 313,
- [Va] Van Erp, E.: PhD dissertation, Penn State University, June 2004.
- [We] Webster, S.: *Pseudo-Hermitian structures on a real hypersurface*. J. Differential Geom. **13** (1978), no. 1, 25–41.
- [Wo1] Wodzicki, M.: *Local invariants of spectral asymmetry*. Invent. Math. **75** (1984) 143–177.
- [Wo2] Wodzicki, M.: *Noncommutative residue. I. Fundamentals. K-theory, arithmetic and geometry* (Moscow, 1984–1986), 320–399, Lecture Notes in Math., 1289, Springer, 1987.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, USA.
 E-mail address: `ponge@math.ohio-state.edu`